

QUASI-ALGEBRAIC GEOMETRY OF CURVES I RIEMANN–ROCH THEOREM AND JACOBIAN

ILYA ZAKHAREVICH

Dedicated to Yu. I. Manin at his 60th birthday, to a Teacher from a pupil

ABSTRACT. We discuss an analogue of Riemann–Roch theorem for curves with an infinite number of handles. We represent such a curve X by its Shottki model, which is an open subset U of $\mathbb{CP}^1$ with infinite union of circles as a boundary. An *appropriate* bundle on X is $\omega^{1/2} \otimes \mathcal{L}$, $\mathcal{L}$ being a bundle with (say) constants as gluing conditions on the circles. An *admissible* section of an appropriate bundle on X is a holomorphic half-form on U with given gluing conditions and $H^{1/2}$ -smoothness condition.

We study the restrictions on the mutual position of the circles and the gluing constants which guarantee the finite dimension of the space of appropriate sections of admissible bundles, and make the Riemann–Roch theorem hold. The resulting Jacobian variety is described as an infinite-dimension analogue of a torus.

CONTENTS

0. Introduction	3
0.1. Integrable systems and algebro-geometric solutions	4
0.2. Universal Grassmannian	5
0.3. Semistable elliptic curves	8
0.4. Compactified moduli spaces	11
0.5. Adjacency of moduli spaces and Shottki model	13
0.6. Second compactification	15
0.7. Growth conditions and divisors	17
0.8. Riemann–Roch and growth control	17
0.9. Handles, Sobolev spaces, and representations of $\mathrm{SL}_2(\mathbb{C})$	18
0.10. Thick infinity	20
0.11. Fight with $H^{1/2}$	21
0.12. Main results	21
0.13. Main tools	22
0.14. Historic remarks	23
1. Geometry of half-forms	24
1.1. Half-forms and holomorphic half-forms	24
1.2. Half-forms on complex curves	24

Date: August 1997 *Printed:* February 5, 2008.

1.3. Restriction onto S^1	25
2. Sobolev spaces	25
2.1. Euclidean case	26
2.2. Sobolev spaces on manifolds	27
2.3. Capacity	28
2.4. Generalized Sobolev spaces	28
2.5. Rescaling on $\mathbb{R}^n$ and $\mathbb{C}^n$	29
2.6. $\mathrm{PGL}(2, \mathbb{C})$ -invariant realization of $H^{1/2}(\mathbb{CP}^1)$.	30
2.7. Hardy space	32
3. Cauchy kernel for $\omega^{1/2}$	33
3.1. $\bar{\partial}$ on piecewise-analytic functions	33
3.2. Self-duality of $\bar{\partial}$	34
3.3. The kernel of $\bar{\partial}^{-1}$	35
3.4. Cauchy kernel and L_2	36
4. Toy theory	36
4.1. Toy global space	36
4.2. Almost-perpendicularity	37
4.3. Topology on the boundary	38
4.4. Space of holomorphic functions	39
4.5. Toy Riemann–Roch theorem	40
5. $H^{1/2}$ -theory	42
5.1. In an ideal world	42
5.2. Mollification	43
5.3. Mollification suitable for $\bar{\partial}$	44
5.4. $\tilde{e} \circ \tilde{r}$ as an approximation to $\bar{\partial}$	45
5.5. Mollification for $L_2(\Omega^{1/2} \otimes \mu)$	47
5.6. Almost perpendicular subspaces	47
5.7. Conformal distance	48
5.8. Subspaces of $\Omega^{3/4}$	49
6. Generalized Hardy space	50
6.1. Hilbert space of holomorphic half-forms	50
6.2. Hilbert operator	51
6.3. Boundary map	52
6.4. Gluing conditions	55
6.5. Strong sections and duality	55
6.6. Weak sections	58
6.7. Finite-degree bundles	59
6.8. Stratification of infinity	59
7. Riemann–Roch theorems	62
7.1. Abstract Riemann–Roch theorem	62
7.2. Riemann problem	63
7.3. Finiteness theorem	65

7.4.	Duality	66
7.5.	Riemann–Roch for curves	67
7.6.	Criterion of admissibility of a curve	69
7.7.	Filling the gap	70
7.8.	Moduli space	70
8.	Set of admissible bundles	72
8.1.	Exceptional indices	73
8.2.	Hilbert–Schmidt bundles	73
8.3.	Involution	73
8.4.	Hilbert–Schmidt criterion	73
8.5.	log-convexity	75
8.6.	Types of admissible bundles	75
8.7.	Multiplicators, equivalence and Jacobians	76
8.8.	Divisors	79
8.9.	Universal Grassmannian and bundles	80
9.	Structure of Jacobian	82
9.1.	Constant Jacobian	82
9.2.	The partial period mapping	85
9.3.	A -Periods	87
9.4.	B -periods	89
9.5.	Space of periods	91
9.6.	Period matrix	92
9.7.	Bounded Jacobian as a torus	95
9.8.	Rigged Hodge structure	97
	References	100

0. INTRODUCTION

The need to carve out a set of curves of infinite genus for which “most” theorems of algebraic geometry are true comes out from the following observations:

1. The existence of algebro-geometric description of solutions of infinite-dimensional integrable systems;
2. The ability to describe the series of perturbation theory for string amplitudes as integrals over moduli spaces;
3. The hope that the union of compactifications of moduli spaces may have a simpler geometry than the moduli spaces themselves.

Different approaches which would result in different sets of curves are possible (as in papers of Feldman, Knörrer and Trubowitz cf. [3]), thus we first motivate our choice of tools (Shottki model, $H^{1/2}$ -topology, capacities and half-forms) as (probably) a best one to fulfill the expectations of the above origins of the theory. Until Section 0.7 we discuss how the above topics motivate the choice of the Shottki model as a way to

describe a curve of infinite genus. After this (up to section 0.10) we describe motivations for the choice of half-forms, $H^{1/2}$ -topology, and generalized-Sobolev-spaces (or capacities) to describe admissible sections on a given curve. In the remaining part of the introduction we do a walk-through the methods and results one can find in this paper, as well as some historic remarks.

0.1. Integrable systems and algebro-geometric solutions. A great break-through in the first topic came with the work [7]. A hyperelliptic curve of genus g with real branching points and a divisor of degree g on this curve allows one to construct a so-called *algebro-geometric* solution of KdV equation

$$u_{xxx} + 12uu_x - 4u_t = 0.$$

Such solutions are called *g-gap potentials*. McKean and Trubowitz studied what substitutes algebraic geometry for solutions of KdV which are *not* finite gap potentials. To such a solution they associated a curve (i.e., a complex manifold of dimension 1) which was hyperelliptic of *infinite genus*, i.e., had infinitely many branching points, and some substitution for the notion of a divisor of degree g . It was shown that for the curves which are related to KdV equation one can construct a well-behaved analogue of 1-dimensional algebraic geometry. One should consider this analogue as a generalization of algebraic geometry to hyperelliptic curves of infinite genus (in fact only to some special curves of this type).

Note that other integrable systems lead to different classes of curves which appear in algebro-geometric solutions for the systems. Thus one may expect that infinite-dimensional integrable systems may lead to generalizations of algebraic geometry to different classes of curves of infinite genus.

For this approach one of the richest systems is so called KP system

$$(u_{xxx} + 12uu_x - 4u_t)_x + 3u_{yy} = 0.$$

To describe an algebro-geometric solution of a KP system one starts with an arbitrary algebraic curve, and an arbitrary linear bundle on this curve. Thus to generalize the approach of [7] to the KP equation, one needs to study curves of infinite genus of *generic* form (as opposed to a hyperelliptic curve) and bundles on them.

Suppose for a minute that we have such a generalization, i.e., a collection of curves and bundles on them. Call the members of these families *admissible*. We assume that to any admissible curve and a bundle on it we can associate some solution of KP equation. Let us investigate what can we deduce about this collection from the known properties of KP system (the properties below are applicable at least to algebro-geometric solutions).

The dynamics of KP leaves the curve the same, but changes the bundle. Thus the collection of admissible bundles should be reach enough to include all the bundles obtained by time-flow of KP. In fact KP can be generalized to include an infinite collection of commuting flows (with different time variables), and they (taken together) can transform any bundle to any other one (at least in finite-genus case). Thus we

should expect that we need our collection to include *all* the possible bundles, thus the whole Jacobian.

The dynamics of KP is described in terms of locally affine structure on the Jacobian, hence one should be able to describe the Jacobian as a quotient of a vector space by a lattice. Algebraic geometry identifies the Jacobian with a quotient of the space of global holomorphic forms by the forms with integer periods. Thus one needs something similar to this description.

There is an alternative description of a solution of KP equation in terms of so called τ -functions. The relation of algebro-geometric description with the description in terms of τ -functions needs a construction of a Laurent series of a meromorphic global section of an admissible bundle. This ceases to be trivial in infinite-genus case, since already the description in [7] shows that one may need to apply this operation at the *infinity* of the curve, i.e., at the points added to the curve to compactify it. These points *are not* smooth points of the curve, the curve is not even a topological manifold near these points.

Finally, to get somewhat *explicit* description of solutions of KP as functions of several variables, one needs a way to explicitly describe global sections of linear bundles. Moreover, the bundles we need to consider should have a finite number (preferably one!) of independent global sections. A tool to construct such bundles in the case of finite genus is the Riemann–Roch theorem, which gives an estimate on the dimension of sections of the given bundles, and this estimate is precise in the case of bundles in generic position. These sections are described in terms of θ -functions, thus we will also need to describe θ -functions.

Collecting all this together, we see that we need to describe curves X which can be equipped with a linear bundle $\mathcal{L}$ such that the space of global sections is finite-dimensional. One should be able to define what does it mean that two bundles $\mathcal{L}$ and $\mathcal{L}'$ are equivalent, describe the equivalence classes of bundles in terms of global holomorphic 1-forms, give a local description of sections of the bundles, and give a global description of sections of the bundles in terms of θ -functions.

In this paper we do not complete this program. However, we describe all the ingredients but the last one.

0.2. Universal Grassmannian. The other two origins, the string theory and geometry of moduli spaces, come into play if we consider the algebro-geometric methods of solving integrable systems in the other direction, as a way to find information of algebro-geometric type from solutions of integrable systems (similar to solution of Shottki problem in [8]).

The key idea is that in the investigated cases the set of algebro-geometric solutions is *dense* in the set of all solutions, thus the set of the solutions is a *completion* of the set of algebro-geometric solutions. Since the the set of algebro-geometric solutions is a moduli space of appropriate structures, and the set of solutions is a linear space

(due to the possibility to solve the Cauchy problem), we see that the moduli space has a completion which has a topology of a linear space.

While the moduli spaces carry remarkable measures [2] used in the integrals of the string theory, this linear space has a symplectic structure, so one may expect that this symplectic structure may have a relationship to the measure on the subsets corresponding to algebro-geometric solutions.

To carry out this program one needs to investigate how an arbitrary solution may be approximated by algebro-geometric solutions. For KdV equation this question was answered by [7] (for a fixed spectrum), ameliorated to describe the inclusion of the subset of finite gap potentials up to a diffeomorphism in [9, 10], up to a symplectomorphism in [1]. The answer is that the above inclusion is isomorphic to inclusion of trigonometric polynomials into the space of all functions (for a very wide range of spaces of functions). Since trigonometric polynomials fill finite-dimensional coordinate subspaces in spaces of functions (using Fourier coefficients as coordinates), we see that this picture is very similar to one of a divisor with normal crossings.

Much simpler problem is how an algebro-geometric solution may be approximated by (simpler) algebro-geometric solutions. A remarkable fact is that such problems for different classes of integrable systems may be solved via a uniform approach: compactification of moduli spaces via *Universal Grassmannian*.

The reason is that the standard way to associate a solution of an integrable system to an algebro-geometric data comes from consideration of the mapping to *Universal Grassmannian*. The Universal Grassmannian gives a convenient way to say that the space of sections of a linear bundle on one curve is close to the space of sections on another linear bundle on another curve.

Consider an algebraic curve X with a fixed point $P \in X$ and a local coordinate system x in a neighborhood of P which maps P to $0 \in \mathbb{C}$. Let $\mathcal{L}$ be a linear bundle on X , fix a trivialization of this bundle in a neighborhood of P . One can associate a Laurent series $\mathfrak{l}(\varphi)$ to any section φ of $\mathcal{L}$ in a punctured neighborhood of P : using a coordinate system on X and the trivialization of $\mathcal{L}$, one can write this section as a function of $z \approx 0$, $z \neq 0$. Let V be the space of meromorphic sections of $\mathcal{L}$ which are holomorphic outside of P . (Abusing divisor notations, one can write V as $\Gamma(X, \mathcal{L}(\infty \cdot P)) = \bigcup_k \Gamma(X, \mathcal{L}(k \cdot P))$.) Then $\mathfrak{l}(V)$ is a subspace of the space $\mathfrak{L}$ of Laurent series. Let the universal Grassmannian $\text{Gr}(\mathfrak{L})$ be the Grassmannian of subspaces of $\mathfrak{L}$. Then $\mathfrak{l}(V) \in \text{Gr}(\mathfrak{L})$ depends on X , $\mathcal{L}$, the coordinate system near P , and on the trivialization of $\mathcal{L}$ near P .

On $\mathfrak{L}$ there are natural actions of the group of formal diffeomorphisms¹ of $(\mathbb{C}, 0)$ and the group of multiplication by invertible Taylor series. Let G be the semidirect product of these groups. The action of G on $\mathfrak{L}$ corresponds (via $\mathfrak{l}$) to changes of coordinate system on X , and a change of the trivialization of $\mathcal{L}$. We see that to a triple $(X, P, \mathcal{L})$ we can naturally associate a point $\mathfrak{l}(X, P, \mathcal{L}) \in \text{Gr}(\mathfrak{L})/G$.

¹I.e., invertible ∞ -jets of mappings $\mathbb{C} \rightarrow \mathbb{C}$ with $0 \in \mathbb{C}$ being a fixed point.

Note that while only the mapping to $\mathrm{Gr}(\mathfrak{L})/G$ is invariantly defined, one can get a canonical lifting to $\mathrm{Gr}(\mathfrak{L})$ as far as $X \neq \mathbb{CP}^1$. The description below does not behave well w.r.t. deformations of the curve/bundle, but can be easily modified to do so. Given a bundle $\mathcal{L}$, we can find $k \in \mathbb{Z}$ such that $\mathcal{L}(k \cdot P)$ has only one independent section, and $\mathcal{L}((k+1) \cdot P)$ has another one. The ratio z of these two sections identifies a neighborhood of P with a neighborhood of $\infty \in \mathbb{CP}^1$, and this identification is defined up to an affine transformation. Now the only section of $\mathcal{L}(k \cdot P)$ gives a local identification of $\mathcal{L}$ with $\mathcal{O}(-k \cdot P)$. Together with a coordinate system in a neighborhood of P it gives a local section of $\mathcal{L}$ defined up to multiplication by a constant.

What remains is to pick up a coordinate system from those which differ by an affine transformation. If $X \neq \mathbb{CP}^1$, then for some $l > k+1$ there is a section of $\mathcal{L}(l \cdot P)$ of the form $z^l + p_{l-1}(z) + az^{-m} + bz^{-m-1} + O(z^{-m-2})$ with $a \neq 0$, p_{l-1} being a polynomial of degree $l-1$, and $m > 0$. Taking the minimal possible l , we see that the condition that $a = 1$, $b = 0$ picks up a coordinate system out of the above class, unique up to multiplication by a root of degree $l+m$ of 1. (If $X = \mathbb{CP}^1$, then of course there is no canonically defined coordinate system, since $(X, P, \mathcal{L})$ has automorphisms.) Now using this coordinate system and the corresponding trivialization of $\mathcal{L}$ one gets a canonically defined image of $(X, P, \mathcal{L})$ in $\mathrm{Gr}(\mathfrak{L})$.

Let the *Teichmüller–Jacoby* space $\mathcal{N}_{g,1,d}$ be the moduli space (i.e., the “set” of equivalence classes) of triples $(X, P, \mathcal{L})$ with $g(X) = g$, and $\deg(\mathcal{L}) = d$. What is of primary importance to us is the fact that $\mathfrak{l}|_{\mathcal{N}_{g,1,d}}$ is an injection (even in the case when X may have double points). Indeed,

1. Meromorphic functions on X may be described as ratios of elements in $\mathfrak{l}(X, P, \mathcal{L})$, thus normalization $\mathrm{Norm}(X)$ of X may be reconstructed basing on $\mathfrak{l}(X, P, \mathcal{L})$;
2. The point P can be reconstructed since we know all the meromorphic functions on X , and know the order of pole at P ;
3. To reconstruct the lifting $\bar{\mathcal{L}}$ of $\mathcal{L}$ to $\mathrm{Norm}(X)$ note that if $\varphi \in V$, then the divisor (of zeros) of φ can be reconstructed as poles of functions in $\mathfrak{l}(\varphi)^{-1} \mathfrak{l}(V)$.
4. Finally, to describe the gluings one needs to perform to get X from $\mathrm{Norm}(X)$ it is sufficient to consider a complement to $\Gamma(\mathcal{L}(\infty \cdot P))$ in $\Gamma(\bar{\mathcal{L}}(\infty \cdot P))$.

As a corollary, the mapping $\mathfrak{l}$ defines an inclusion of $\mathcal{N}_{g,1,d}$ into $\mathrm{Gr}(\mathfrak{L})/G$. Since the image has a natural structure of a topological space, this inclusion defines some natural *compactification* $\bar{\mathcal{N}}_{g,1,d}$ of the Teichmüller–Jacoby space. Indeed, consider the closure of $\mathfrak{l}(\mathcal{N}_{g,1,d})$ in $\mathrm{Gr}(\mathfrak{L})/G$. As we will see it shortly, this closure is an image of a compact smooth manifold². The description of this compactification is very similar to the description of Deligne–Mumford compactification, it is carried out by adding to $\mathcal{N}_{g,1,d}$ so called *semistable* objects. We will see that these object are non-smooth curves and sheaves on them.

²In fact we will see that already the *normalization* of the image is smooth.

The mapping to Universal Grassmannian is a way to collect all the moduli spaces for different genera into “one big heap”. It is relevant to curves of infinite genus since on a back yard of this heap one expects to find moduli spaces of curves of infinite genus.

0.3. Semistable elliptic curves. In this section we consider the case $g = 1$, where one can calculate the mapping to the Universal Grassmannian explicitly. We will see that in this case the closure of the image is smooth³.

Fix an elliptic curve E . Then the set of classes of equivalence of linear bundles of degree 0 (i.e., the Jacobian variety) is isomorphic to the curve itself (after we fix a point P on the curve). Given any point $Q \in E$, the divisor $Q - P$ determines a linear bundle $\mathcal{O}(Q - P)$ on E , and any bundle of degree 0 is isomorphic to exactly one bundle of this form.

In the case of $g = 1$ it is not instructive to consider the Teichmüller–Jacoby manifold *literally*, since it is tainted by the fact that an involution σ of an elliptic curve E sends any bundle $\mathcal{L}$ of degree 0 to $\mathcal{L}^{-1}$, thus the moduli space of bundles of degree 0 on an elliptic curve fixed *up to an isomorphism* is E/σ , i.e., a rational curve. To fix the problem one can add some harmless discrete parameter which would prohibit σ to be an automorphism, say, consider collections $(X, P, \mathcal{L}, \alpha)$, α being a homology class modulo $\mathbb{Z}/3\mathbb{Z}$. However, knowing that we can fix this problem, we are going to ignore it whatsoever, since it can also be avoided by considering curves close to the given one, what we are going to do anyway.

For every elliptic curve E there is a unique number $j \in \mathbb{C}$ such that E is isomorphic to the curve

$$y^2 = (500j - 1)x^3 - 15jx - j.$$

The number j is called the *j-invariant* of the curve. The moduli (Teichmüller) space of curves of genus 1 may be identified with $\mathbb{C}$ via j -invariant. The Teichmüller–Jacoby space is fibered over the Teichmüller space with the fiber being the Jacobian, i.e., the elliptic curve itself (i.e., in the standard notations this space coincides with $\mathcal{M}_{1,2}$). To compactify the Teichmüller space one adds a point with $j = \infty$, obtaining $\mathbb{CP}^1$ as the compactified Teichmüller space.

To visualize the compactification of the Teichmüller space it is more convenient to consider the family of elliptic curves X_ε given by $y^2 = x^2 - x^3 - \varepsilon$. When $\varepsilon \rightarrow 0$ (so $j = \frac{1}{15^3\varepsilon(4-27\varepsilon)} \rightarrow \infty$) one gets a rational curve with a double point. Below we discuss how the corresponding compactification of Teichmüller–Jacoby space looks like. We will see that instead of adding one point, we need to add a rational curve with a selfintersection. The compactified Teichmüller–Jacoby space maps to the compactified Teichmüller space, with the fiber over ∞ being the above singular curve.

Consider an arbitrary elliptic curve E with large a j -invariant, and a bundle of degree 0 on E . One can identify E with X_ε for an appropriate small ε . Fix a

³As we will see it later, $g = 1$ is last case when one does not need any normalization of the image.

point P_ε on X_ε , then for any bundle $\mathcal{L}$ of degree 0 on X_ε one can identify $\mathcal{L}$ with $\mathcal{O}(Q_\varepsilon - P_\varepsilon)$, Q_ε being an appropriate point on X_ε . Fix P_ε to be the point of X which has y -coordinate being 0, and is close to $(1, 0) = P_0$. Suppose that $\mathcal{L} \neq \mathcal{O}$, thus $Q_\varepsilon \neq P_\varepsilon$.

Let us describe meromorphic sections of $\mathcal{L} = \mathcal{O}(Q_\varepsilon - P_\varepsilon)$ with the only pole being at P_ε . It is the same as to describe sections of $\mathcal{O}(Q_\varepsilon + kP_\varepsilon)$, $k \gg 0$, i.e., meromorphic functions on X_ε with a (possible) simple pole at Q_ε , and any pole at P_ε . These sections are uniquely determined by their singular part at P_ε (since there is no meromorphic function on an elliptic curve X_ε which has only a simple pole at Q_ε). Thus to describe the space of these sections, it is sufficient to describe functions f_k on X_ε which have a pole of prescribed order k at P_ε and possibly an additional pole at Q_ε . (Note that $f_0 \equiv 1$ should be considered as a section of $\mathcal{O}(Q_\varepsilon - P_\varepsilon)$ with a pole at P_ε , since holomorphic sections of $\mathcal{O}(Q_\varepsilon - P_\varepsilon)$ are identified with functions having a zero at P_ε .)

Note that y is a meromorphic function on X_ε with a triple pole at infinity of X_ε , x has a double pole at infinity of X_ε . Let P_ε have coordinates $(a_\varepsilon, 0)$, the line through P_ε and Q_ε have the equation $c_\varepsilon x + y = d_\varepsilon$, the third point R_ε of intersection of this line and X_ε has x -coordinate b_ε . If $k = 2k_1$ is even, one can take $f_k = \frac{1}{(x - a_\varepsilon)^{k_1}}$. If $k = 2k_1 - 1$, and $k_1 > 1$, then $y f_{k+1}$ has no pole at infinity, thus can be considered as f_k . What remains is to describe f_1 . Note that $\frac{1}{c_\varepsilon x + y - d_\varepsilon}$ has a pole at P_ε , Q_ε , and at R_ε . Moreover, it has a zero of third order at infinity. Multiplying this function by $x - b_\varepsilon$, we kill the pole at R_ε , thus get $f_1 = \frac{x - b_\varepsilon}{c_\varepsilon x + y - d_\varepsilon}$.

Let us investigate how the space spanned by f_i , $i \geq 0$, depends on X_ε and Q_ε . First of all, ε depends smoothly on Q_ε (if coordinates of Q_ε are q_1, q_2 , then $\varepsilon = q_2^2 - q_1^2 - q_1^3$). From this moment on, we may denote Q_ε by just Q , since ε is a function of Q . Second, x depends smoothly on y and ε in a neighborhood of P_ε on X_ε , *including* the case $\varepsilon = 0$. Thus P_ε depends smoothly on Q , *including* cases $\varepsilon = 0$ and $Q = (0, 0) \in \mathbb{C}^2$. Hence $c_\varepsilon, d_\varepsilon$ depend smoothly on Q . Substituting $y = -c_\varepsilon x + d_\varepsilon$ into $y^2 = x^2 - x^3 - \varepsilon$, we get a cubic equation in x , which has q_1 as a root, and two well-separated roots at a_ε and b_ε . In particular, b_ε depends smoothly on Q . Note that one can consider y as a local coordinate near P_ε , thus the functions f_k can be written as function of y if y is small. We see that all the functions $f_k(y)$ depend smoothly on (q_1, q_2) (at least as far as q_1, q_2 are small, y is small).

If $q_2^2 - q_1^2 - q_1^3 \neq 0$, the described above space $V_{q_1, q_2} = \langle f_i(y) \rangle_{i \geq 0} \subset \mathfrak{L}$ is the image of $(X_\varepsilon, P_\varepsilon, \mathcal{L})$ in the Universal Grassmannian, i.e., it is in $\mathfrak{l}(\mathcal{N}_{1,1,0})$. We see that the closure of $\mathfrak{l}(\mathcal{N}_{1,1,0})$ contains at least the subspaces V_{q_1, q_2} with $q_2^2 - q_1^2 - q_1^3 = 0$. What remains is to give a description of these subspaces as spaces of sections of sheaves on non-smooth curves. We will see that if $(q_1, q_2) \neq (0, 0)$, it is sufficient to consider *linear bundles* on non-smooth curves, but if $(q_1, q_2) = (0, 0)$, a consideration of a sheaf is unavoidable.

It is obvious *which* non-smooth curve we need to consider: the limit X_0 of X_ε when $\varepsilon \rightarrow 0$, with the equation $y^2 = x^2 - x^3$. There is a parameterization mapping $\mathbb{CP}^1 \rightarrow X_0$, $\lambda \mapsto (x, y)$ if $y = \lambda x$. It is a bijection outside of $\lambda = \pm 1$, these two points are both mapped to $(0, 0)$. Thus X_0 is a rational curve with two points glued together. To describe a linear bundle $\mathcal{L}$ on X_0 one needs to describe its lifting to $\mathbb{CP}^1$, and the identification of two fibers over $\lambda = \pm 1$. If the lifting has degree 0, it is a trivial bundle, thus one can describe the gluing by a number $\theta \in \mathbb{C}^*$. Denote this bundle $\mathcal{L}_\theta$, it has a global holomorphic section iff $\theta = 1$.

Note that if $P_0 = (1, 0) \in X_0$, $Q \in X_0$, $Q \neq (0, 0)$, then the mapping $f(\lambda) \rightarrow \frac{\lambda - \lambda_0}{\lambda} f(\lambda)$ gives an isomorphism of $\mathcal{O}(Q - P_0)$ to $\mathcal{L}_\theta$ if $\theta = \frac{1 - \lambda_0}{1 + \lambda_0}$, λ_0 being the coordinate of Q . The defined above functions $f_k(y)$ form a basis of meromorphic sections of $\mathcal{O}(Q - P_0)$, thus we see that for $Q \in X_0$ the defined above subspace $V_{q_1, q_2} \subset \mathfrak{L}$ may be interpreted as $\mathfrak{l}(X_0, P_0, \mathcal{L}_\theta)$.

Now investigate what happens if $Q \rightarrow (0, 0)$ along X_0 , i.e., $\theta \rightarrow 0$ or $\theta \rightarrow \infty$. Instead of y , consider the coordinate λ in a neighborhood of P_0 on X_0 . In this coordinate the space $V(X_0, P_0, \mathcal{L}_\theta)$ is the set of polynomials $p(\mu)$ in $\mu = \lambda^{-1}$ which satisfy $p(1) = \theta p(-1)$. Denote this subspace of $\mathbb{C}[\mu]$ by V_θ . Obviously, the subspace V_θ depends smoothly on $\theta \in \mathbb{CP}^1$, V_0 consists of polynomials with a zero at 1, V_∞ consists of polynomials with a zero at -1 . Moreover, images of V_0 and V_∞ in $\text{Gr}(\mathfrak{L})/G$ coincide, since $V_\infty = \frac{\mu+1}{\mu-1}V_0$, and $\frac{\mu+1}{\mu-1}$ is smooth near $\mu = \infty$, therefore $f(\mu) \rightarrow \frac{\mu+1}{\mu-1}f(\mu)$ corresponds to an element of G .

Returning back to the representation of V_{q_1, q_2} , $q_1 = q_2 = 0$, in terms of f_k , we see that $f_1 = x/y = \lambda^{-1}$, $f_2 = (x-1)^{-1} = -\lambda^{-2}$, thus $f_{2k} = \pm \lambda^{-2k}$; $y = \lambda(1 - \lambda^2)$, thus $f_{2k+1} = \frac{1 - \lambda^2}{\lambda^{2k+1}} = \lambda^{-2k-1} - \lambda^{-2k+1}$. We see that this space is $\mathbb{C}[\lambda^{-1}]$. Hence the limit of spaces of sections of $\mathcal{O}(Q - P)$ when $Q \rightarrow (0, 0)$ is isomorphic to the space of meromorphic functions on the normalization $\mathbb{CP}^1$ of X_0 with the only pole at $\lambda = 0$, i.e., with $\mathfrak{l}(\mathbb{CP}^1, 0, \mathcal{O})$. If we want to describe this space in the same terms as we described the mapping $\mathfrak{l}$, we should consider sections of $\mathcal{O}$ on $\mathbb{CP}^1$ as sections of the direct image $\pi_*\mathcal{O}$ on X_0 , π being the projection of $\mathbb{CP}^1$ to X_0 . Since $\pi_*\mathcal{O}$ is no longer a sheaf of sections of a linear bundle, we conclude that we *need* to consider sheaves instead of bundles.

The only thing which remains to prove is that the completion of the image $\mathfrak{l}(\mathcal{N}_{1,1,0})$ is smooth. We had already shown the part of the completion we consider here is an image of a neighborhood of 0 in $\mathbb{C}^2$. What remains to prove is that the mapping $\mathbb{C}^2 \xrightarrow{V_\bullet} \text{Gr}(\mathfrak{L})/G$ has a non-degenerate derivative at $(0, 0)$. It is sufficient to pick up a section of the projection $\text{Gr}(\mathfrak{L}) \rightarrow \text{Gr}(\mathfrak{L})/G$, and show that a lifting of $V_\bullet$ to this section has a non-degenerate derivative at $(0, 0)$. Again, it is sufficient to restrict this mapping to X_0 and show that the derivatives along two branches of X at $(0, 0)$ are independent.

One branch of X_0 gives a family of subspaces V_θ , $\theta \approx 0$, $V_\theta \subset \mathbb{C}[\lambda^{-1}]$. It is a simple but tedious calculation to show that the lifting defined in Section 0.2 results

in a smooth lifting of $V_\bullet$, and that the smooth liftings resulting from two branches are non-degenerate, and have different derivatives indeed.

We see that to describe the elements in the completion of $\mathfrak{l}(\mathcal{N}_{1,1,0})$ in a way similar to the description of $\mathfrak{l}$ one needs to consider both degenerated curves and degenerated bundles (i.e., sheaves) on these curves. Using the fact that the space of sections of $\pi_*\mathcal{O}$ is close to the set of sections of $\mathcal{O}(Q)$ if Q is close to the double point $(0,0)$ of X_0 , it is reasonable to abuse notations and write $\pi_*\mathcal{O}$ as $\mathcal{O}(O)$, here $O = (0,0)$.

0.4. Compactified moduli spaces. We start with a description of elements of the would-be compactified moduli spaces. As we have seen it in Section 0.3, we need to allow the curves to have double points, and allow a generalization of a notion of bundle. We call such bundles *non-smooth* bundles, in fact they are sheaves which are (locally) direct images of bundles on the normalization⁴ of the curve.

Definition 0.1. Let Y be a connected curve with only singularities $\text{Sing}(Y)$ being double points, let $\tilde{Y}$ be the normalization of Y . Define $g(Y) = g(\tilde{Y}) + \text{card}(\text{Sing}(Y))$, for a subset $S_0 \subset \text{Sing}(Y)$ let Y_{S_0} be the curve obtained by ungluing points in S_0 . A *smooth linear bundle* $\mathcal{L}$ on Y is a linear bundle $\tilde{\mathcal{L}}$ on $\tilde{Y}$ with a fixed identification of fibers at points of $\tilde{Y}$ over the same point on Y . Define *degree* of a smooth linear bundle as the $\deg(\tilde{\mathcal{L}})$. A *section* of $\mathcal{L}$ is a section of $\tilde{\mathcal{L}}$ compatible with identifications of fibers.

A (not necessarily smooth) *linear bundle* $\mathcal{L}$ on Y is a collection $(S_0, \bar{\mathcal{L}})$ of $S_0 \subset \text{Sing}(Y)$ and a smooth linear bundle $\bar{\mathcal{L}}$ on Y_{S_0} . A *section* of $\mathcal{L}$ is a section of $\bar{\mathcal{L}}$. Let $\deg(\mathcal{L}) = \deg(\bar{\mathcal{L}}) + \text{card}(S_0)$, call S_0 the set of *double points* of $\mathcal{L}$. Let $P \subset \text{Smooth}(Y)$ be a finite collection of points. Call the collection $(Y, P, \mathcal{L})$ *semistable* if $(Y, P, \mathcal{L})$ has no infinitesimal automorphisms, i.e., if any rational connected component of $\tilde{Y}$ has at least three points in P or over double points of Y , and if Y is a (smooth) elliptic curve, then either $P \neq \emptyset$, or $\deg \mathcal{L} \neq 0$.

In other words, for a “non-smooth linear bundle” $\mathcal{L}$ the double points of the curve are broken into two subsets: for one (smooth points of $\mathcal{L}$) we fix identifications of fibers of the lifting to normalizations, for another one (as we have seen in Section 0.5, one should consider them as *poles* of $\mathcal{L}$) local sections of $\mathcal{L}$ may have “different” values at two branches of Y which meet at a given point.

Note that the smooth structures on the Teichmüller space and on the Jacobian of a smooth curve provide an atlas on the set of equivalence classes of smooth curves/bundles. The explicit deformation which we are going to describe now will provide an atlas in a neighborhood of a class of a semistable curve.

Consider a curve Y as in the definition, let $Q \in Y$ be a double point, and y_1, y_2 be coordinates on the branches of Y near Q , so Y is locally isomorphic to $y_1 y_2 = 0$.

⁴Recall that in the case $\dim = 1$ *normalization* coincides with ungluing double points and straightening out cusps.

Let $\mathcal{L}$ be a linear bundle over Y with a fixed trivialization near Q . Suppose that Q is a double point of $\mathcal{L}$, thus local sections of $\mathcal{L}$ can be written as (unrelated) functions $f_1(y_1), f_2(y_2)$. Note that trivialization of $\mathcal{L}$ induces identification of fibers of $\tilde{\mathcal{L}}$ over Q . Let $\mathring{\mathcal{L}}$ be the bundle on Y (smooth near Q) obtained by addition of this identification to $\mathcal{L}$. Extend the bundle $\mathring{\mathcal{L}}$ from Y to $Y \cup \{(y_1, y_2) \mid |y_1|, |y_2| \ll 1\}$ trivially using the trivialization of $\mathcal{L}$ near Q , denote this extension as $\tilde{\mathcal{L}}$.

Define a deformation $(Y_{\varepsilon_1, \varepsilon_2}, \mathcal{L}_{\varepsilon_1, \varepsilon_2})$ of $(Y, \mathcal{L})$ (here $\varepsilon_1, \varepsilon_2 \in \mathbb{C}$ are fixed small numbers) by the following recipe:

1. if $\varepsilon_1 \varepsilon_2 = 0$, then $Y_{\varepsilon_1, \varepsilon_2} = Y$;
2. if $\varepsilon_1 = \varepsilon_2 = 0$, then $\mathcal{L}_{\varepsilon_1, \varepsilon_2} = \mathcal{L}$;
3. if $\varepsilon_1 \neq 0$, but $\varepsilon_2 = 0$, then $\mathcal{L}_{\varepsilon_1, \varepsilon_2}$ is the bundle $\mathring{\mathcal{L}}(P_{\varepsilon_1})$, here P_{ε_1} is the point on Y with $y_1 = \varepsilon_1$; similarly for $\varepsilon_1 = 0$, but $\varepsilon_2 \neq 0$;
4. If $\varepsilon_1 \neq 0, \varepsilon_2 \neq 0$, then $Y_{\varepsilon_1, \varepsilon_2}$ is obtained by gluing the hyperbola $y_1 y_2 = \varepsilon_1 \varepsilon_2$ to Y (with a small neighborhood of Q removed) via coordinate projections, and $\mathcal{L}_{\varepsilon_1, \varepsilon_2} = \mathring{\mathcal{L}}(P_{\varepsilon_1, \varepsilon_2})$, here $\mathring{\mathcal{L}}_{\varepsilon_1, \varepsilon_2}$ is the restriction of $\tilde{\mathcal{L}}$ to $Y_{\varepsilon_1, \varepsilon_2}$, the point $P_{\varepsilon_1, \varepsilon_2}$ is the point on hyperbola with coordinates $(\varepsilon_1, \varepsilon_2)$.

If Y had some marked points $\{P_i\} \subset \text{Smooth}(Y)$, then $Y_{\varepsilon_1, \varepsilon_2}$ has the same marked points (correctly defined since $Y_{\varepsilon_1, \varepsilon_2}$ is identified with Y outside a small neighborhood of Q).

Note that if $P \in \text{Smooth}(Y)$, then $\mathfrak{l}(Y, P, \mathcal{L})$ is well-defined in $\text{Gr}(\mathfrak{L})/G$.

Lemma 0.2. *$g(Y_{\varepsilon_1, \varepsilon_2}) = g(Y)$, $\deg \mathcal{L}_{\varepsilon_1, \varepsilon_2} = \deg \mathcal{L}$. If $(Y, P, \mathcal{L})$ is semistable, then $(Y_{\varepsilon_1, \varepsilon_2}, P, \mathcal{L}_{\varepsilon_1, \varepsilon_2})$ is semistable too. If $P \in \text{Smooth}(Y)$, then $\mathfrak{l}(Y_{\varepsilon_1, \varepsilon_2}, P, \mathcal{L}_{\varepsilon_1, \varepsilon_2}) \in \text{Gr}(\mathfrak{L})/G$ depends smoothly on $\varepsilon_1, \varepsilon_2$.*

Similarly, if Q is a double point of Y , but not a double point of $\mathcal{L}$, define a one-parametric deformation $(Y_\varepsilon, \mathcal{L}_\varepsilon)$ by gluing to Y a hyperbola $y_1 y_2 = \varepsilon$ without changing $\mathcal{L}$ (far from Q) and the trivialization of $\mathcal{L}$. A statement similar to the above lemma continues to be true.

Let $\bar{\mathcal{N}}_{g,1,d}$ be the set of equivalence classes of semistable collections $(Y, P, \mathcal{L})$, let $\bar{\mathcal{N}}_{g,1,d}^{(m,k)} \subset \bar{\mathcal{N}}_{g,1,d}$ be the subset of $\bar{\mathcal{N}}_{g,1,d}$ consisting of curves with exactly m double points and bundles with exactly k double points ($k \leq m$). Note that $\bar{\mathcal{N}}_{g,1,d}^{(0,0)} = \mathcal{N}_{g,1,d}$, and the complement to $\mathcal{N}_{g,1,d}$ in $\bar{\mathcal{N}}_{g,1,d}$ is a disjoint union of $\bar{\mathcal{N}}_{g,1,d}^{(m,k)}$. Call these subsets *strata* of $\bar{\mathcal{N}}_{g,1,d}$. Moreover, note that each $\bar{\mathcal{N}}_{g,1,d}^{(m,k)}$ carries a natural smooth structure (inherited from the smooth structures on Teichmüller spaces and Jacobians). (In fact the above smooth structures can be refined to structures of orbifolds, but we ignore this refinement here.)

Let $(Y, y_0, \mathcal{L}) \in \bar{\mathcal{N}}_{g,1,d}^{(m,k)}$, $\varepsilon \in \mathbb{C}^{m+k}$ is small. Taking coordinate systems in neighborhoods of double points, and trivializations of $\mathcal{L}$ in these neighborhoods, we obtain a

deformation $(Y_\varepsilon, P, \mathcal{L}_\varepsilon) \in \bar{\mathcal{N}}_{g,1,d}$. Since $\bar{\mathcal{N}}_{g,1,d}^{(m,k)}$ has a natural smooth structure, we see that a piece of $\bar{\mathcal{N}}_{g,1,d}$ is fibered over $\bar{\mathcal{N}}_{g,1,d}^{(m,k)}$ with fibers being small balls in $\mathbb{C}^{m+k}$.

Define a structure of an manifold on $\bar{\mathcal{N}}_{g,1,d}$ using the above fibration as an atlas in a neighborhood of a point in $\bar{\mathcal{N}}_{g,1,d}^{(m,k)}$.

Proposition 0.3. $\bar{\mathcal{N}}_{g,1,d}$ is a smooth compact manifold (orbifold). $\mathfrak{l}(\bar{\mathcal{N}}_{g,1,d})$ coincides with the closure of $\mathfrak{l}(\bar{\mathcal{N}}_{g,1,d})$.

Proposition 0.4. $\mathfrak{l}(X, P, \mathcal{L}) = \mathfrak{l}(X', P', \mathcal{L}')$ iff $(X, P, \mathcal{L})$ becomes isomorphic to $(X', P', \mathcal{L}')$ after ungluing of double points of $\mathcal{L}$ and $\mathcal{L}'$.

This means that $\bar{\mathcal{N}}_{g,1,d}$ is a natural smooth compactification of $\mathcal{N}_{g,1,d}$. Moreover, the explicit coordinates in a neighborhood of a stratum show that the boundary $\bar{\mathcal{N}}_{g,1,d} \setminus \mathcal{N}_{g,1,d}$ is a divisor with normal intersections. As it follows from Proposition 0.4, the image $\mathfrak{l}(\bar{\mathcal{N}}_{g,1,d})$ can be obtained from $\bar{\mathcal{N}}_{g,1,d}$ by contracting smooth submanifolds into points, thus the normalization of the image coincides with $\bar{\mathcal{N}}_{g,1,d}$.

0.5. Adjacency of moduli spaces and Shottki model. We have seen that the consideration of the mapping to the universal Grassmannian leads to a remarkable compactification of Teichmüller–Jacoby space. Moreover, one can use this compactification to define inclusions of compactified Teichmüller–Jacoby spaces for different g, d one into another. These inclusions are going to be compatible with the mapping $\mathfrak{l}$ to Universal Grassmannian, and may be constructed studying the properties of the mapping $\mathfrak{l}$.

Consider a curve X with a bundle $\mathcal{L}$ (we allow X and $\mathcal{L}$ to have double points). Given two smooth points Q_1, Q_2 on X one can construct a curve $X_{Q_1 Q_2}$ obtained by gluing Q_1 with Q_2 . This curve has one more double point, and genus $g(X) + 1$. One can consider the bundle $\mathcal{L}_{Q_1 Q_2}$ on $X_{Q_1 Q_2}$ which has an extra double point at $Q_1 \sim Q_2$, otherwise coincides with $\mathcal{L}$. Note that sections of $\mathcal{L}$ can be considered as sections of $\mathcal{L}_{Q_1 Q_2}$ and visa versa, hence $\mathfrak{l}(X, P, \mathcal{L}) = \mathfrak{l}(X_{Q_1 Q_2}, P, \mathcal{L}_{Q_1 Q_2})$ for any P .

Using this remark one can easily include the compactification of one moduli space into the boundary of another one. Indeed, one can generalize the construction of $\bar{\mathcal{N}}_{g,1,d}$ to the case of n marked points $\{P_i\}$ (instead of one P), the only change being that one needs to consider $\text{Gr}(\mathfrak{L}^n)$ instead of $\text{Gr}(\mathfrak{L})$. Now associate an element of $\bar{\mathcal{N}}_{g+l,n-2l,d+l}$ to an element $(Y, \{P_i\}, \mathcal{L}) \in \bar{\mathcal{N}}_{g,n,d}$ as follows: take first $2l$ points out of $\{P_i\}$, and glue them pairwise. Since we do not glue fibers of $\mathcal{L}$ at these points, the resulting double points on the resulting curve Y_1 are double points of a bundle $\mathcal{L}_1$. Clearly, $(Y_1, \{P_{2l+i}\}, \mathcal{L}_1) \in \bar{\mathcal{N}}_{g+l,n-2l,d+l}$. Thus we obtain a mapping $\bar{\mathcal{N}}_{g,n,d} \rightarrow \bar{\mathcal{N}}_{g+l,n-2l,d+l}$. Note that this correspondence does not change the space of global sections of $\mathcal{L}$, thus the image under the map $\mathfrak{l}$.

Since we are especially interested in $\bar{\mathcal{N}}_{g,1,d}$, let us modify the above construction to get a mapping $\bar{\mathcal{N}}_{g,1,d} \rightarrow \bar{\mathcal{N}}_{g+l,1,d+l}$. To do this it is sufficient to define a mapping $\bar{\mathcal{N}}_{g,1,d} \rightarrow \bar{\mathcal{N}}_{g,1+2l,d}$. We will define such a mapping for every *most degenerate* curve,

which is a semistable curve with the class on a stratum of $\dim = 0$ on the moduli space. The mapping will be given by *gluing* this curve to the given curve. Since there are finitely many most degenerate curves (they are enumerated by appropriately colored trees), we will obtain a finite number of mappings $\bar{\mathcal{N}}_{g,1,d} \rightarrow \bar{\mathcal{N}}_{g,1+2l,d}$.

A most degenerate curve $(Z, \{z_j\})$ of genus 0 with $2l + 2$ marked points z_j , $0 \leq j \leq 2l + 1$ has $2l - 1$ double points, and its normalization has $2l$ connected components. Any bundle of degree 0 over Z is trivial. Given a curve $(Y, P_0, \mathcal{L})$, glue Z to Y by identifying P_0 and z_0 and identify $\mathcal{L}|_{P_0}$ with $\mathcal{O}_Z|_{z_0}$ arbitrarily, denote the resulting bundle $\mathcal{L}'$. The resulting collection $(Y \cup Z, \{z_j\}, \mathcal{L}')$ is obviously an element of $\bar{\mathcal{N}}_{g,1+2l,d}$, and gluing together z_j , $1 \leq j \leq 2l$, we obtain an element of $\bar{\mathcal{N}}_{g+l,1,d+l}$.

We see that $\mathfrak{l}(X, P, \mathcal{L})$ can be approximated by $\mathfrak{l}(X', P', \mathcal{L}')$ (here X and X' are smooth curves) with $g(X') < g(X)$ if $(X, P, \mathcal{L})$ is close to a subset of codimension 2 which consists of semistable curves/bundles such that a bundle has a double point.

To visualize better the above surgery we improve the description in two ways. First, let us revisit the process of ε -deformation.

We glue a hyperbola $y_1 y_2 = \varepsilon$ to the coordinate lines $y_1 = 0$, $y_2 = 0$ via the coordinate projections. One can momentarily see that this is equivalent to gluing together two regions $\{|y_1| \geq \sqrt{|\varepsilon|}\}$, $\{|y_2| \geq \sqrt{|\varepsilon|}\}$ along the boundary via $y_2 = \varepsilon/y_1$.

Given a bundle $\mathcal{L}$ with trivializations near Q_1 and Q_2 one obtains a bundle $\mathring{\mathcal{L}}$ on the resulting curves (its sections are sections of $\mathcal{L}$ outside of the disks with compatible restrictions to the disk boundaries). The deformed bundle is $\mathring{\mathcal{L}}(Q)$, here Q is close to the circle $|y_1| = |y_2| = \sqrt{|\varepsilon|}$ on the deformed curve.

There is an alternative description of the deformed curve. Above we were removing two disks of radii ε_1 and ε_2 near points y_1 , y_2 and gluing the boundaries. The radii of disks were variable, so it was hard to visualize the picture. Remove instead bigger disks of (fixed) radii δ_1 , δ_2 . What remains is to glue two annuli $\{\varepsilon_i < |z| < \delta_i\}$, $i = 1, 2$, to the resulting manifold with a boundary, and glue inner boundaries of annuli together. Each annulus is conformally equivalent to a cylinder with ratio length/radius being $\log \frac{\delta_i}{\varepsilon_i}$. Thus we need to glue in a cylinder $S^1 \times (0, L)$ of *conformal length* $L = \log \frac{\delta_1 \delta_2}{\varepsilon_1 \varepsilon_2}$. If $\varepsilon_{1,2}$ decrease, it is equivalent to gluing longer and longer *handles* between circles of radii $\delta_{1,2}$. Note also that we modify $\mathcal{L}$ by adding a pole inside this handle.

Second, note that the above gluing mapping $y_2 = \varepsilon/y_1$ is defined not only on the boundary of the above regions, but in the regions themselves. Thus continuing the identification of boundaries, one can identify the region $\{|y_2| \geq \sqrt{|\varepsilon|}\}$ with $\{|y_1| \leq \sqrt{|\varepsilon|}\}$. If a part of curve corresponds to a subset of $\{|y_2| \geq \sqrt{|\varepsilon|}\}$, then it is identified with a subset of $\{|y_1| \leq \sqrt{|\varepsilon|}\}$. Suppose that the curve we deform has rational smooth components. If one component Y_i contains l_i double points, then to perform the above deformation we remove l_i disks of radii ε_{ik} , $k = 1, 2$, around these

points. What remains is a sphere without l_i disks, and we need to glue several such spheres together identifying boundaries by fraction-linear mappings. Say, if $i \neq j$ and Y_j is glued to Y_i via gluing φ_i , then $\varphi(Y_j)$ is a subset of a removed disk for Y_i . Consider now the union $Y_i \cup \varphi_i(Y_j)$. It is again a sphere with several disks removed, and the boundaries of this disks are identified with the boundaries of other components (or different circles on $Y_i \cup \varphi_i(Y_j)$) via fraction-linear mappings.

Continuing this process as long as we can, the result is *one* sphere with several disks removed, and boundaries of these disks are identified pairwise via fraction-linear mappings. This is so called *Shottki model* of the curve, and we see that the analogue of Deligne–Mumford compactification we consider here leads naturally to this model of deformed curves. One can do a similar thing in the case when the initial curve contains non-rational components as well. In this case one should restrict attention to the “connected part” of the curve which consists of rational components.

We conclude that in the simplest case the above process of deformation may be described as this (for one particular choice of degenerate rational curve with 3 marked points): Take two points Q_1, Q_2 which are close to the marked point P and much closer to each other than to P . Now remove two non-intersecting disks around Q_1 and Q_2 , and glue a very long handle between boundaries of these disks. Here a disk which contains P, Q_1 and Q_2 plays the rôle of the “old” marked point, we assume that z_0 and z_4 (using the notations from the beginning of this section) are on the same smooth component of the degenerate rational curve.

0.6. Second compactification. The previous section shows that compactified moduli spaces are included one into another, thus one can consider the direct limit, i.e., the *union* of these spaces. The mapping $\mathfrak{l}$ identifies this union $\mathfrak{N}_{d-g} = \bigcup_l \mathcal{N}_{g+l,1,d+l}$, with a subset of the Universal Grassmannian, and (at last!) we have the ingredients necessary for the discussion what the closure of this subset may look like.

Consider an element $(Y, P, \mathcal{L})$ of $\mathcal{N}_{g+l,1,d+l}$ for a very big l . Suppose that it is close to an element $(Y', P', \mathcal{L}')$ of $\mathcal{N}_{g,1,d}$. To get Y we glue a most degenerate rational curve to Y' , the result is a curve with $2l$ components and $2l + 2$ marked points. Next we glue marked points pairwise, and ε -deform the resulting curve at the double points. However, the double points are naturally broken into two categories: double points on the attached rational curve, and doubled marked points. Let us change the order of gluing and deformations: glue in the degenerate curve, ε -deform the double points on this curve, then glue marked points together and ε -deforming them. Note that the former double points are not double points of the linear bundle, but the latter ones are double points for the bundle.

Deformation of double points on a degenerate rational curve leads to a Shottki model for some curve, i.e., a sphere with several disks removed. However, since the genus is 0, there is no removed disks at all, thus we get a sphere with $2l + 2$ marked points. The condition that the parameters of deformation are small is translated into the fact that double ratios of marked points on the sphere are very big, i.e., that

conformally the centers are *well separated*. Deform now the double point where the degenerated curve is glued to Y' . This leads to the part of the above sphere being identified (by a fraction-linear mapping) with a disk around P' , and the marked points go to points of this disk. We conclude that this part of deformation corresponds to picking up a collection of points which are close to P' and conformally well separated.

The second part of the deformation is the removing of small disks around the marked points, and gluing together the boundaries of these disks (or, what is the same, gluing long handles to boundaries of bigger fixed disks). In the most important case $g = 0$, thus the description of a neighborhood of a point on $\mathfrak{N}_k$ is related to studying curves of genus g obtained by gluing together $2g$ small conformally well separated disks on $\mathbb{CP}^1$. Note that the notion of being *well separated* depends on the combinatorics of degenerated curves we glue in to get the limit point of $\mathfrak{N}_k$.

Since we expect that $I(\mathfrak{N}_k)$ is dense in the space of solutions of an integrable system, one can approximate a point in this space by a sequence of curves and bundles. Assuming the best case scenario, we can obtain a term of this sequence by a small deformation of the previous term (here a small deformation is taken in the sense of algebraic geometry, i.e., one takes a couple of points on the curve, glues them together and deforms the double point slightly). We see that such a point of the space can be described as a *curve of potentially infinite* genus, i.e., a sequence of the curves where the next one is obtained by gluing long handles to the previous one.

In this paper we show that one can consider a *curve of actually infinite* genus instead. Such a curve is obtained in the same manner as the above sequence of finite-genus curves: one takes an infinite collection of “well-separated” points on $\mathbb{CP}^1$, removes a disk of a small radius around each point, and glues the boundaries pairwise.⁵ We show that under suitable conditions the standard theorems of algebraic geometry hold for the resulting curves as well. This shows that the *completed* moduli space $\mathfrak{M}_{-1}$ of such curves+bundles may be important in algebraic geometry. If this moduli space coincides with the whole space of solutions (i.e., the best case scenario has place indeed), one can see that the completed moduli space has a very simple topology. Indeed, by Cauchy–Kovalevskaya theorem the space of solutions is identified with the space of initial data for the solutions, which is a topological linear space.

One can see that under assumption that the moduli space described in this paper coincides with the set of solutions of integrable systems, the study of compactified moduli spaces simplifies a lot by a transition to the case of infinite genus. This simplification may suggest additional approaches to the problem of studying the moduli spaces in finite genus as well.

⁵The exact meaning of the above terms “well-separated collection” and circles being “of small radii” is the main topic of this paper. The answers vary a little bit depending on what problem one considers, for the list of results see Section 0.12.

Unfortunately, at this moment it is unclear whether the best case scenario is applicable to the moduli space as a whole. As we noted above, the results of [7] show that this is true for the real part of the hyperelliptic subset of the moduli space.

0.7. Growth conditions and divisors. The main target of this paper is to describe how to quantify the conditions on the collection of circles to be “conformally well separated” and “of small radii” so that the main theorems of algebraic geometry are still valid for the resulting curves of infinite genus. However, before we discuss the conditions on the nonlinear data (curves and bundles), we should discuss the simpler conditions on the linear data (sections of above bundles).

The basic theorem of 1-dimensional algebraic geometry states that any linear bundle on a compact analytic curve has a finite-dimensional space of global sections. The moment we try to drop the compactness condition this theorem breaks, since any non-compact analytic curve is a Stein manifold, thus any sheaf on it has a giant space of global sections.

One way to fix this situation is to consider growth conditions. Let $\bar{X}$ be a compact curve with a linear bundle $\bar{\mathcal{L}}$. Suppose that $\bar{X}$ and $\bar{\mathcal{L}}$ are provided with Hermitian metrics. Pick up a point $\infty \in \bar{X}$, let $X = \bar{X} \setminus \{\infty\}$, $\mathcal{L} = \bar{\mathcal{L}}|_X$. Then bounded sections of $\mathcal{L}$ can be uniquely extended to analytic sections of $\bar{\mathcal{L}}$, thus $\mathcal{L}$ has a finite-dimensional space of bounded sections. Similarly, if we consider sections of $\mathcal{L}$ with magnitude going to 0 when $x \rightarrow \infty$, it is the same as to consider sections of $\bar{\mathcal{L}}(-1 \cdot \infty)$ on $\bar{X}$, here $-1 \cdot \infty$ is a divisor on $\bar{X}$.

Going in a different direction, consider L_2 -sections of $\mathcal{L}$. It is easy to see that this is also equivalent to consideration of sections of $\bar{\mathcal{L}}$. If we consider a different metric on X , say, $\frac{dx^2}{\text{dist}(x, \infty)^{2\alpha}}$, then to consider holomorphic on X L_2 -sections is the same as to consider $\bar{\mathcal{L}}([\alpha] \cdot \infty)$. Similarly, different choice of metric on $\mathcal{L}$ will lead to different shift of $\bar{\mathcal{L}}$ by a divisor at ∞ .

We see that different growth conditions applied to sections of $\mathcal{L}$ lead to different “effective” continuations of $\mathcal{L}$ to $\bar{X}$. Obviously, the situation becomes more complicated when $\bar{X} \setminus X$ consists of more than one point. If $\bar{X} \setminus X$ is discrete, different possible choices of growth conditions lead to a lattice which is isomorphic to the lattice of divisors on $\bar{X}$ concentrated on $\bar{X} \setminus X$. Situation goes out of control if $\bar{X} \setminus X$ is a “massive” set, or $\bar{X}$ is not a smooth manifold at all. In such cases different choices of metrics on X and $\mathcal{L}$ (or some other data controlling the growth of sections) form a very complicated lattice. However, there is a way to select “good” elements of this lattice.

0.8. Riemann–Roch and growth control. The Riemann–Roch theorem says that on a compact analytic curve there is a relationship between a *degree of $\mathcal{L}$* , which is an easily calculable geometric characteristics of $\mathcal{L}$, and the dimensions of spaces of global sections of $\mathcal{L}$ and $\omega \otimes \mathcal{L}^{-1}$, ω being the linear bundle of holomorphic forms:

$$\dim \Gamma(\mathcal{L}) - \dim \Gamma(\omega \otimes \mathcal{L}^{-1}) = \deg \mathcal{L} - g + 1.$$

Let us show how this theorem might be applied to picking up “correct” growth conditions on non-compact analytic curves.

Consider some fixed growth conditions for X . They select a new sense for the functor $\Gamma(X, \bullet)$. Suppose again that $\bar{X} \setminus X$ is discrete, $\mathcal{L} = \bar{\mathcal{L}}|_X$. We have seen above that consideration of these growth conditions is equivalent to consideration of $\bar{\mathcal{L}}(D)$, here D is some divisor at infinity, i.e., $\Gamma(X, \mathcal{L}) = \Gamma(\bar{X}, \bar{\mathcal{L}}(D))$. One should expect that D does not depend on $\mathcal{L}$ (at least when $\mathcal{L}$ does not change a lot). Suppose that there is a way to determine $\deg \bar{\mathcal{L}}$ basing on $\mathcal{L}$, say, we describe bundles by divisors on X .

If the dimensions of sections of linear bundles on X with given growth conditions satisfy the Riemann–Roch theorem, then an easy calculation shows that $2D \sim 0$. In particular, suppose that $\mathcal{L}$ was a restriction of a bundle $\bar{\mathcal{L}}$ of half-forms on $\bar{X}$, i.e., $\bar{\mathcal{L}} \otimes \bar{\mathcal{L}} \simeq \omega$. Then we see that the bundle on $\bar{X}$ which corresponds to $\bar{\mathcal{L}}|_X$ with given growth conditions is $\bar{\mathcal{L}}(D)$. Note that $\bar{\mathcal{L}}(D) \otimes \bar{\mathcal{L}}(D) = \omega$. In particular, the growth conditions which satisfy Riemann–Roch theorem “preserve” the set of bundles of half-forms (or θ -characteristics): the sections on X of one θ -characteristic $\bar{\mathcal{L}}$ which satisfy the growth conditions can be naturally identified with sections on $\bar{X}$ of some other θ -characteristic $\bar{\mathcal{L}}(D)$.

Suppose now that the growth conditions on sections of $\mathcal{L}$ are picked up in some invariant way—whatever it means. Because of that we expect that D is a linear combination of points on $\bar{X} \setminus X$ with the same coefficients. Since $\deg D = 0$, thus D is 0.

We see that Riemann–Roch provides a selection criterion for picking up growth conditions which “do not add” points at infinity to a given divisor. By analogy, one can apply the same criterion in cases when $\bar{X} \setminus X$ is massive or $\bar{X}$ does not exist: If there is an invariant way to describe the growth conditions, and the Riemann–Roch theorem holds for a class of bundles, then one may describe the part at infinity of the divisor of a section of such a bundle: a section has no “poles” at infinity if it satisfies the growth conditions.

0.9. Handles, Sobolev spaces, and representations of $\mathrm{SL}_2(\mathbb{C})$. Let us apply heuristics from the previous section to the case of a curve X of infinite genus, i.e., a Riemannian sphere with an infinite number of disks removed, and infinite number of handles glued in along the cut lines (or just any surface glued in from “pants” in such a way that the graph of gluing is of infinite genus). This is not a compact Riemannian surface, so one needs to consider “infinities” of this surface, and fix some growth conditions near these infinities.

Moreover, if the genus is infinite, one needs to be especially careful with Riemann–Roch theorem, since the right-hand side contains genus $g(X)$ (i.e., number of handles) of the curve, which is not a number any more, but infinity. To save the theorem, let us consider the quantity $\deg \mathcal{L} - g + 1$ as a unity. One way to do it is to consider some fixed bundle $\mathcal{M}$ on X which is “naturally constructed” and satisfies the condition

$\deg \mathcal{M} = g - 1$ for finite-genus X , and consider sections of $\mathcal{L} \otimes \mathcal{M}$ instead of sections $\mathcal{L}$. Then the Riemann–Roch theorem may be rewritten as

$$\dim \Gamma(\mathcal{L} \otimes \mathcal{M}) - \dim \Gamma(\omega \otimes \mathcal{M}^{-1} \otimes \mathcal{L}^{-1}) = \deg \mathcal{L}.$$

Note that $\deg \omega \otimes \mathcal{M}^{-1} = \deg \mathcal{M}$, so $\mathcal{N} = \omega \otimes \mathcal{M}^{\otimes -2}$ is a “naturally constructed” bundle of degree 0. There are 2^g different square roots of a given bundle of degree 0, and fixing one solution $\mathcal{R}$ to $\mathcal{R}^2 = \omega \otimes \mathcal{M}^{\otimes -2}$ we may change $\mathcal{M}$ to $\mathcal{M} \otimes \mathcal{R}^{-1}$. After such a change the formula simplifies to

$$\dim \Gamma(\mathcal{L} \otimes \mathcal{M}) - \dim \Gamma(\mathcal{L} \otimes \mathcal{M}^{-1}) = \deg \mathcal{L},$$

which has an additional advantage of being symmetric w.r.t. $\mathcal{L} \mapsto \mathcal{L}^{-1}$. Moreover, $\mathcal{M}$ is now a solution of $\mathcal{M}^2 = \omega$.

We come to the following heuristic: to consider Riemann–Roch theorem for a curve X with infinitely many handles one should fix a bundle of half-forms $\mathcal{M} = \omega^{1/2}$ on X , consider a linear bundle $\mathcal{L}$ of finite degree, and apply appropriate growth conditions to sections of $\mathcal{L} \otimes \mathcal{M}$. We expect that the choice of growth conditions is very restricted, since any possible change is equivalent to consideration of different square root of ω (see Section 0.8).

One should expect that thus obtained growth conditions are in some way “invariant” (since so rigid), thus geometrically defined. It is still meaningful to apply these conditions in the case $\mathcal{L} = \mathcal{O}$, when $\mathcal{L} \otimes \mathcal{M}$ is the sheaf of half-forms. We come to the following conclusions:

Conditions of boundness (i.e., L_∞ -topology) cannot be applied (since there is no geometrically-defined norm on the fibers of the bundle of half-forms), as well as L_2 -type restrictions (since a square of a holomorphic half-form is a half-form on $X_{\mathbb{R}}$, thus cannot be invariantly integrated). One can easily see that the fourth degree of a half form α (more precise, $\alpha^2 \otimes \bar{\alpha}^2$) is a top-degree-form on $X_{\mathbb{R}}$, thus a good candidate is the condition of integrability of fourth degree, i.e., L_4 -topology.

It turns out that generic Banach spaces (like L_4) are not convenient for cut-and-glue operations we are going to perform, but there is a way to get a Hilbert structure instead of the Banach one. Note that s -th derivative of α -form changes as a $(\alpha + s)$ -form under coordinate transformations (in the main term), thus its square is a top-degree form on $X_{\mathbb{R}}$ (in the main term) if $\alpha + s = 1$. We conclude that the notion that *1/2-th derivative of a half-form is square-integrable* has a chance to be geometrizable. This notion leads us to consideration of $H^{1/2}$ -Sobolev spaces with values in (holomorphic) half-forms. (We discuss the basics of the theory of Sobolev spaces in Section 2) In fact taking different s one gets an entire hierarchy of Banach spaces $W_{\frac{2}{s+1/2}}^s$ which interpolate between L_4 and $H^{1/2}$. However, only one of these spaces is a Hilbert one, and having a Hilbert space is very convenient for our method of divide and conquer.

As we will see it in Section 2.7, the spaces of functions we obtain in such a way are closest possible analogues of Hardy spaces. Note also that they are very small

modifications of a particular representation of $\mathrm{SL}_2(\mathbb{C})$ from a supplementary series (see Section 2.6). Indeed, the supplementary series is realized in s -forms on $(\mathbb{CP}^1)_{\mathbb{R}}$, and the Hilbert structure in these spaces is equivalent to H^{1-2s} -structure. Taking $s = 1/4$, we see that the $H^{1/2}$ -structure on sections of $\omega^{1/4} \otimes \bar{\omega}^{1/4}$ may be defined⁶ to be invariant w.r.t. fraction-linear transformations. One should compare this result with the above heuristic about $H^{1/2}$ -structure on sections of $\omega^{1/2}$ being “invariant in the main term” w.r.t. diffeomorphisms (we use an additional heuristic that $\omega^{1/4} \otimes \bar{\omega}^{1/4}$ is “close” to $\omega^{1/2}$).

0.10. Thick infinity. We conclude that a working definition of *admissible* sections on a sphere with some disks removed is that the section is a half-form on the domain which can be extended as a $H^{1/2}$ -section of $\omega^{1/2}$ to the whole sphere (though a usual heuristic about $H^{1/2}$ is that it is a smoothness class, it restricts the growth as well).

The usual definition of H^s on a manifold with a boundary (see Section 2.2) allows one to work with function within the interior of the manifold only. In our case the smooth part of the boundary (union of circles) is not closed (if the number of circles is infinite), so one loses the info about what happens on the *dust*, i.e., in the accumulation points of the disks. This has some very inconvenient consequences. Let S be the sphere with interiors of disks removed, V be the closure of $\mathbb{CP}^1 \setminus S$. Then the sections we consider are elements of $H^{1/2}(\mathbb{CP}^1 \setminus V)$. The condition of such a φ being holomorphic can be written as $\bar{\partial}\varphi = 0 \in H^{-1/2}(\mathbb{CP}^1 \setminus V)$ (since taking derivative moves one notch down on Sobolev scale). However, if the dust $S \cap V$ is massive enough, there are functions of smoothness $H^{-1/2}$ with support in $S \cap V$, thus there functions φ of smoothness $H^{1/2}$ such that $\bar{\partial}\varphi$ has support on the dust only. Since $\bar{\partial}\varphi = 0 \in H^{-1/2}(\mathbb{CP}^1 \setminus V)$, such functions look like holomorphic ones, but in fact the dust is the part of infinity on our curve of infinite genus, thus one should consider the divisor of these functions as having components at infinity (since $\bar{\partial}\varphi$ is not zero there). This ruins our preparations to count the divisor at infinity correctly.

One possible way out of this deadlock is to prohibit collections of circles with a massive dust, say, restrict the Hausdorff dimension of the dust to be smaller than 1. We pick up a different approach: we massage the definition of the Sobolev space for a subset of a manifold in such a way that it now takes into account the behaviour on the boundary even if the boundary is massive. These are so called *generalized* Sobolev spaces. As a corollary, we can consider pretty monstrous curves of infinite genus.

For example, one can take a Sierpinsky carpet⁷ on a complex plane, and take one disk inside each thrown away triangle/rectangle. Gluing the boundaries of disk

⁶Recall that H^s -structure on sections of bundles on manifolds is defined only up to equivalence of topologies.

⁷Which is a two-dimensional analogue of the Cantor set, obtained by repeated removing of a smaller rectangle inside a bigger one (or removing a triangle with vertices on the sides of a bigger triangle).

pairwise gives a curve we can deal with (if radii of disks decrease quick enough). In particular, the dust can have a positive measure (in fact in our theory the only condition on the dust is that it is nowhere dense, so can be a set of accumulation points).

One can see that smooth points on the resulting curve have infinitely many connected components, and it is the dust which keeps these components together. The components are tubes, and the boundaries of these tubes are glued to the dust, which is connected. We see that points of the dust need to be considered as legitimate points on the curve of infinite genus, and the curve is *not a topological manifold* in neighborhoods of these points.

The striking fact is that it is possible to strengthen the restriction on the radii of the circles in such a way that any global holomorphic function has an *asymptotic* Taylor expansion near any point of the dust (see Section 6.8)! Thus the dust would consist of points which one has a full right to call *smooth points*.

0.11. Fight with $H^{1/2}$. While consideration of half-forms of smoothness $H^{1/2}$ *enormously* simplifies the work with infinities on the curve, it is one of the worst choices when we consider gluing conditions on sections on the glued together circles on the complex sphere. The reason is that Sobolev spaces with half-integer indices do not satisfy a lot of properties of other Sobolev spaces when restrictions to hypersurfaces is considered (this is the case when one usually considers Besov spaces instead of Sobolev ones).

To fight with this, we define a notion of *mollified restriction* to a hypersurface. Applying this restriction to the case of one removed disk $\{|z| > 1\}$ one can momentarily see that the space $H^{1/2}$ we consider coincides with the Hardy space $\mathcal{H}^2$ for the circle $\{|z| = 1\}$! We see that the space of “global holomorphic functions” we consider is a generalization of the Hardy space to the case when the boundary consists of many circles. What is more, the Hardy space $\mathcal{H}^\infty$ (of multipliers in $\mathcal{H}^2$) also plays an important rôle when we consider equivalence of bundles in Section 8.7.

0.12. Main results. Since technical details take a lot of place in the course of discussion, we make a concise list of main results (without discussing what the different conditions on the curve/bundle mean).

In Section 4.5 we prove (an analogue of) Riemann–Roch theorem for $\bar{\partial}: \mathcal{O} \rightarrow \bar{\omega}$ in assumption that the matrix $\mathbf{M}_2 = (e^{-l_{ij}} - \delta_{ij})$ (formed basing on conformal distances l_{ij} between disks) gives a compact mapping $l_2 \rightarrow l_2$. In Section 9.1 we show that the mapping $\bar{\partial}: \mathcal{O} \rightarrow \bar{\omega}$ has a maximal rank compatible with its index. In Section 9.2 we show an analogues result for $\bar{\partial}: \omega \rightarrow \omega \otimes \bar{\omega}$ under an additional assumption that disks have a *thickening* (in fact we show that the mapping of taking A -periods of global homomorphic forms is an isomorphisms). In Section 9.5 we describe the image of the (A, B) -period mapping. In all these cases the mappings in question have an infinite index, so we massage these mappings to get a mapping of index 0 (by extending the domain or target spaces) which is in fact an isomorphism.

In Section 7.3 we show that the space of global holomorphic sections of $\omega^{1/2} \otimes \mathcal{L}$ is finite-dimensional provided the matrix $\mathbf{M}_1 = (a_i e^{-l_{ij}/2} - a_i \delta_{ij})$: give a compact mapping $l_2 \rightarrow l_2$. Here (a_i) is some sequence associated to the bundle $\mathcal{L}$ (it consists of norms of gluing cocycle for $\mathcal{L}$). In the section 7.4 we show that the duality

$$\text{Coker } (\bar{\partial}: \omega^{1/2} \otimes \mathcal{L} \rightarrow \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega}) \simeq \text{Ker } (\bar{\partial}: \omega^{1/2} \otimes \mathcal{L}^{-1} \rightarrow \omega^{1/2} \otimes \mathcal{L}^{-1} \otimes \bar{\omega})$$

holds as far as $\mathbf{M}_1$ gives a bounded operator $l_2 \rightarrow l_2$.

In Section 7.5 we show that the above mappings satisfy the Riemann–Roch theorem as far as both $(a_i e^{-l_{ij}/2} - a_i \delta_{ij})$ and $(a'_i e^{-l_{ij}/2} - a'_i \delta_{ij})$ give compact operators $l_2 \rightarrow l_2$ (here a'_i is the sequence associated to $\mathcal{L}^{-1}$). In the same section we show that these mappings are Fredholm under an additional condition that (a_i) , (a'_i) are bounded.

In Section 7.6 we show that the above conditions give no restrictions on the *dust* of the corresponding Möbius group. In Section 7.7 we discuss heuristics which show that our version of the Riemann–Roch theorem is close to the strongest possible.

In Section 8.4 we show that an existence of so called *Hilbert–Schmidt* bundle on a curve shows that $\omega^{1/2}$ is also Hilbert–Schmidt. In Section 8.5 we show that the set of Hilbert–Schmidt bundles is convex (in the sense of multiplication of bundles), thus it is possible to define a *group of strong Hilbert–Schmidt* bundles. In Section 8.7 we define the notion of two bundles being *equivalent*, and define the Jacobian. In Section 8.8 we define divisors, and in Section 8.9 the mapping to the Universal Grassmannian (using asymptotic series at points of the dust defined in Section 6.8).

In Section 9.1 we show that if $\sum_{i \neq j} e^{-l_{ij}} < \infty$, then any admissible bundle of degree 0 can be described by locally-constant cocycles (and some amplifications of this result). In Section 9.6 we show that the period matrix is symmetric, and the imaginary part is positive. In Section 9.7 we show that the Jacobian coincides with the quotient of the space of possible periods by the lattice of *integer* periods, as far as $\sum_{i \neq j} e^{-l_{ij}} < \infty$. Finally, in Section 9.8 we show how to modify the notion of Hodge structure to describe the curves we study in this paper.

0.13. Main tools. We already noted that to deal with massive infinities we need the notion of generalized Sobolev space (introduced in Section 2.4), and to deal with gluing conditions along the boundaries of disks we need mollified restriction/extension mappings (defined in Section 5.3).

We discuss the relationship of the “invariant” growth conditions (we need them to have a good enumeration of divisor at infinity, see Section 0.8) with the supplementary series of representations of $\text{SL}_2(\mathbb{C})$ in Section 2.6. In Section 2.7 we demonstrate that the classical Hardy space $\mathcal{H}^2$ is a particular case of spaces we consider here (for “curves of genus $1/2$ ”, when the fundamental domain is a disk). In Section 8.7 we show that the space of multipliers (needed for defining equivalence of bundles) gives $\mathcal{H}^\infty$ in the particular case of a disk as a fundamental domain.

To handle divide-and-conquer strategy of dealing with curves of infinite genus we consider *almost perpendicular* families of subspaces (studied in Section 5.6). To study

duality we employ spaces of *strong sections* of bundles (see Section 6.5), to study $\bar{\partial}$ as a Fredholm operator we employ spaces of *weak sections* (see Section 5.6). (In most important cases these spaces coincide.)

To define B -periods on a curve we apply a method of averaging (Section 9.4) of paths connecting two sides of an A -cycle.

0.14. Historic remarks. In the spring of 1981 Yu. I. Manin was overwhelmed by unusual amount of sophomore students who asked him to be their undergraduate advisor. Contrary to his habits of the time, Yuri Ivanovich presented a list of topics he thought would be interesting and instructive for us to work on. One of these topics was a generalization of algebro-geometric description of solutions of KP to a curve of infinite genus represented by its Shottki model. (It was R. Ismailov who started to work on this topic.)

Five years later, when integrable systems reached the peak of their popularity, we obtained first results on the Riemann–Roch theorem which were similar to the results of this paper. The curves were represented in the way that was in fashion that time: as collections of pants glued along boundaries. The restrictions on the curves/bundles were jaw-breaking and very ad-hockish. Since Manin’s suggestions were completely forgotten already, there was nothing similar to the Shottki model approach. However, half-forms and $H^{1/2}$ -topology were already present at that time.

When in the end of the eighties the first variant of this paper was taking shape, it became clear that the conditions on curves/sheaves simplify *enormously* if one restricts attention to gluings of boundary circles which are fraction-linear, and gluing conditions for the bundles are given by locally constant cocycles (relative to half-forms). To our great surprise, with these restrictions the pants would glue together into the Shottki model of the curve (see Section 0.5), and sheaves glue into the bundle of half-forms on $\mathbb{CP}^1$. This finished a complete circle returning the settings back into context of the question of Manin’s of 1981!

Note that though the importance of Shottki model, of half-forms and of $H^{1/2}$ -growth conditions was clear long time ago, it was the formalization of the mollified restriction mapping which emphasized the capacity of infinity and the parallelism with the usual study of Hardy spaces $\mathcal{H}^2$ and $\mathcal{H}^\infty$. This formalization (together with the rest of this paper) would not appear if not the fruitful discussions with A. Tyurina in spring of 1996.

During the last several years Feldman, Knörrer and Trubowitz made a major breakthrough using an unrelated approach (cf. [3]).

The author is most grateful to I. M. Gelfand, A. Givental, A. Goncharov, D. Kazhdan, M. Kontsevich, Yu. I. Manin, H. McKean, V. Serganova, A. Tyurina and members of A. Morozov’s seminar for discussions which directed many approaches applied here. It was the patient work of V. Serganova which improved readability of the most obscure pieces of this paper.

This work was partially supported by NSF grant and Sloan foundation fellowship.

1. GEOMETRY OF HALF-FORMS

1.1. Half-forms and holomorphic half-forms. Consider an oriented real manifold M . For $\alpha \in \mathbb{C}$ consider a (complex) linear bundle Ω_M^α with transition functions $(\det \frac{\partial Y}{\partial X})^\alpha$ if $Y = Y(X)$. We will write a section of this bundle as $\varphi(X) dX^\alpha$, so $\varphi(X) dX^\alpha = \psi(Y) dY^\alpha$ if $Y = Y(X)$ and $\varphi(X) (\det \frac{\partial Y}{\partial X})^\alpha = \psi(Y)$. When M is clear from context, we will denote Ω_M^α by Ω^α .

In what follows we will be most interested in $\Omega^{1/2}$. Note that $\Omega^{1/2} \otimes \Omega^{1/2}$ is canonically isomorphic to the bundle of top de Rham forms. For any two global sections φ, ψ of $\Omega^{1/2}$ the product $\varphi\bar{\psi}$ is a top de Rham form, thus the pairing

$$(\varphi, \psi) \mapsto \int \varphi\bar{\psi}$$

gives a structure of pre-Hilbert space on $\Gamma(M, \Omega^{1/2})$. In particular, a notion of L_2 -section of $\Omega^{1/2}$ is canonically defined.

If M is a complex-analytic manifold, the power $(\det \frac{\partial Y}{\partial X})^\alpha$ is not uniquely defined (unless $\alpha \in \mathbb{Z}$), thus Ω^α is not canonically defined. We use an ad hoc definition of an analogue of $\Omega^{1/2}$ as follows:

Denote by ω the top holomorphic de Rham bundle for M . Consider a linear bundle l with an isomorphism $i: l \otimes l \simeq \omega$. We call sections of such a bundle *holomorphic half-forms*, and will denote $l = \omega^{1/2}$. In what follows we will consider at most one bundle of half-forms on a given manifold.

Note that any local diffeomorphism $M \rightarrow N$ induces a bundle of half-forms on M given a bundle of half-forms on N .

1.2. Half-forms on complex curves. In what follows we are most interested in complex curves. On a compact complex curve of genus g there are 2^g different bundles of half-forms. If $g = 0$, then there is a uniquely defined bundle of half-forms, isomorphic to $\mathcal{O}(-1)$.

To pick up an isomorphism of $\mathcal{O}(-1)^{\otimes 2} = \mathcal{O}(-2)$ with ω , one must fix an element $v \in \Lambda^2 V^*$, here V is 2-dimensional space, projectivization of which is $\mathbb{CP}^1$. Indeed, let $\mathbf{e}$ be the Euler vector field on V , $\mathbf{e} = z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2}$. Consider v as a constant 2-form on V , then $\mathbf{e} \lrcorner v$ is a 1-form on V which induces a global section of $\omega \otimes \mathcal{O}(2)$ on PV . Multiplication on this section gives an isomorphism of $\mathcal{O}(-2)$ and ω .

This shows that there should be an action of $\mathrm{SL}(2, \mathbb{C})$ on the bundle of half-forms, compatible with the action of $\mathrm{SL}(2, \mathbb{C})$ on $\mathbb{CP}^1$. Indeed, in affine coordinate z on $\mathbb{CP}^1 \setminus \{\infty\}$, a half-form may be written as $\varphi(z) dz^{1/2}$. Here if domain of φ contains ∞ , then $\varphi(\infty) = 0$. To write an action of $m = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2, \mathbb{C})$, note that the derivative of the corresponding action $z' = \frac{az+b}{cz+d}$ on $\mathbb{CP}^1$ has a canonical square root $\frac{1}{cz+d}$. Thus $\varphi(z') dz'^{1/2} \sim \varphi(z) \frac{1}{cz+d} dz^{1/2}$.

Note that it is the restriction of this action on $\mathrm{SL}(2, \mathbb{R})$ what is used to define automorphic forms on $\{\mathrm{Im} z > 0\}$. Note also that this action cannot be pushed down to an action of $\mathrm{PGL}(2, \mathbb{C})$.

1.3. Restriction onto S^1 . Consider now an annulus $B = \{r < |z| < R\}$ on a complex plane. Obviously, there are only two different bundles of half-forms on B . The inclusion of B into $\mathbb{CP}^1$ induces a bundle $\omega^{1/2}$ of half-forms on B . Another bundle of half-forms is $\omega^{1/2} \otimes \mu$, here μ is a locally constant sheaf with monodromy -1 on B , obviously μ^2 is the constant sheaf $\mathbb{C}$. Note that the holomorphic form $\frac{dz}{iz}$ on B has no square root in $\omega^{1/2}$, but has two in $\omega^{1/2} \otimes \mu$, one may write these square roots symbolically as $\pm \frac{(dz)^{1/2}}{i^{1/2} z^{1/2}}$.

Consider now a restriction of $\omega^{1/2} \otimes \mu$ onto $S^1 = \{|z| = 1\} \subset B$ (we assume that $r < 1 < R$). Consider a coordinate $t = \mathrm{Im} \ln z$ on S^1 , and identify $\frac{dz^{1/2}}{i^{1/2} z^{1/2}} \in \Gamma(S^1, \omega^{1/2} \otimes \mu)$ with $dt^{1/2} \in \Gamma(S^1, \Omega_{S^1}^{1/2})$. A simple calculation shows that this identification gives an isomorphism of $\omega^{1/2}|_{S^1}$ with $\Omega_{S^1}^{1/2} \otimes \mu$, which is preserved by real-analytic diffeomorphisms of S^1 with a given lift to μ .

In general, an immersion of S^1 in a complex curve X with a bundle of half-forms ω induces an isomorphism of $\omega^{1/2}|_{S^1}$ either with $\Omega_{S^1}^{1/2} \otimes \mu$, or with $\Omega_{S^1}^{1/2}$. One may call immersions of the first type A -cycles, of the second one B -cycles. Until Section 9 the immersions we consider are going to be A -cycles, thus we will always have a factor μ .

Note that the above isomorphism is defined up to multiplication by ± 1 , and both $\Omega_{S^1}^{1/2} \otimes \mu$ and $\Omega_{S^1}^{1/2}$ have natural pre-Hilbert structures, thus $\omega^{1/2}|_{S^1}$ has a natural pre-Hilbert structure as well.

2. SOBOLEV SPACES

In what follows $\mathcal{S}$ denotes the space of rapidly decreasing smooth functions, $\mathcal{D}$ denotes the space of smooth functions with compact support, and $\mathcal{S}'$ and $\mathcal{D}'$ are the dual spaces. l_2 denotes the Hilbert space of square-integrable sequences, and L_2 denotes the space of square-integrable functions. The symbol $\bigoplus V_i$ denotes the space of sequences with finite number of non-zero terms. If V_i are Hilbert spaces, $\bigoplus_{l_2} V_i$ denotes the space of square-integrable sequences (v_i) with $v_i \in V_i$.

The material of this section is mostly standard, however, note that we also discuss some topics which are not parts of standard curriculum of Sobolev spaces: in Section 2.4 we introduce *generalized* Sobolev spaces, in Sections 2.6 and 2.7 we give descriptions of supplementary series of representations of $\mathrm{SL}(2, \mathbb{C})$ and of Hardy space in terms of Sobolev spaces. In addition to the results which are covered in this section, in Section 5.2 we introduce *mollifications* of mappings of restriction to a submanifold (in the case $s = 1/2$), and of mapping of extension from submanifold (in the case $s = 0$), both in the cases when the non-mollified mappings are not continuous.

2.1. Euclidean case. Consider a vector space $\mathbb{R}^n$ with the standard Euclidean structure. Consider $s \in \mathbb{R}$ and a norm $\|\bullet\|_s$ on the set $\mathcal{S}$ of rapidly decreasing C^∞ -functions:

$$\|f(x)\|_s = \int (1 + |\xi|^2)^s |\widehat{f}(\xi)|^2 d\xi,$$

here $\widehat{f}(\xi)$ is the Fourier transform of f . By definition, the Sobolev space $H^s(\mathbb{R}^n)$ is the completion of $\mathcal{S}$ w.r.t. this norm. It is naturally isomorphic to the set of L_2 -functions $g(\xi)$ w.r.t. the measure $(1 + |\xi|^2)^{s/2} d\xi$. Since the inverse Fourier transform of $g(\xi)$ is a well-defined generalized function on $\mathbb{R}^n$, there is a natural inclusion $H^s(\mathbb{R}^n) \hookrightarrow \mathcal{S}'$, compatible with the inclusion $\mathcal{S} \hookrightarrow H^s(\mathbb{R}^n)$.

The pairing

$$(f, g) \mapsto \int f \bar{g} dx$$

extends to a pairing $H^s(\mathbb{R}^n) \otimes H^{-s}(\mathbb{R}^n) \rightarrow \mathbb{C}$ which is a pairing of Hilbert spaces. This pairing is compatible with the pairing between $\mathcal{S}$ and $\mathcal{S}'$.

Let D be a closed subset of $\mathbb{R}^n$, U be an open subset of $\mathbb{R}^n$. Define a Hilbert subspace

$$\mathring{H}^s(D) = H^s(\mathbb{R}^n) \cap \{f \in \mathcal{S}' \mid \text{Supp } f \subset D\},$$

and a quotient Hilbert space

$$H^s(U) = H^s(\mathbb{R}^n) / \mathring{H}^s(\mathbb{R}^n \setminus U).$$

The following properties of Sobolev spaces are most important in analysis:

1. If D is compact, V, U are open, $V \subset D \subset U \subset \mathbb{R}^n$, and $\varphi: U \rightarrow \varphi(U)$ is a diffeomorphism, then φ^* induces invertible bounded operators $\mathring{H}^s(\varphi(D)) \rightarrow \mathring{H}^s(D)$, $H^s(V) \rightarrow H^s(U)$.
2. If D is compact, then any differential operator A of degree d with smooth coefficients gives a bounded operators $\mathring{H}^s(D) \rightarrow \mathring{H}^{s-d}(D)$, $H^s(V) \rightarrow H^{s-d}(V)$.
3. In the same way a pseudodifferential operator of degree d gives a bounded operators $\mathring{H}^s(D) \rightarrow H^{s-d}(U)$.
4. The mapping of restriction of smooth functions onto $\mathbb{R}^{n-k} \subset \mathbb{R}^n$ extends to a bounded operator

$$r: H^s(\mathbb{R}^n) \rightarrow H^{s-\frac{k}{2}}(\mathbb{R}^{n-k}) \text{ if } s > \frac{k}{2}.$$

5. Dually, extension-by- δ -function of generalized functions on $\mathbb{R}^{n-k}$ to generalized functions on $\mathbb{R}^n$ gives a bounded operator $e: H^s(\mathbb{R}^{n-k}) \rightarrow H^{s-\frac{k}{2}}(\mathbb{R}^n)$ if $s < 0$.
6. If D is compact and has a smooth boundary, then $\mathring{H}^s(D)^\perp \subset H^{-s}(\mathbb{R}^n)$ coincides with $\mathring{H}^{-s}(\overline{\mathbb{R}^n \setminus D})$.

2.2. Sobolev spaces on manifolds. If M is a paracompact C^∞ -manifold, then $H^s(M)$ can be defined as a subspace of generalized functions on M which consists of functions f such that for any $U \subset M$ and $\varphi: \tilde{U} \xrightarrow{\sim} \tilde{V} \subset \mathbb{R}^n$ the restriction $f|_{\tilde{U}}$ satisfies $(\varphi^{-1})^* f \in H^s(V)$ (here $\tilde{U}$ is a neighborhood of $\bar{U}$, $V = \varphi(U)$).

An alternative definition is that for an appropriate partition of unity $(U_\alpha, \sigma_\alpha)$, $\text{Supp } \sigma_\alpha \subset\subset U_\alpha$, $\sum_\alpha \sigma_\alpha = 1$, the products $\sigma_\alpha f$ satisfy $(\varphi_\alpha^{-1})^* (\sigma_\alpha f) \in \mathring{H}^s(V_\alpha)$, here $\varphi_\alpha: U_\alpha \xrightarrow{\sim} V_\alpha \subset \mathbb{R}^n$.

In the same way one may define H^s -sections of C^∞ -vector bundles over M . We denote the space of H^s -section of a bundle E by $H^s(M, E)$. Obviously, if M is compact, one may define a structure of Hilbert space on $H^s(M)$, but this structure is not uniquely defined. However, the corresponding topology on $H^s(M)$ is well-defined. For a general manifold M one defines a topology on $H^s(M)$ as an inverse limit w.r.t. topologies on $H^s(U)$, U being open subsets of M diffeomorphic to bounded open subsets of $\mathbb{R}^n$.

If D is a closed subset of M , and U is an open subset, define $\mathring{H}^s(D) \subset H^s(M)$ as subspace of function with support in D , and $H^s(U)$ as $H^s(M) / \mathring{H}^s(M \setminus U)$.

Note that if $M = \mathbb{R}^n$, the above definition produces a different space $H_{\text{loc}}^s(\mathbb{R}^n)$ of functions on $\mathbb{R}^n$ than the space $H^s(\mathbb{R}^n)$ defined in Section 2.1. A generalized function f on $\mathbb{R}^n$ is in $H_{\text{loc}}^s(\mathbb{R}^n)$ if in any bounded domain it is equal to a function from $H^s(\mathbb{R}^n)$.

The properties of Sobolev spaces $H^s(\mathbb{R}^n)$ have direct analogues for $H^s(M)$, thus diffeomorphisms of manifolds and differential operators act on $H^s(M)$, and one can restrict/extend Sobolev sections to/from submanifolds. In particular, the existence of a parametrix for an elliptic differential operator shows that on a compact manifold M an elliptic operator A of degree d gives a Fredholm operator $H^s(M) \rightarrow H^{s-d}(M)$ (with suitable changes if A acts in vector bundles).

Note that on compact manifolds one can define the Sobolev spaces using arbitrary elliptic (pseudo)differential operators:

Proposition 2.1. *Consider a compact manifold M with a metric and a positive self-adjoint elliptic operator A of degree d . Then the s -Sobolev norm is equivalent to the norm*

$$\|f\| = \int_M |(1 + A)^{s/d} f|^2 d\mu.$$

We will also use the following statement:

Proposition 2.2. *Consider two linear bundles $\mathcal{L}_1$ and $\mathcal{L}_2$ on a compact manifold M such that $\mathcal{L}_1 \otimes \mathcal{L}_2 \simeq \Omega^{\text{top}}(M)$. Then the pairing $\int \alpha\beta$ between smooth sections of $\mathcal{L}_1$ and $\mathcal{L}_2$ can be extended to non-degenerate pairing between Hilbert spaces $H^s(M, \mathcal{L}_1)$ and $H^{-s}(M, \mathcal{L}_2)$ for an arbitrary $s \in \mathbb{R}$.*

2.3. Capacity. Consider a manifold M . We say that s -capacity of a closed subset $S \subset M$ is non-zero, if there is a non-zero function $f \in H^s(M)$ such that $\text{Supp } f \subset S$. We say that S has a *capacity dimension* $\geq d$, if s -capacity of S is non-zero for $s = -\frac{\dim M - d}{2}$.

Since δ -function of a point belongs to $H^s(M)$ with $s < -\frac{\dim M}{2}$, a capacity dimension of a point is $\geq d$ if $d < 0$. It is clear that this estimate cannot be improved. In the same way the capacity dimension of a submanifold S is $\geq d$ iff $d < \dim S$.

For a general subset S the same is true if one considers Hausdorff dimension [6].

2.4. Generalized Sobolev spaces. In Section 6.1 we consider the space of $H^{1/2}$ -half-forms which are holomorphic outside of a closure $\bar{U}$ of a union U of disks in $\mathbb{CP}^1$. We will see that this space is too big for our purposes if $\bar{U}$ is “much bigger” than U . We will need to consider half-forms which “are holomorphic” in $\bar{U} \setminus U$ as well as in $\mathbb{CP}^1 \setminus \bar{U}$.

To have this we need the half-forms to be *defined* in $\mathbb{CP}^1 \setminus U$ instead of $\mathbb{CP}^1 \setminus \bar{U}$. The problem with this is that U is not closed, so the usual definition of $H^{1/2}$ -half-forms as of a quotient-space does not work. This shows the need for the following

Definition 2.3. Consider a subset U of the manifold M . Let $\mathring{H}^s(U)$ denotes the closure of the subspace $L \subset H^s(M)$

$$L = \{f \in H^s(M) \mid \text{Supp } f \subset U\}.$$

Consider $V \subset M$. Let $H^s(V) = H^s(M) / \mathring{H}^s(M \setminus V)$.

Note that we do not require that the subset U is closed, and V is open. If they are, then we get the standard definitions of $\mathring{H}^s$ and H^s . Obviously, $\mathring{H}^s(U) \subset \mathring{H}^s(\bar{U})$, but this inclusion may be proper, as the following construction shows⁸.

Consider a disjoint family of closed subsets $V_i \subset M$, Let $\mathcal{V} = \overline{\bigcup V_i} \setminus \bigcup V_i$. Suppose that

1. The natural mapping $\bigoplus_{i_2} \mathring{H}^s(V_i) \xrightarrow{\iota} H^s(M)$ is a (continuous) injection⁹.
2. The s -capacity of $\mathcal{V}$ is positive.

The first condition insures that the space $\mathring{H}^s(\bigcup V_i)$ is the image of the mapping ι . Hence any non-zero function $f \in \mathring{H}^s(\bigcup V_i)$ satisfies the condition $\text{Supp } f \cap \bigcup V_i \neq \emptyset$.

On the other hand, the second condition shows that there is a non-zero function $f \in H^s(M)$ such that $\text{Supp } f \subset \mathcal{V}$. Obviously, $f \in \mathring{H}^s(\overline{\bigcup V_i})$, but $f \notin \mathring{H}^s(\bigcup V_i)$.

In Section 7.6 we show how to construct a family of disks V_i which satisfy the first condition. The centers of these disks may be an arbitrary prescribed locally discrete set. Moreover, one can easily find an appropriate set of centers such that the

⁸The construction can be simplified, but in the current form it is an example of domains we are going to deal with.

⁹I.e., the image is closed, and the mapping is an isomorphism on the image.

corresponding set $\mathcal{V}$ does not depend on radii and coincides with an arbitrary given closed set $\mathcal{V}_0$ with empty interior.

In particular, if there exist a closed set $\mathcal{V}$ with an empty interior and non-zero s -capacity, then one can construct a subset U of $\mathbb{C}P^1$ such that the inclusion $\mathring{H}^s(U) \subset \mathring{H}^s(\bar{U})$ is proper. Taking appropriate Cantor sets, one gets that this is possible with any $s < 0$.

On the other hand, if U has smooth boundary, then $\mathring{H}^s(U) = \mathring{H}^s(\bar{U})$.

2.5. Rescaling on $\mathbb{R}^n$ and $\mathbb{C}^n$. Consider an H^s -function $f(x)$ on $\mathbb{R}^n$ with compact support. The Fourier transform $\widehat{f}(\xi)$ is real-analytic, thus the integral

$$(2.1) \quad \int |\xi|^{2s} |\widehat{f}(\xi)|^2 d\xi,$$

can be defined in the sense of generalized functions (i.e., analytic continuation in s) near $\xi = 0$. Since $f \in H^s$, the integral converges near ∞ . Moreover, if $s > -\frac{n}{2}$, the integral converges near $\xi = 0$, thus its value is a limit of Riemann sums, hence is non-negative.

In any case the integral defines a quadratic form on $\mathring{H}^s(D)$ for any compact D .

Fact . *If $s > -\frac{n}{2}$, the integral defines a norm on $\mathring{H}^s(D)$, and this form is equivalent to the Hilbert norm.*

The advantage of the norm (2.1) is the fact that it is covariant with respect to dilatations. Define a mapping $\mathfrak{D}_a^s: \mathcal{S} \rightarrow \mathcal{S}$ by

$$(\mathfrak{D}_a^s f)(x) = a^s f(ax), \quad a, s \in \mathbb{R}, \quad a \neq 0.$$

Then $\mathfrak{D}_a^{-s+\frac{n}{2}}$ is an isometry w.r.t. the norm (2.1). Note also that that $\mathfrak{D}_a^{tn}$ is a natural action of a dilatation on sections of the linear bundle Ω^t .

Combining this with the previous statement, we conclude that any similarity transform¹⁰ $\mathbf{T}$ acts as a uniformly bounded operator in the space of H^s -sections of Ω^α

$$\mathbf{T}^*: \mathring{H}^s(\mathbf{T}D, \Omega^\alpha) \rightarrow \mathring{H}^s(D, \Omega^\alpha), \quad \alpha = \frac{1}{2} - \frac{s}{n},$$

as far as both D and $\mathbf{T}D$ remain in the same disk $\{|x| < R\}$, and R is fixed.

If we consider $\mathbb{C}^n = \mathbb{R}^{2n}$ and the linear bundle $\omega^{1/2}$ over $\mathbb{C}^n$, then the action of dilatations¹¹ on this bundle is the same¹² as on $\Omega_{\mathbb{R}^{2n}}^{1/4}$. We conclude that similarity transforms act as uniformly bounded transformations

$$\mathring{H}^{n/2}(\mathbf{T}D, \omega^{1/2}) \rightarrow \mathring{H}^{n/2}(D, \omega^{1/2})$$

as far as both D and $\mathbf{T}D$ remain in the same disk $\{|z| < R\}$, and R is fixed.

¹⁰I.e., a composition of translations, rotations and dilatations.

¹¹In fact one needs to consider 2-sheeted covering of the group of similarity transforms.

¹²Strictly speaking, differs on a multiplication by a constant of magnitude 1.

2.6. **PGL(2, $\mathbb{C}$)-invariant realization of $H^{1/2}(\mathbb{CP}^1)$.** Consider double ratio

$$(a : b : c : d) = \frac{a - b}{b - d} : \frac{a - c}{c - d}, \quad a, b, c, d \in \mathbb{CP}^1.$$

Double ratio $(z_1 : z_1 + \delta z_1 : z_2 : z_2 + \delta z_2)$ gives a section ρ of the bundle $T^*\mathbb{CP}^1 \boxtimes T^*\mathbb{CP}^1$ over $\mathbb{CP}^1 \times \mathbb{CP}^1$. In affine coordinates it can be written as $\frac{1}{(z_1 - z_2)^2} dz_1 dz_2$. By construction this section is invariant with respect to the action of $\text{PGL}(2, \mathbb{C})$.

Consider $K_s = \frac{\rho^s \bar{\rho}^s}{\Gamma(-2s+1)}$. It is a section of $\Omega^s(\mathbb{CP}_{\mathbb{R}}^1) \boxtimes \Omega^s(\mathbb{CP}_{\mathbb{R}}^1)$, which in local coordinates looks like $K_s(z_1, z_2) = \frac{1}{|z_1 - z_2|^{4s} \Gamma(-2s+1)} dZ_1^s dZ_2^s$ and has no singularity outside of $z_1 = z_2$. Here $dZ = dx \wedge dy$ if $z = x + iy$. The operator with kernel K_s defines a pairing

$$(\alpha, \beta)_s = \int_{\mathbb{CP}^1 \times \mathbb{CP}^1} K_s(z_1, z_2) \alpha(z_1) \bar{\beta}(z_2)$$

on the space $\Gamma(\Omega^{1-s}(\mathbb{CP}_{\mathbb{R}}^1))$ if $s < \frac{1}{2}$, this pairing depends on s analytically [4], and may be continued to an arbitrary s .

Proposition 2.4. $(\cdot, \cdot)_s$ is a positive-definite Hermitian form on $\Gamma(\Omega^{1-s}(\mathbb{CP}_{\mathbb{R}}^1))$, if $|s - \frac{1}{2}| < \frac{1}{2}$. It is equivalent to the H^{-1+2s} -norm on this space.

Proof. For $s = \frac{1}{2}$ we get a standard pairing on $\Omega^{1/2}$ (since $K_{1/2}$ is a δ -function), so it is sufficient to consider $s \neq \frac{1}{2}$. Take an affine coordinate system z on $\mathbb{CP}^1 \setminus \{\infty\}$. A smooth section of Ω^t near ∞ is represented by a smooth t -form $f(z) dZ^t$ with an asymptotic

$$|z|^{-4t} g(1/z) dZ^t,$$

near ∞ . Here $g(w)$ is a smooth function of x and y , $w = x + yi$. Thus the Fourier transform $\hat{f}(\zeta)$ of f is rapidly decreasing, smooth outside of $\zeta = 0$, and has an asymptotic expansion

$$h(\zeta) + |\zeta|^{2-4t} h_1(\zeta) + O(|\zeta|^N)$$

near 0, here $N > 0$ is an arbitrary integer, h and h_1 are smooth functions, and $t \neq \frac{1}{2}$.

The pairing $(\cdot, \cdot)_s$ written in terms of $\hat{\alpha}, \hat{\beta}$ is the pairing

$$(2.2) \quad \text{const} \cdot \int \hat{\alpha}(\zeta) \bar{\hat{\beta}}(\zeta) |\zeta|^{4s-2} d\zeta d\bar{\zeta}$$

(in the sense of generalized functions) and we see that for $t = 1 - s$ and $|s - \frac{1}{2}| < \frac{1}{2}$ the integral converges, thus the pairing is positive.

On the other hand, the local equivalence of the norm (2.2) with H^{2s-1} is already proved in the previous section (cf. Equation (2.1)). $\square$

Remark 2.5. Note that what we got is a complementary series of unitary representations of $\text{SL}(2, \mathbb{C})$.

Remark 2.6. The defined pairings are compatible with duality $\int \alpha\beta$ between Ω^s and Ω^{1-s} . Note also that in the case $s = 0$ the pairing has $\text{rk} = 1$, dually, in the case $s = 1$ the pairing becomes $-(\Delta\alpha, \beta)$, which has constants in the null-space.

Put $s = \frac{1}{4}$. We get a $\text{PGL}(2, \mathbb{C})$ -invariant implementation¹³ of the space $H^{-1/2}(\Omega^{3/4})$. Note now that $\Omega^{3/4} = \omega^{3/4} \otimes \bar{\omega}^{3/4}$ differs not so much from the space $\omega^{1/2} \otimes \bar{\omega}$ we are most interested in. The difference $\omega^{1/4} \otimes \bar{\omega}^{-1/4}$ is a bundle with transitions functions $(\frac{D}{\bar{D}})^{1/4}$ of magnitude 1 (here $D = dz/dw$). In particular, a choice of affine coordinate z on $\mathbb{CP}^1$ gives an identification of $\omega^{1/2} \otimes \bar{\omega}$ with $\Omega^{3/4}$ (outside of infinity). This identification commutes with natural actions of $\text{Aff}(\mathbb{C})$ on $\Omega^{3/4}$ and $\omega^{1/2} \otimes \bar{\omega}$ (up to multiplication by a constant of magnitude 1). Here $\text{Aff}(\mathbb{C})$ is the group of affine transformations on $\mathbb{C}$ (in fact to get an action on $\omega^{1/2} \otimes \bar{\omega}$ one needs to consider 2-covering of $\text{Aff}(\mathbb{C})$).

This explains the almost-invariance of $H^{-1/2}(\mathbb{C}, \omega^{1/2} \otimes \bar{\omega})$ with respect to affine transformations, discussed in Section 2.5. Indeed, consider a disk $K \subset \mathbb{CP}^1$. On $\mathbb{CP}^1 \setminus K$ multiplication by $dz^{1/4}d\bar{z}^{-1/4}$ gives an isomorphism of the sheaves $\omega^{1/2} \otimes \bar{\omega}$ and $\Omega^{3/4}$. Taking ∞ as a center of K , we obtain the results of Section 2.5 in the case $n = 1$.

In what follows the following result is sufficient for us to show conformal invariance of the objects we introduce:

Amplification 2.7. *Fix a metric on $\mathbb{CP}^1$ and $\varepsilon > 0$. Consider $\varphi \in \text{SL}(2, \mathbb{C})$, let $A \subset \mathbb{CP}^1$, $B = \varphi(A)$. If both $\mathbb{CP}^1 \setminus A$ and $\mathbb{CP}^1 \setminus B$ contain disks of radius ε with centers at c_A, c_B , then the norm of $\varphi^*: \mathring{H}^{-1/2}(B, \omega^{1/2} \otimes \bar{\omega}) \rightarrow \mathring{H}^{-1/2}(A, \omega^{1/2} \otimes \bar{\omega})$ is bounded by*

$$C(\varepsilon) \left(\frac{\text{diam}(A)}{\text{dist}(\varphi^{-1}(c_B), A)} \right)^{1/2}.$$

By duality, the same bound is valid for action of φ^ in $H^{1/2}(\bullet, \omega^{1/2})$.*

Proof. Indeed, $SU(2)$ is compact, thus acts on $H^{1/2}(\mathbb{CP}^1, \omega^{1/2} \otimes \bar{\omega})$ by uniformly bounded operators. Thus it is sufficient to consider the case when a disk K of radius ε is fixed (say, if $c_A = c_B = \infty$, thus $|z| > 1/\varepsilon$), and $A \cap K = B \cap K = \emptyset$.

Let α, β be sections of $\omega^{1/4} \otimes \bar{\omega}^{-1/4}$ and $\omega^{-1/4} \otimes \bar{\omega}^{1/4}$ which coincide with $\bar{d}z^{1/4}d\bar{z}^{-1/4}$ and $dz^{-1/4}d\bar{z}^{1/4}$ inside $\mathbb{CP}^1 \setminus K$ correspondingly, and have a compact support. Then $\varphi^*|_A f$, $f \in \omega^{1/2} \otimes \bar{\omega}$, is equal to

$$\mu_\varphi \beta \varphi_{3/4}^*|_A(\alpha f), \quad \mu_\varphi = \left(\frac{\varphi'}{|\varphi'|} \right)^{1/2}$$

¹³I.e., a Hilbert structure which is equivalent to the Sobolev Hilbert structure which is in turn defined up to equivalence (compare with Section 2.2).

(here $\varphi_{3/4}^*$ acts on $\Omega^{3/4}$). On the other hand, multiplications by α and β are bounded operators, and $\varphi_{3/4}^*$ is unitary, so what remains to prove is a bound on operator of multiplication by μ_φ in $H^{-1/2}$ or in $H^{1/2}$. It is sufficient to consider $H^{1/2}$.

Let us apply interpolation theorem now: the norm of multiplication by μ_φ in $H^{1/2}$ is no more than geometric mean of norms in $H^0 = L_2$ and in H^1 . Since $|\mu_\varphi| = 1$, it is unitary in H^0 , so it is sufficient to estimate the norm in H^1 , which is bounded by $\text{const} \cdot \max |\mu'_\varphi|$, i.e., by $\text{const} \cdot \text{dist}(\varphi^{-1}(\infty), A)^{-1}$.

To get the estimate in the theorem, note that we have a freedom of rescaling A , thus may assume that $\text{diam}(A) = 1$. $\square$

2.7. Hardy space. Let $K = \{z \mid |z| \leq 1\}$. The Hardy space $\mathcal{H}$ is the subspace of $L_2(\partial K)$ consisting of functions with Fourier coefficients (a_n) which vanish for $n < 0$. For a function $f \in \mathcal{H}$, $f = \sum a_k z^k$, $|z| = 1$, let $c_f = \sum a_k z^k$ be defined for $|z| < 1$. The latter series converges, and defines a holomorphic function inside K .

Lemma 2.8. *The mapping $f \mapsto c_f$ is an injection $L_2(\partial K) \rightarrow H^{1/2}(K)$. The image of this injection coincides with*

$$\text{Ker} \left(H^{1/2}(K) \xrightarrow{\bar{\partial}} H^{-1/2}(K) \right).$$

Proof. It is sufficient to prove that $\|z^k\|_{H^{1/2}(K)}$ is bounded from above and from below when $k \in \mathbb{N}$. Consider a concentric disk K_1 of radius $R < 1$. Since $\|z^k\|_{H^{1/2}(K_1)} \leq \|z^k\|_{H^1(K_1)} = O(kR^k)$, it is $o(1)$, thus it is sufficient to consider the norm of z^k in a narrow annulus with external boundary ∂K .

In turn, taking coordinate $\log z = a + ib$, $b \in \mathbb{R}/2\pi\mathbb{Z}$, it is sufficient to consider $H^{1/2}$ -norm of e^{ka} on the half-line $a \leq 0$, more precise, it is sufficient to consider L_2 -norm of $(\frac{d}{da} + k)^{1/2} e^{ka}$, $a \leq 0$. Indeed, we need to show that there is a continuation of $e^{k(a+ib)}$ outside of $a \leq 0$ such that the $H^{1/2}$ -norm of this continuation is bounded, and that the $H^{1/2}$ -norm of any such continuation is bounded from below. One can suppose that the continuation is $e^{ikb}\varphi(a)$, so one needs to estimate $\int (1 + k^2 + \alpha^2) |\widehat{\varphi}(\alpha)|^2 d\alpha$. The operator $(\frac{d}{da} + k)^{1/2}$ maps this norm to a norm equivalent to L_2 -norm, moreover, it sends functions with support in $a \geq 0$ into itself (since $k \geq 0$). Thus it defines an invertible operator $H^{1/2}(\mathbb{R}_{\leq 0}) \rightarrow H^0(\mathbb{R}_{\leq 0}) = L_2(\mathbb{R}_{\leq 0})$.

This means that $(\frac{d}{da} + k)^{1/2} e^{ka}|_{a \leq 0} = (\frac{d}{da} + k)^{1/2} \varphi(a)|_{a \leq 0}$, thus we should not care about the choice of continuation φ . Moreover, since $(\frac{d}{da} + k)^{1/2}$ does not move support to the left, $(\frac{d}{da} + k)^{1/2} e^{ka} = C(k) e^{ka}$, $a \leq 0$. Since $(\frac{d}{da} + k)^{1/2}$ is an operator of convolution with $\frac{a_+^{-3/2} e^{-ka}}{\Gamma(-1/2)}$ (in the sense of generalized functions, x_+^s is 0 if $x < 0$, x^s if $x > 0$ and $s > 0$, and is defined by analytic continuation from the region $s > 0$, where the above description defines it completely [4]), to estimate $C(k)$ one needs to

calculate

$$\begin{aligned} - \int_0^\infty x_+^{-3/2} e^{-kx} e^{-kx} dx &\stackrel{\text{def}}{=} - \int_0^\infty x^{-3/2} (e^{-2kx} - 1) dx \\ &= k^{1/2} \int_0^\infty (kx)^{-1/2} \frac{1 - e^{-2kx}}{kx} d(kx) = k^{1/2} C. \end{aligned}$$

Thus we need to show that $k^{1/2} \|e^{ka}\|_{a \leq 0} \|_{L_2}$ is bounded from above and from below, which is obvious. $\square$

Remark 2.9. It is clear that if $f \in \mathcal{H} \cap C^\infty$, then $f = c_f|_{\partial K}$. Moreover, if $f \cdot \delta_{\partial K}$ is the continuation of f to $\mathbb{C}$ by δ -function, then $c_f = \text{const} \cdot \bar{\partial}^{-1}(f \cdot \delta_{\partial K})$ (see Section 3.3 for description of $\bar{\partial}^{-1}$). Several following sections are dedicated to defining something similar in the case $f \in \mathcal{H}$.

Remark 2.10. In Section 6 we will construct a generalization of the following modification of this result: instead of K we consider an isomorphic domain $\mathbb{CP}^1 \setminus K$. Instead of functions on $\mathbb{CP}^1$ and ∂K we consider sections of $\omega^{1/2}$ and $\Omega^{1/2} \otimes \mu$ correspondingly.

3. CAUCHY KERNEL FOR $\omega^{1/2}$

3.1. $\bar{\partial}$ on piecewise-analytic functions. Let M be a complex curve with a linear bundle $\mathcal{L}$. Consider a domain $D \subset M$ with a smooth boundary $\partial D = \gamma$. Let $f(z)$, $z \in \bar{D}$, be an analytic section of $\mathcal{L}$ in D which continues as a smooth section to $\bar{D}$. Extend f outside of D as 0.

Consider $\bar{\partial}f$. It is a section of $\mathcal{L} \otimes \bar{\omega}$, which vanishes on $M \setminus \gamma$. Since f has a jump of the first kind along γ , and $\bar{\partial}$ is a differential operator of the first order, $\bar{\partial}f$ has at most a δ -function singularity along γ . One may locally write

$$\bar{\partial}f = g(z) \cdot \delta_{h(z)=0}.$$

Here $g(z)$ is a smooth section of $\mathcal{L} \otimes \bar{\omega}$, h is a local equation of γ .

Let us calculate $g|_\gamma$ in terms of f . First of all, if one changes the equation $h = 0$ by a different equation $hh_0 = 0$ (here h_0 has no zeros close to the point in question), the coefficient at $g(z)$ changes to

$$\delta_{hh_0=0} = \frac{1}{h_0} \delta_{h=0},$$

since δ -function has homogeneity degree -1 . Thus $g(z)|_\gamma$ should be multiplied by $h_0|_\gamma$ to preserve the same value of the product. This shows that to get an invariant description of $g(z)|_\gamma$ one needs to write

$$g(z)|_\gamma = G(z) \otimes dh|_\gamma.$$

Here $dh|_\gamma$ is considered as a section of the conormal bundle $N^*\gamma$ to γ , thus $G(z)$ should be a section of $(\mathcal{L} \otimes \bar{\omega})|_\gamma \otimes (N^*\gamma)^* = (\mathcal{L} \otimes \bar{\omega})|_\gamma \otimes N\gamma$.

Now $G(z)$ does not depend on the parameterization of a neighborhood of γ , so it should be directly expressible in terms of $f(z)$.

Lemma 3.1. $\bar{\omega}|_\gamma$ is canonically isomorphic to $(N^*\gamma) \otimes \mathbb{C}$.

Proof. Indeed, $\bar{\omega}$ is defined as a quotient of $(T^*M_{\mathbb{R}}) \otimes \mathbb{C}$ by holomorphic forms, moreover, $\bar{\omega}_{\mathbb{R}}$ is an isomorphic image of $T^*M_{\mathbb{R}}$. Since $N^*\gamma \subset T^*M_{\mathbb{R}}$, there is a natural non-zero mapping $N^*\gamma \rightarrow \bar{\omega}_{\mathbb{R}}$, which gives the required isomorphism after complexification. $\square$

As a corollary, $(\mathcal{L} \otimes \bar{\omega})|_\gamma \otimes N\gamma \simeq \mathcal{L}|_\gamma$. Now the following fact is obvious from calculations in local coordinates:

Proposition 3.2. $G(z) = -f(z)|_\gamma$.

In particular, we see that one can write

$$(3.1) \quad \bar{\partial}f = -e \circ r(f)$$

in terms of operators r of restriction and e of extension-by- δ -function. Thus we can consider $-e \circ r$ as an “approximation” to $\bar{\partial}$ which is good on functions which are analytic far from γ . This approximation is exact on functions with a jump of the first kind.

Yet another way to treat this identity is to write it as

$$-\bar{\partial}^{-1} \circ e \circ r(F) = \vartheta_D F.$$

Here F is a function which is holomorphic in a neighbourhood of $\bar{D}$, ϑ_D is a function which is 1 on D and 0 outside of D . We give a mollified version of this statement in Section 5.5.

3.2. Self-duality of $\bar{\partial}$. Consider a mapping $\bar{\partial}: \omega^{1/2} \rightarrow \omega^{1/2} \otimes \bar{\omega}$. Note that the spaces of sections of these bundles are dual w.r.t. the pairing

$$(3.2) \quad (\varphi, \psi) = \int_M \varphi \psi.$$

Lemma 3.3. $\bar{\partial}$ is skew-symmetric w.r.t. the above pairing,

$$(\varphi_1, \bar{\partial}\varphi_2) + (\varphi_2, \bar{\partial}\varphi_1) = 0.$$

Proof. It is sufficient to show that $(\varphi_1, \bar{\partial}\varphi_1) = 0$, i.e., to study

$$\int_M \varphi \bar{\partial}\varphi = \frac{1}{2} \int_M \bar{\partial}\varphi^2 = \frac{1}{2} \int_M d\varphi^2$$

which obviously vanishes. $\square$

Note that this supports the heuristic that the bundle $\omega^{1/2}$ is “the best one” of the powers of ω .

Used literally, the above considerations were applicable to the space of smooth sections of the bundles in question. On the other hand, the pairing (3.2) extends to a continuous pairing between $H^{1/2}(M, \omega^{1/2})$ and $H^{-1/2}(M, \omega^{1/2} \otimes \bar{\omega})$, and the operator $\bar{\partial}$ maps one of these spaces to another. Using the facts from Section 2.2 We get a

Proposition 3.4. *The operator $\bar{\partial}$ gives a canonically defined Fredholm symplectic form on $H^{1/2}(M, \omega^{1/2})$. This form is non-degenerate if $M = \mathbb{CP}^1$.*

Here we call a bilinear form α on a Hilbert space $\mathcal{H}$ a *Fredholm form*, if the corresponding operator $\alpha: H \rightarrow H^* = H$ is Fredholm.

Recall that in the case $M = \mathbb{CP}^1$ the space $H^{1/2}(M, \omega^{1/2})$ is the “best” of Sobolev spaces for $\omega^{1/2}$, since it allows the action of $\mathrm{SL}(2, \mathbb{C})$ (at least “in small”) by bounded operators. Here we see another “nice” property of this particular Sobolev space: there is a canonically defined invertible pairing on this space (invertible in the sense that the corresponding operator is bounded).

Remark 3.5. If we could construct a self-adjoint operator which enjoys the above properties of the operator $\bar{\partial}$, i.e., is $\mathrm{SL}(2, \mathbb{C})$ -invariant and Fredholm, one would be (almost) able to define an invariant Sobolev space structure on $H^{1/2}(M, \omega^{1/2})$. The only thing missing would be positive definiteness.

3.3. The kernel of $\bar{\partial}^{-1}$. Suppose $M = \mathbb{CP}^1$. In this case the operator $\bar{\partial}: \mathcal{D}'(\omega^{1/2}) \rightarrow \mathcal{D}'(\omega^{1/2} \otimes \bar{\omega})$ is invertible, thus the operator $\bar{\partial}^{-1}$ is canonically defined. Consider a kernel $K(x, y)$ of this operator,

$$(\bar{\partial}^{-1}f)(x) = \int K(x, y) f(y).$$

Obviously, $K(x, y)$ is a section of $\omega^{1/2} \boxtimes \omega^{1/2}$.

To calculate $K(x, y)$ consider a δ -section δ_{z_0} of $\omega^{1/2}$ at z_0 (defined up to a scalar multiple). Then $K(\bullet, z_0) = \text{const} \cdot \bar{\partial}^{-1}\delta_{z_0}$ is a holomorphic section of $\omega^{1/2}$ on $\mathbb{CP}^1 \setminus \{z_0\}$. It is easy to see that it has a simple pole at $z = z_0$, thus has no zeros (since $\omega^{1/2} \simeq \mathcal{O}(-1)$). We conclude that $K(x, y)$ has a simple pole at $x = y$. There is only one such a section of $\omega^{1/2} \boxtimes \omega^{1/2}$. Now one may easily recognize the kernel for $\bar{\partial}^{-1}$ in the Cauchy formula

$$\bar{\partial}^{-1}\delta_{z=z_0}dz^{1/2}d\bar{z} = \frac{1}{2\pi i} \frac{1}{z - z_0} dz^{1/2}$$

The right-hand side $f(z, z_0)$ has no pole at ∞ , and is uniquely determined by this condition and the equation

$$\bar{\partial}f = \delta_{z=z_0}dz^{1/2}d\bar{z}$$

in $\mathbb{CP}^1 \setminus \{\infty\}$. We conclude that

$$K(x, y) = \frac{1}{2\pi i} \frac{1}{x - y} dx^{1/2} dy^{1/2}.$$

3.4. Cauchy kernel and L_2 . Consider a smooth embedded curve $\gamma \hookrightarrow \mathbb{CP}^1$ (we do not suppose that γ is compact). Any smooth function f on γ with a compact support may be extended as a generalized δ -function $e(f)$ on $\mathbb{CP}^1$ with a support on γ , same for half-forms on γ . Applying $\bar{\partial}^{-1}$ to the result, we obtain a form on $\mathbb{CP}^1$ which is holomorphic outside of γ . Here we discuss when this mapping $\bar{\partial}^{-1} \circ e$ allows $f \in L_2(\gamma)$ instead of $f \in \mathcal{D}(\gamma)$.

Remark 3.6. In what follows the main example of γ is a disjoint union of infinite number of circles.

Fix a point $z \in \mathbb{CP}^1$ and suppose that some neighborhood of z does not intersect γ . Then for a half-form f

$$\begin{aligned} \left. \frac{\bar{\partial}^{-1} \circ e(f)}{dz^{1/2} d\bar{z}} \right|_{z_0} &\stackrel{\text{def}}{=} \langle \bar{\partial}^{-1} \circ e(f), \delta_{z_0} dz^{1/2} \rangle \\ &= - \langle e(f), \bar{\partial}^{-1} (\delta_{z_0} dz^{1/2}) \rangle \\ &= - \langle e(f), K(\bullet, z_0) dz^{-1/2} \rangle = \int_C f(x) K(x, z_0) dz^{-1/2}. \end{aligned}$$

We conclude that the linear functional of calculating $\bar{\partial}^{-1} \circ e(f)$ at z_0 is given by the kernel $K(\bullet, z_0)|_{\bullet \in C}$. However, since $K(\bullet, z_0)$ is bounded outside of a neighborhood of z_0 , we see that

Proposition 3.7. *The mapping $\bar{\partial}^{-1} \circ e$ extends to a mapping from $L_2(\gamma)$ to the space $\text{Hol}(\mathbb{CP}^1 \setminus \bar{\gamma})$ of forms holomorphic outside of γ iff the length of γ is finite.*

4. TOY THEORY

In this section we study an example which shows the main framework of our approach without any of the complications related to the necessity to consider $H^{1/2}$ -spaces. This is an example related to dealing with *partial period mapping* $\Gamma(M, \omega) \rightarrow \mathbb{C}^g$ of taking periods of global holomorphic forms along A -cycles on the Riemann surface. In fact, we start from studying a related space of global holomorphic functions which are allowed to have constant jumps along A -cycles.

4.1. Toy global space. Consider a complex curve M and the mapping $\partial\bar{\partial}: \Omega^0 \rightarrow \Omega^1$ (let us recall that $\Omega^{\text{top}} = \Omega^1$ in our notations for fractional forms). The integral $-\int_M if \cdot \partial\bar{\partial}\bar{f}$ defines a sesquilinear form on functions with compact support, it is Hermitian, and the only functions which are in the null-space of this form are constants, since the integral is equal to $\int_M i\partial f \cdot \bar{\partial}\bar{f} \geq 0$.

This integral defines a pre-Hilbert structure on global sections of Ω^0 modulo constants, this structure is compatible with the Sobolev H^1 -topology. Moreover, if

$M = \mathbb{CP}^1$, this structure is $\mathrm{PGL}(2, \mathbb{C})$ -invariant, and coincides with the structure of supplementary series of representations of $\mathrm{SL}(2, \mathbb{C})$ (in notations of Section 2.6 it is the case $s = 1$, compare with Remark 2.6). As a result, we obtain a $\mathrm{PGL}(2, \mathbb{C})$ -invariant realization of Hilbert space $H^1(\mathbb{CP}^1)/\mathrm{const}$.

The dual space is the subspace $H_{f=0}^{-1}(\mathbb{CP}^1, \Omega^1)$ of $H^{-1}(\mathbb{CP}^1, \Omega^1)$ consisting of forms with integral 0, thus the latter space also carries an $\mathrm{PGL}(2, \mathbb{C})$ -invariant Hilbert structure. To describe it, note that Remark 2.6 is still applicable, thus in the limit $s \rightarrow 0$ the pre-Hilbert structure of Section 2.6 becomes a form of $\mathrm{rk} = 1$ with $H_{f=0}^{-1}(\mathbb{CP}^1, \Omega^1)$ in the null-space. As a corollary, $\frac{d}{ds}|_{s=0}(\alpha, \beta)_s$ gives a correctly defined positive form on $H_{f=0}^{-1}(\mathbb{CP}^1, \Omega^1)$ which should be dual to the pairing on $H^1(\mathbb{CP}^1)/\mathrm{const}$.

Taking the s -derivative of the kernel K_s from Section 2.6, we get $\log|z_1 - z_2|$ (when restricted on $\mathbb{C} \subset \mathbb{CP}^1$), which is indeed scaling-invariant up to addition of a constant. (Note that the constant is irrelevant, since the forms α and β we are going to pair K with have vanishing integral.) What the above argument shows is that the continuation of this kernel to $\mathbb{CP}^1$ is invariant w.r.t. fraction-linear mappings.

4.2. Almost-perpendicularity. The description of the pairing on $H_{f=0}^{-1}(\mathbb{CP}^1, \Omega^1)$ immediately implies

Proposition 4.1. *Consider two disjoint disks K_1, K_2 on $\mathbb{CP}^1$ of conformal distance l . Let $H_{1,2} = \overset{\circ}{H}_{f=0}^{-1}(K_{1,2}, \Omega^1) \subset H_{f=0}^{-1}(\mathbb{CP}^1, \Omega^1)$. Let P_l be the orthogonal projector from one subspace to another. Then*

$$\|P_l\| \sim e^{-l}.$$

The equivalence means that the quotient of two sides remains bounded and separated from 0 when l varies.

Proof. Since the natural norm on $H_{f=0}^{-1}(\Omega^1)$ is invariant with respect to PGL , the angle between the subspaces $H_{1,2}$ depends on l only. Note that it is sufficient to prove the statement in the case $l > \varepsilon$ for some fixed $\varepsilon > 0$. Now proceed as in the proof of Proposition 5.13. The only change is that the kernel is now $\log|L + z_1 - z_2|$, and on forms with integral 0 it is equivalent to $(\log|L + z_1 - z_2| - \log L - \mathrm{Re} \frac{z_1 - z_2}{L})$, which is L^{-2} times an operator of rank 1, plus much smaller operator. $\square$

Remark 4.2. Note that norm of this projector is much smaller than the norm of the projector from Proposition 5.13 (if l is big enough).

Corollary 4.3. *Consider a family of disjoint disks $\{K_i\}$ in $\mathbb{CP}^1$ with pairwise conformal distances l_{ij} , $i \neq j$. Let $l_{ii} = 0$. If the matrix $(e^{-l_{ij}})$ gives a bounded operator*

$l_2 \rightarrow l_2$, then the natural extension-by-0 mapping

$$\bigoplus_{l_2} \mathring{H}_{f=0}^{-1}(K_i, \Omega^1) \rightarrow H_{f=0}^{-1}(\mathbb{CP}^1, \Omega^1)$$

is a continuous injection, and, dually, the natural restriction mapping

$$H^1(\mathbb{CP}^1) / \text{const} \rightarrow \bigoplus_{l_2} H^1(K_i) / \text{const}$$

is a continuous surjection.

4.3. Topology on the boundary. Consider a disk $K \subset \mathbb{CP}^1$. Given a function f on ∂K , one can consider its decomposition into a sum $f_+ + f_-$ of functions which can be analytically extended into/outside of K . Such a decomposition exists if f is in L_2 , and the summands are uniquely defined up to addition of a constant.

If $f \in H^{1/2}(\partial K)$, then f_+ and f_- are H^1 -functions on K and on $\mathbb{CP}^1 \setminus K$ (compare with Section 2.7), if $f \in H^{1/2}(\partial K) / \text{const}$, then $f_+ \in H^1(K) / \text{const}$, $f_- \in H^1(\mathbb{CP}^1 \setminus K) / \text{const}$, and $f_{\pm}$ are uniquely defined by f . Thus H^1 -norms of $f_{\pm}$ are correctly defined. In the same way as in Section 2.7 one can prove

Lemma 4.4. *The described above extension mapping*

$$H^{1/2}(\partial K) \rightarrow H^1(K) / \text{const} \oplus H^1(\mathbb{CP}^1 \setminus K) / \text{const}$$

is a continuous injection.

Since the Hilbert structure on the right-hand side can be made $\text{PGL}(2, \mathbb{C})$ -invariant, the extension mapping defines the $\text{PGL}(2, \mathbb{C})$ -invariant realization of the Hilbert structure on $H^{1/2}(\partial K)$. The property of invariance can be described in the following way: let φ be a fraction-linear mapping, $\varphi(K) = K'$. Then φ^* defines a unitary operator

$$\varphi^*: H^{1/2}(\partial K') \rightarrow H^{1/2}(\partial K).$$

Remark 4.5. In the same way as we defined a $\text{PGL}(2, \mathbb{C})$ -invariant pairing on sections of Ω^{1-s} on $\mathbb{CP}^1$ (in Section 2.6), one can define a $\text{PGL}(2, \mathbb{R})$ -invariant pairing on section of $\Omega_{\mathbb{RP}^1}^{1-s}$ on $\mathbb{RP}^1 = S^1$. In this way one gets so called supplementary series of representations of $\text{SL}(2, \mathbb{R})$. By expressing the pairing in appropriate coordinate systems on $\mathbb{RP}^1 = S^1$ one can easily see that the described above $\text{SL}(2, \mathbb{R})$ -invariant Hilbert structure coincides with one of these structures.

Definition 4.6. The representation $f = f_+ + f_-$ decomposes $H^{1/2}(\partial K) / \text{const}$ into a direct sum of two subspaces which we denote $H_+^{1/2}(\partial K) / \text{const}$ and $H_-^{1/2}(\partial K) / \text{const}$.

Combining the above description of $H^{1/2}(\partial K)$ with results of the previous section, one obtains

Corollary 4.7. *In the conditions of Corollary 4.3 the natural restriction mapping*

$$H^1(\mathbb{CP}^1) / \text{const} \rightarrow \bigoplus_{l_2} H^{1/2}(K_i) / \text{const}$$

is a continuous surjection.

4.4. Space of holomorphic functions. Consider a family $\{K_i\}$ of disjoint disks in $\mathbb{CP}^1$.

Definition 4.8. We say that a generalized function f on $\mathbb{CP}^1$ is H^1 -holomorphic in $\mathbb{CP}^1 \setminus \bigcup K_i$ if $f \in H^1(\mathbb{CP}^1 \setminus \bigcup K_i)$, and $\bar{\partial}f = 0 \in H^0(\mathbb{CP}^1 \setminus \bigcup K_i, \bar{\omega}) = L_2(\mathbb{CP}^1 \setminus \bigcup K_i, \bar{\omega})$. Denote the the space of H^1 -holomorphic functions in $\mathbb{CP}^1 \setminus \bigcup K_i$ by $\mathcal{H}^{(1)}$.

Note that the Sobolev spaces in this definition are generalized ones. Note also that the Hilbert norm on $L_2(\mathbb{CP}^1 \setminus \bigcup K_i, \bar{\omega})$ is canonically defined by $\|\alpha\|^2 = -i \int \bar{\alpha} \alpha$.

Theorem 4.9. *Suppose that the conformal distance between ∂K_i and ∂K_j is l_{ij} , and the matrix $(e^{-l_{ij}})$ gives a bounded operator $l_2 \rightarrow l_2$. Consider the mapping of taking the boundary value:*

$$b: \mathcal{H}^{(1)} / \text{const} \rightarrow \bigoplus_{l_2} H^{1/2}(\partial K_i) / \text{const}.$$

Let b_- be the component of this mapping going into $\bigoplus_{l_2} H_-^{1/2}(\partial K_i) / \text{const}$. Then b_- is a bounded invertible mapping.

Proof. We already know that b_- is bounded. To show that one can reproduce a function by the component b_- of its restriction to a boundary consider a function $f \in \mathcal{H}^{(1)}$. By definition, it is a restriction of some function $g \in H^1(\mathbb{CP}^1)$, and this function g is defined up to a function with support in $\bigcup K_i$. By the condition of f being holomorphic, $\bar{\partial}g$ is an element of $L_2(\bigcup K_i, \bar{\omega}) = \bigoplus_{l_2} L_2(K_i, \bar{\omega})$, and $\bar{\partial}g$ is defined up to addition of $\bar{\partial} \overset{\circ}{H}^1(\bigcup K_i)$. On the other hand, the topology on $H^1(X)$ (here X is a manifold with coordinates x_k) can be defined by the norm $\|f\|_{L_2}^2 + \sum \|\partial_{x_k} f\|_{L_2}^2$, thus $\overset{\circ}{H}^1(\bigcup K_i) = \bigoplus_{l_2} \overset{\circ}{H}^1(K_i)$.

We conclude that $\bar{\partial}g$ is a canonically defined element of

$$\bigoplus_{l_2} L_2(K_i, \bar{\omega}) / \bar{\partial} \overset{\circ}{H}^1(K_i).$$

Lemma 4.10. *Consider a disk $K \subset \mathbb{CP}^1$. Then there exists a canonical isomorphism*

$$L_2(K, \bar{\omega}) / \bar{\partial} \overset{\circ}{H}^1(K) \simeq H_-^{1/2}(\partial K) / \text{const}.$$

Proof. Let us start with a left-to-right mapping. Let $\alpha \in L_2(K, \bar{\omega}) / \bar{\partial} \mathring{H}^1(K)$. Since $\bar{\partial}$ is an elliptic operator without cokernel and with 1-dimensional null-space, $\bar{\partial}^{-1}\alpha$ is an H^1 -function defined up to addition of a constant and addition of an element of $\mathring{H}^1(K)$. Thus the restriction of f to ∂K is defined up to a constant, so it is an element of $H^{1/2}(\partial K) / \text{const}$. Moreover, it is in $H_-^{1/2}(\partial K) / \text{const}$ since $\bar{\partial}^{-1}\alpha$ is holomorphic outside K .

To get right-to-left mapping note that any element of $H_-^{1/2}(\partial K) / \text{const}$ can be (by definition) continued to a holomorphic outside of K function f , and this function is of class H^1 . Thus to this continuation one can apply the arguments which precede the lemma. $\square$

To finish the proof of the theorem note that the knowledge of $r_-(f)$ allows one to construct $\bar{\partial}f$ (by lemma), so the application of $\bar{\partial}^{-1}$ reconstructs f . $\square$

4.5. Toy Riemann–Roch theorem. Consider a family of disjoint disks K_i , $i \in I$, and the corresponding space of H^1 -holomorphic functions. Suppose that I has an involution $'$ which interchanges two halves of $I = I_+ \amalg I'_+$. Fix automorphisms φ_i of $\mathbb{CP}^1$, $i \in I$, such that $\varphi_{i'} = \varphi_i^{-1}$, and $\varphi_i(\partial K_i)$ is $\partial K_{i'}$ with inverted orientation. Note that if the set I is finite, then after gluing ∂K_i via φ_i one gets a curve of genus $\text{card}(I_+) = \text{card}(I) / 2$.

Associate to a function $f \in \mathcal{H}^{(1)}$ the jump of its boundary value after such a gluing:

$$(4.1) \quad \mathcal{J}: f \mapsto \left(f|_{\partial K_j} - \varphi_j^* \left(f|_{\partial K_{j'}} \right) \right), \quad j \in I_+.$$

Theorem 4.11. *Suppose that the conformal distance between ∂K_i and ∂K_j is l_{ij} , and the matrix $(e^{-l_{ij}} - \delta_{ij})$ gives a compact operator $l_2 \rightarrow l_2$. Then the mapping*

$$\mathcal{J}: \mathcal{H}^{(1)} / \text{const} \rightarrow \bigoplus_{j \in I_+}^{l_2} H^{1/2}(\partial K_j) / \text{const}$$

is a Fredholm operator of index 0.

Proof. Consider the composition $\tilde{\mathcal{J}} = \mathcal{J} \circ b_-^{-1}$. Let us show that $\tilde{\mathcal{J}}$ is Fredholm of index 0, this would immediately imply the statement of the theorem. The mapping φ_j^* interchanges $\pm$ -components of $H^{1/2}(\partial K_j) / \text{const}$ and $H^{1/2}(\partial K_{j'}) / \text{const}$, thus one can identify

$$H^{1/2}(\partial K_j) / \text{const} = H_-^{1/2}(\partial K_j) / \text{const} \oplus H_+^{1/2}(\partial K_j) / \text{const}$$

with $H_-^{1/2}(\partial K_j) / \text{const} \oplus H_-^{1/2}(\partial K_{j'}) / \text{const}$. Denote the composition of $\tilde{\mathcal{J}}$ and this identification by $\mathcal{K}: \bigoplus_{l_2} H_-^{1/2}(\partial K_j) / \text{const} \rightarrow \bigoplus_{l_2} H_-^{1/2}(\partial K_j) / \text{const}$. For $f \in \mathcal{H}^{(1)}$ denote by $f_{j\pm}$ the $\pm$ -components of $f|_{\partial K_j}$. One can easily see that $\mathcal{K} \left((f_{j-})_{j \in I} \right) = (f_{j-} - \varphi_j^*(f_{j'+}))_{j \in I}$, in other words, $\mathcal{K} = \text{id} - \varphi^* \circ b_+ \circ b_-^{-1}$, here $b_+ = b - b_-$. Since

$\varphi^* = \bigoplus_{l_2} \varphi_j^*$ is an isometry, to prove the theorem it is enough to show that $b_+ \circ b_-^{-1}$ is compact. Now we investigate how to reconstruct $H_+^{1/2}$ -components of restriction to a boundary via $H_-^{1/2}$ -components.

Consider the operator with the Cauchy kernel $(y - x)^{-1} dy$, $x, y \in \mathbb{CP}^1$. Restrict this kernel to $\bigcup \partial K_i$. One gets an operator which sends functions on $\bigcup \partial K_i$ to functions on $\bigcup \partial K_i$. Put zeros instead of the diagonal terms (which send functions on ∂K_i to functions on ∂K_i), and call the resulting operator $\mathbf{K}$. Obviously, the functions in the image can be holomorphically extended to K_i . Moreover, if a function was non-zero on ∂K_i only, and could be analytically continued inside K_i , then this function is in the zero-space of $\mathbf{K}$.

Thus the operator $\mathbf{K}$ maps $\bigoplus H_-^{1/2}(\partial K_i) / \text{const}$ to $\bigoplus H_+^{1/2}(\partial K_i) / \text{const}$. Moreover, as the proof of Theorem 4.9 shows, $b_+ \circ b_-^{-1} = \frac{\mathbf{K}}{2\pi i}$, since the pseudodifferential operator $\bar{\partial}^{-1}$ has $\frac{dy}{2\pi i(x-y)}$ as the null-space.

We conclude that the image $V_1 = b(\mathcal{H}^{1/2} / \text{const})$ in $\bigoplus H^{1/2}(\partial K_i) / \text{const}$ is a graph of operator $\frac{\mathbf{K}}{2\pi i}: \bigoplus H_-^{1/2}(\partial K_i) / \text{const} \rightarrow \bigoplus H_+^{1/2}(\partial K_i) / \text{const}$. On the other hand, consider the subspace V_2 of $\bigoplus H^{1/2}(\partial K_i) / \text{const}$ consisting of functions which are in the null-space of $\mathcal{J}$, i.e., the subspace given by the condition

$$f|_{\partial K_j} = \varphi_j^* \left(f|_{\partial K_{j'}} \right), \quad j \in I_+.$$

Since φ_j^* interchanges $H_-^{1/2}(\partial K_j) / \text{const}$ and $H_+^{1/2}(\partial K_{j'}) / \text{const}$ and is unitary, we conclude that the subspace V_2 is a graph of a unitary mapping $\bigoplus \varphi_i^*|_{H_+^{1/2}(\partial K_i)}$.

Thus we are in conditions of the abstract Riemann–Roch theorem (Theorem 7.5), which finishes the proof. $\square$

Remark 4.12. This theorem is an infinite-genus variant of Riemann–Roch theorem for the case of the bundle $\mathcal{O}$ on an algebraic curve M . Indeed, the latter theorem says that the mapping $\bar{\partial}: H^s(M, \mathcal{O}) \rightarrow H^{s-1}(M, \bar{\omega})$ between Sobolev spaces is a Fredholm mapping of index $1 - g(M)$. The theory of elliptic operators says that in a case of smooth compact M the value of s is irrelevant, but in our context we are forced to use the value $s = 1$.

The relation of the mapping $\bar{\partial}$ to the mapping $\mathcal{J}$ is the standard (in mathematical physics) trick of *reduction to boundary*. We do cuts in the curve M to obtain a region $S = \mathbb{CP}^1 \setminus \bigcup K_i$. Instead of $H^s(M, \mathcal{O})$ we consider subspace V of $H^s(S, \mathcal{O})$ which consists of functions which satisfy gluing conditions on the boundary. These gluing conditions are similar to $\mathcal{J}f = 0$, but are applied to functions (instead of functions modulo constants), thus there is an extra condition per cut, total g extra conditions. Let

$$\mathcal{J}_{H^1}: H^1(S, \mathcal{O}) / \text{const} \rightarrow \bigoplus_{l_2} H^{1/2}(\partial K_j) / \text{const}$$

be defined by the formula (4.1). Thus V/const is a subspace of $\text{Ker } \mathcal{J}_{H^1}$ of codimension $g - 1$. Now the translation of the classical Riemann–Roch theorem is that the mapping $\bar{\partial}: \text{Ker } \mathcal{J}_{H^1} \rightarrow H^{s-1}(S, \bar{\omega})$ is of index 0. (Note that the choice $s = 1$ insures that there should be no gluing conditions for elements of $H^{s-1}(M, \bar{\omega})$.)

Now to solve the equation $\bar{\partial}f = h$, $\mathcal{J}_{H^1}f = 0$ we reduce it to a boundary value problem: we use the fact that $\bar{\partial}$ is a surjection on $\mathbb{CP}^1$, thus one can immediately find a function f_0 such that $\bar{\partial}f_0 = h$. Thus $f_1 = f - f_0$ should satisfy $\bar{\partial}f_1 = 0$, $\mathcal{J}_{H^1}f_1 = -\mathcal{J}_{H^1}f_0$. This shows that the index of the mapping $\mathcal{J}$ is 0 in the algebraic case $g < \infty$. We see that Theorem 4.11 is indeed an infinite-dimensional analogue of Riemann–Roch theorem for the bundle $\mathcal{O}$.

Remark 4.13. It is possible to modify Theorem 4.11 to make it applicable to deformations of the bundle $\mathcal{O}$ as well (see Section 6.4 for details). Since to describe Jacobian we need the special case $\mathcal{O}$ only, we do not pursue this venue here.

Remark 4.14. For the particular case of the bundle $\mathcal{O}$ it is possible to prove a much stronger result than Riemann–Roch theorem: that $\mathcal{J}$ is an isomorphism. However, since this result is not true out of context of toy theory, we postpone its proof until Section 9.1, when it is needed for description of Jacobian.

5. $H^{1/2}$ -THEORY

As we have seen in Section 4, Riemann–Roch theorem can be easily proven given appropriate ingredients, such as *almost perpendicularity* of functions with far-separated supports, ability to *restrict* a global section to the boundary, ability to *invert* $\bar{\partial}$ basing on boundary values, and an ability to glue a function provided we know the values inside and outside the given curve with appropriate compatibility conditions along the curve. However, when one tries to apply the same technique to the topology we are most interested in, i.e., $H^{1/2}$ -topology, significant difficulties arise.

5.1. In an ideal world. In what follows we use the (complex) case $n = 1$ of Section 2.5. We have seen that the space $H^{1/2}(M, \omega^{1/2})$ should have a special significance in studying the holomorphic half-forms. Moreover, the results of Section 2.7 suggest that one would be able to describe holomorphic elements of $H^{1/2}(D, \omega^{1/2})$, $D \subset M$, by their restrictions on ∂D . If our world were the ideal world, then the following properties would be satisfied:

1. The action of $\text{SL}(2, \mathbb{C})$ on $H^{1/2}(\mathbb{CP}^1, \omega^{1/2})$ would be an action by uniformly bounded operators;
2. For an embedded curve $\gamma \rightarrow \mathbb{CP}^1$ the restriction mappings $r: H^s(\mathbb{CP}^1, \omega^{1/2}) \rightarrow H^{s-1/2}(\gamma, \Omega^{1/2} \otimes \mu)$ (defined for $s > \frac{1}{2}$) would be defined for $s = \frac{1}{2}$ as well;
3. For the same curve the mapping $e: H^s(\gamma, \Omega^{1/2} \otimes \mu) \rightarrow H^{s-1/2}(\mathbb{CP}^1, \omega^{1/2} \otimes \bar{\omega})$ of continuation by δ -function (defined for $s < 0$) would be defined for $s = 0$ as well.

4. If a curve γ divides a domain S into two parts S_1, S_2 , then the natural restriction mapping $H^{1/2}(S) \rightarrow H^{1/2}(S_1) \oplus H^{1/2}(S_2)$ would be an isomorphism.

To make a long story short, in the ideal world the main technical tools of this paper would behave in a civilized manner, which would simplify the exposition a lot.

Identification of the Hardy space with the subspace of holomorphic functions would be provided by the restriction r . The composition

$$H^{1/2}(\mathbb{CP}^1, \omega^{1/2}) \xrightarrow{r} H^0(\gamma, \Omega^{1/2} \otimes \mu) \xrightarrow{e} H^{-1/2}(\mathbb{CP}^1, \omega^{1/2} \otimes \bar{\omega})$$

would be a “model” of $\bar{\partial}$ -operator¹⁴

$$H^{1/2}(\mathbb{CP}^1, \omega^{1/2}) \xrightarrow{\bar{\partial}} H^{-1/2}(\mathbb{CP}^1, \omega^{1/2} \otimes \bar{\omega}),$$

so that the composition (Cauchy formula)

$$H^{1/2}(\mathbb{CP}^1, \omega^{1/2}) \xrightarrow{r} H^0(\gamma, \Omega^{1/2} \otimes \mu) \xrightarrow{e} H^{-1/2}(\mathbb{CP}^1, \omega^{1/2} \otimes \bar{\omega}) \xrightarrow{\bar{\partial}^{-1}} H^{1/2}(\mathbb{CP}^1, \omega^{1/2})$$

would be continuous. By construction the functions in the image of this operator are holomorphic outside of γ , and the operator r gives an injection of this subspace into $H^0(\gamma, \Omega^{1/2} \otimes \mu)$. Now one may consider this as a Cauchy formula, since the composition is an identity operator on the subspace of holomorphic functions in the domain D bounded by γ .

All these operators would be canonically defined by D , and compatible with projective mappings $\mathbb{CP}^1 \rightarrow \mathbb{CP}^1$, $D \rightarrow D_1$. Thus one would be able to consider the image of r on subspace of holomorphic forms in $H^{1/2}(D, \omega^{1/2})$ as a “model” of this space.

The last property in the list would allow us to glue together the forms which are provided by different means on pieces S_1 and S_2 .

Since we are confined to the current world, the above program will not work, thus we need some workarounds against above three non-facts. We will consider a Riemannian structure on $\mathbb{CP}^1$, and will need to control the size of domains in question, in order for results of Section 2.5 to be applicable. We will also need mollifications of operators r and e , and will need some restrictions on what we can glue together.

5.2. Mollification. Here we introduce a mapping

$$H^0(S^1, \Omega^{1/2} \otimes \mu) \xrightarrow{\tilde{e}} H^{-1/2}(S^1 \times (-\varepsilon, \varepsilon), \omega^{1/2})$$

which a closest existing analogue for the (non-existing) mapping e from the previous section. It will be an invertible mapping onto its image, and it will depend on additional parameters a_n , $n \in \mathbb{Z}$. (Later we will need some particular choice of parameters (a_n) , appropriate for the elliptic operator $\bar{\partial}$ we study.)

¹⁴In the sense of Section 3.1, i.e., it would coincide with $-\bar{\partial}$ on piecewise holomorphic functions.

Fix $\varepsilon > 0$ and a sequence $(a_n)_{n \in \mathbb{Z}}$ such that $0 < a < a_n < A$ for fixed constants a and A . Fix a smooth function $\sigma(y)$, $y \in (-\varepsilon, \varepsilon)$, with a compact support and integral 1. Now map a half-form $e^{2\pi i k x} dx^{1/2}$, $x \in S^1 = \mathbb{R}/\mathbb{Z}$, $k \in \mathbb{Z} + \frac{1}{2}$, into

$$a_k e^{2\pi i k x} \frac{\sigma(|k|y)}{|k|} dz^{1/2}, \quad x \in S^1, y \in (-\varepsilon, \varepsilon), \quad z = x + iy,$$

and continue this mapping linearly to $L_2(S^1, \Omega^{1/2} \otimes \mu) = H^0(S^1, \Omega^{1/2} \otimes \mu)$.

Lemma 5.1. *This mapping is an injection, i.e., an invertible mapping onto its image.*

The dual mapping $H^{1/2}(S^1 \times (-\varepsilon, \varepsilon), \omega^{1/2}) \xrightarrow{\tilde{r}} H^0(S^1, \Omega^{1/2} \otimes \mu)$ given by

$$f(x, y) dz^{1/2} \mapsto g(x) = \sum_k g_k e^{2\pi i k x} dx^{1/2},$$

$$g_k = \int f(x, y) e^{-2\pi i k x} \frac{\sigma(|k|y)}{|k|} dx dy$$

has a similar property: it is a surjection, i.e., an invertible mapping from the quotient by its null-space. It is a close analogue of the (non-existing) mapping r from the previous section.

Suppose $a_k = 1$ for any k . If $\varepsilon \rightarrow 0$, then in weak topology for mappings $\mathcal{D} \rightarrow \mathcal{D}'$ the constructed mappings converge to the mappings of extension as δ -function and restriction, but on Sobolev spaces the norms of these mappings go to ∞ ,

Remark 5.2. In what follows we will use this mapping with following modifications: we suppose that $\text{Supp } \sigma \subset [0, \varepsilon]$, thus we map a half-form on S^1 to a half-form concentrated on a small collar to the *right* of S^1 in $S^1 \times [-\varepsilon, \varepsilon]$. In fact $S^1 \times [-\varepsilon, \varepsilon]$ will be identified with an annulus $\{R e^{-2\pi\varepsilon} < |z - z_0| < R e^{2\pi\varepsilon}\}$ in $\mathbb{C}$, and the mapping would send half-forms on $\{|z| = R\}$ to half-forms concentrated on the *outside* collar.

Similarly, the dual mapping

$$H^{1/2}(R e^{-2\pi\varepsilon} < |z - z_0| < R e^{2\pi\varepsilon}) \xrightarrow{\tilde{r}} H^0(\{|z| = R\})$$

will depend only on value of $f(z)$ on the outside collar.

5.3. Mollification suitable for $\bar{\partial}$. Here we are going to fix the values for the coefficients (a_n) from Section 5.2 which are most suitable for the operator $\bar{\partial}$.

Proposition 5.3. *Fix $\varepsilon > 0$. Let $\gamma = \{z \mid |z| = 1\}$, $\tilde{K} = \{z \mid |z| < e^{2\pi\varepsilon}\}$. For appropriate A and a there exists a sequence (a_n) , $0 < a < a_n < A$, such that the corresponding restriction mapping $\tilde{r}: H^{1/2}(\tilde{K}, \omega^{1/2}) \rightarrow L_2(\gamma, \Omega^{1/2} \otimes \mu)$ coincides with the usual restriction on half-forms which are holomorphic between γ and $\tilde{K}$. The numbers a_n , thus the operator $\tilde{r}$, are uniquely determined.*

Dually,

Proposition 5.4. *Let $U = K \cup (\mathbb{CP}^1 \setminus \tilde{K})$. Consider the Cauchy kernel restricted to $\tilde{K}$. It gives two operators: an operator $\tilde{H}^{-1/2}(\tilde{K}) \xrightarrow{\bar{\partial}^{-1}} H^{1/2}(U)$, and an operator $C^\infty(\gamma) \xrightarrow{\bar{\partial}^{-1}} \mathcal{D}'(U)$. Consider the mollification $\tilde{e}$ of the extension mapping e corresponding to a sequence (a_n) . With appropriate choice of the sequence (a_n) the composition*

$$C^\infty(\gamma) \hookrightarrow L_2(\gamma) \xrightarrow{\tilde{e}} \tilde{H}^{-1/2}(\tilde{K}) \xrightarrow{\bar{\partial}^{-1}} H^{1/2}(U) \hookrightarrow \mathcal{D}'(U)$$

coincides with the mapping $C^\infty(\gamma) \xrightarrow{\bar{\partial}^{-1}} \mathcal{D}'(U)$. The numbers a_n , thus the operator $\tilde{e}$, are uniquely determined by the above condition.

Remark 5.5. Note that the mapping $\tilde{r}$ is a left inverse to the mapping $f \mapsto c_f$ from Section 2.7.

5.4. $\tilde{e} \circ \tilde{r}$ as an approximation to $\bar{\partial}$. Consider a circle γ in $\mathbb{C}$. In what follows we consider particular mollifications of the δ -inclusion $C^\infty(\gamma) \xrightarrow{e} \mathcal{D}'(\mathbb{C})$ and restriction $\mathcal{D}(\mathbb{C}) \xrightarrow{r} C^\infty(S^1)$, which correspond to the only sequences (a_n) which satisfy the conditions of Propositions 5.3, 5.4. The notations $\tilde{e}_\gamma$ and $\tilde{r}_\gamma$ are reserved for these two mappings, we may denote them $\tilde{e}$ and $\tilde{r}$ if the circle to apply them for is clear from context.

A central tool in the following discussion is the mollification of identity (3.1):

Proposition 5.6. *Consider a circle γ or radius ρ which bounds a disk K . Let $\tilde{K}$ be a concentric disk of radius $\rho e^{2\pi\varepsilon}$. Let F be a holomorphic half-form in $\tilde{K} \setminus K$. Let*

$$G = \bar{\partial}^{-1} \circ \tilde{e}_\gamma \circ \tilde{r}_\gamma(F).$$

If F is a holomorphic half-form in $\mathbb{CP}^1 \setminus K$, then G coincides with F outside of $\tilde{K}$, and is 0 in U . If F is a holomorphic half-form in $\tilde{K}$, then G coincides with $-F$ inside of K , and is 0 outside of $\tilde{K}$.

Amplification 5.7. *Fix $\varepsilon, R > 0$. Consider disks $K_\rho = \{|z| < \rho\}$, $K'_\rho = \{|z| < \rho e^{2\pi\varepsilon}\}$, $\tilde{K}_\rho = \{|z| < \rho e^{4\pi\varepsilon}\}$ in $\mathbb{C}$. Let $\gamma = \partial K_\rho$. There exists a mapping*

$$\lambda_\rho: H^{1/2}(\tilde{K}_\rho, \omega^{1/2}) \rightarrow H^{1/2}(\tilde{K}_\rho, \omega^{1/2})$$

such that

1. *For $0 < \rho < R$ the mapping λ_ρ is continuous with the norm uniformly bounded by a constant depending on ε and R only;*
2. *$\lambda_\rho(f)|_{K_\rho} = 0$ for any $f \in H^{1/2}(\tilde{K}_\rho)$;*
3. *$(f - \lambda_\rho(f))|_{\tilde{K}_\rho \setminus K'_\rho} = 0$ for any $f \in H^{1/2}(\tilde{K}_\rho)$;*

4. if f is holomorphic in $\tilde{K}_\rho \setminus K_\rho$, then $\bar{\partial}\lambda_\rho(f) = (\tilde{e}_\gamma \circ \tilde{r}_\gamma)(f)$.

Proof. During the proof we abuse notations and do not mention the bundle $\omega^{1/2}$ in notations for Sobolev spaces.

Note that for a half-form f which is holomorphic outside of K all the statements of the amplification are true if we take $\lambda_\rho(f) = (\bar{\partial}^{-1} \circ \tilde{e} \circ \tilde{r})(f)$. If f is holomorphic inside of $\tilde{K}$, then $\lambda_\rho(f) = f - (\bar{\partial}^{-1} \circ \tilde{e} \circ \tilde{r})(f)$ works. This uniquely defines $\lambda_\rho(f)$ if f is holomorphic inside of $\tilde{K}_\rho \setminus K_\rho$. What we need is to adjust this formula to the case of non-holomorphic half-forms.

Since R is fixed, we can consider the norm from Section 2.5 instead of the equivalent Sobolev $H^{1/2}$ -norm. Since for the former norm the Sobolev spaces $H^{1/2}(\tilde{K}_\rho)$ with different ρ are naturally isomorphic, it is enough to consider $\rho = 1$.

Note that for a holomorphic f the image $\lambda_\rho(f)$ depends on $f|_{\tilde{K}_\rho \setminus K_\rho}$ only. We are going to define λ_ρ in general case such that it satisfies the same condition. Thus λ_ρ is a mapping $H^{1/2}(\tilde{K}_\rho \setminus K_\rho) \rightarrow H^{1/2}(\tilde{K}_\rho)$ such that a half-form in the image is 0 inside K_ρ . We may substitute conformally equivalent domain $S^1 \times (-2\varepsilon, 2\varepsilon)$ instead of $\tilde{K}_\rho \setminus K_\rho$.

Now $H^{1/2}(S^1 \times (-2\varepsilon, 2\varepsilon))$ is an orthogonal sum of subspaces L_n , $n \in \mathbb{Z} + \frac{1}{2}$, spanned by half-forms of the form $\varphi(y) e^{2\pi i n x} dz^{1/2}$, $(x, y) \in S^1 \times (-2\varepsilon, 2\varepsilon)$, $z = x + iy$, thus it is enough to construct uniformly bounded mappings in these subspaces. For a given λ let $\lambda^{(n)}$ be defined as

$$\lambda(\varphi(y) e^{2\pi i n x} dz^{1/2}) = \lambda^{(n)}(\varphi(y)) e^{2\pi i n x} dz^{1/2}.$$

Note that the conditions on $\lambda^{(n)}$ are: $\lambda^{(n)}(\varphi(y))$ vanishes if $y < 0$, $\lambda^{(n)}(\varphi(y)) = \varphi(y)$ if $y > \varepsilon$, and $\lambda^{(n)}(e^{ny})$ has a prescribed value (obtained basing on the last condition of the amplification).

Define $\tilde{\sigma}_n(y)$ via

$$\lambda^{(n)}(e^{ny}) = \tilde{\sigma}_n(y) e^{ny}.$$

Obviously, $\tilde{\sigma}_n(y) = 0$ if $y < 0$, $\tilde{\sigma}_n(y) = 1$ if $y > \varepsilon$. The function $\tilde{\sigma}_n(y)$ is uniquely determined by the last condition of the theorem.

After all these remarks define $\lambda|_{L_n}$ by

$$\varphi(y) e^{2\pi i n x} dz^{1/2} \rightarrow \sigma_n(y) \varphi(y) e^{2\pi i n x} dz^{1/2}.$$

This mapping satisfies all the conditions of the amplification, with a possible exception of uniform boundness of these mappings for different n . To prove the boundness, note that

$$\tilde{\sigma}_n(y) = \tilde{\sigma}_1(ny),$$

and the mappings

$$L_1 \xrightarrow{m_n} L_n: \varphi(y) e^{2\pi i x} \mapsto n^{-1/2} \varphi(ny) e^{2\pi i n x}$$

are uniformly bounded together with their inverse mappings. $\square$

5.5. Mollification for $L_2(\Omega^{1/2} \otimes \mu)$. Consider a family of circles γ_i on $\mathbb{CP}^1$. Fix a projective isomorphism of γ_i with $\{|z| = 1\}$ for every i . This isomorphism identifies an annulus $\{e^{-2\pi\varepsilon} < |z| < e^{2\pi\varepsilon}\}$ with some neighborhood U_i of γ_i .

The $\bar{\partial}$ -adjusted mollification of e gives a mapping

$$\tilde{e}_i: L_2(\gamma_i, \Omega^{1/2} \otimes \mu) \rightarrow H^{-1/2}(\mathbb{CP}^1, \omega^{1/2} \otimes \bar{\omega}).$$

This mapping is a injection of topological vector spaces. In the following sections we discuss under which conditions $\sum_i \tilde{e}_i$ is an injection. Here $\sum_i \tilde{e}_i$ is considered as a mapping from $L_2(\coprod \gamma_i, \Omega^{1/2} \otimes \mu)$ to $H^{-1/2}(\mathbb{CP}^1, \omega^{1/2} \otimes \bar{\omega})$.

This injection is going to be a central tool in Section 6.1, which allows one to identify the space of “admissible” holomorphic half-forms with the space of their boundary values. To provide the criterion for this mapping to be an injection, we need to introduce a notion of “almost perpendicular” subspaces, to study relative position of Sobolev subspaces corresponding to different domains, and to investigate the relative position of domains inside $\mathbb{CP}^1$.

First, we suppose that ε is small enough and the subsets U_i do not intersect. Then it is clear that $\sum_i \tilde{e}_i$ has no null-vectors. Obviously, $\sum_i \tilde{e}_i$ were an injection if the images of $\tilde{e}_i$ were perpendicular for different i . We are going to investigate when these images are *almost perpendicular* for big i . In other words, when the sum of images is direct in the sense of Hilbert topology (i.e., it is an orthogonal sum after an appropriate change of the Hilbert norm to an equivalent one).

5.6. Almost perpendicular subspaces.

Lemma 5.8. *Consider a Hilbert space H and a family of subspaces H_i , $i \in \mathbb{N}$. Denote by a_{ij} the orthogonal projector $H_i \rightarrow H_j$. Let $A = (a_{ij})$ be the matrix of an operator $\mathbf{A}: \bigoplus H_i \rightarrow \prod H_i$, $\prod H_i$ being the space of arbitrary sequences (h_i) , $h_i \in H_i$. Then the natural mapping $\bigoplus H_i \rightarrow H$ extends to a Fredholm mapping*

$$\bigoplus_{l_2} H_i \xrightarrow{i} \text{Im } i \subset H, \quad (h_i) \mapsto \sum h_i,$$

onto its image iff $\mathbf{A}$ induces a Fredholm mapping $\bigoplus_{l_2} H_i \rightarrow \bigoplus_{l_2} H_i$.

Proof. Consider $\mathbf{A}$ as $i^* \circ i$. $\square$

Corollary 5.9. *Suppose that the conditions of the previous lemma the matrix $\alpha = (\|a_{ij}\| - \delta_{ij})$ corresponds to a compact mapping $l_2 \rightarrow l_2$. Then the mapping $\bigoplus_{l_2} H_i \xrightarrow{i}$*

H is a Fredholm operator onto its image. In particular, this is true if

$$\sum_{i \neq j} \|a_{ij}\|^2 < \infty.$$

Moreover, if there is a number C and a family of unit vectors $v_i \in V_i$ such that

$$\|a_{ij}\| < C |(v_i, v_j)|,$$

then the mapping i is continuous iff α provides a bounded mapping $l_2 \rightarrow l_2$.

5.7. Conformal distance. In the applications we consider the curve γ is a union of countably many connected components $\gamma = \coprod \gamma_i$, and all the components γ_i but a finite number are circles. Let $\tilde{e}_i = \tilde{e}_{\gamma_i}$ be the mollified extension mapping, and

$$H_i = \text{Im } \tilde{e}_i.$$

If H_i satisfy the conditions of Lemma 5.8, then the mapping $\sum_i \tilde{e}_i$ from Section 5.5 is a Fredholm mapping onto its image. So the next thing to study is how to calculate the norm of the projection

$$(5.1) \quad \text{Im } \tilde{e}_i \rightarrow \text{Im } \tilde{e}_j.$$

The subspace $\text{Im } \tilde{e}_i$ lies inside a subspace $\mathring{H}(\tilde{K}_i)$, $\tilde{K}_i$ being any disk which contains an appropriate neighborhood of γ_i . Thus to majorate the norm of the projection (5.1) we start with the case of two disjoint disks $\tilde{K}_i, \tilde{K}_j$ on $\mathbb{CP}^1$, and consider the orthogonal projection from $\mathring{H}^{-1/2}(\tilde{K}_i)$ to $\mathring{H}^{-1/2}(\tilde{K}_j)$ inside $H^{-1/2}(\mathbb{CP}^1)$. As we will see, the norm of this projection can be majorated by a number which depends only on some kind of *distance* between $\tilde{K}_i$ and $\tilde{K}_j$.

Two disjoint simple curves γ_1, γ_2 on $\mathbb{CP}^1$ bound a tube U , which is conformally equivalent to exactly one tube $S^1 \times [0, l]$, $l > 0$. Here S^1 is a circle of circumference 2π .

Definition 5.10. We call the number l the *conformal distance* between γ_1 and γ_2 .

Proposition 5.11. Consider three simple curves $\gamma_1, \gamma_2, \gamma_3$ such that γ_2 separates γ_1 and γ_3 . Then the conformal distance $l(\gamma_1, \gamma_3) \geq l(\gamma_1, \gamma_2) + l(\gamma_2, \gamma_3)$.

Proof. Conformal distance between γ and γ' is $\geq l$ iff there exists a function φ defined between γ and γ' such that $\varphi|_\gamma = 0$, $\varphi|_{\gamma'} = l$, and the “energy” $\int \partial\varphi\bar{\partial}\varphi \leq l$. Combining two such functions, one defined between γ_1 and γ_2 , another between γ_2 and γ_3 , we obtain the statement. $\square$

Lemma 5.12. The only $\text{PGL}_2(\mathbb{C})$ -invariant of a couple of disjoint disks on $\mathbb{CP}^1$ is the conformal distance.

5.8. Subspaces of $\Omega^{3/4}$. Consider two disjoint disks K_1, K_2 on $\mathbb{CP}^1$ of conformal distance l . Since the natural norm on $H^{-1/2}(\Omega^{3/4})$ (see Section 2.6) is invariant with respect to PGL , the angle between the subspaces $\mathring{H}^{-1/2}(K_1, \Omega^{3/4})$ and $\mathring{H}^{-1/2}(K_2, \Omega^{3/4})$ depends on l only. Let P_l be the orthogonal projector from one subspace to another.

Proposition 5.13. *Fix $\varepsilon > 0$. If $l \geq \varepsilon$, then*

$$\|P_l\| \sim e^{-l/2}.$$

The equivalence means that the quotient of two sides remains bounded and separated from 0.

Proof. The norm $\|P_l\|$ is a smooth function of l , thus one needs to prove only the asymptotic when $l \rightarrow \infty$. One may represent two disks of conformal distance $\approx l \gg 1$ as $|z| \leq 1$ and $|z - e^{l/2}| \leq 1$. Let $L = e^{l/2}$. Consider the coordinate $z_1 = z$ in the first disk, and $z_2 = z - L$ in the second one. What we need to prove is that the kernel $K_1(z_1, z_2) = \frac{1}{|z_1 - z_2 + L|}$ in $\{(z_1, z_2) \mid |z_1|, |z_2| \leq 1\}$ gives an operator of the norm $\sim 1/L$.

Since the radii are fixed now, we can consider functions instead of $3/4$ -forms. Since the norm of operator with the kernel $K_0 = \frac{1}{L}$ is $O(\frac{1}{L})$, it is enough to prove that the kernel $K_2(z_1, z_2) = \frac{1}{|z_1 - z_2 + L|} - \frac{1}{L}$ corresponds to an operator of norm $o(1/L)$. Let us estimate Hilbert–Schmidt norm of this operator. It is equal to the $H^{1/2} \otimes_{l_2} H^{1/2}$ -norm of K_2 . On the other hand, the last norm is bounded by H^1 -norm, which is obviously $O(L^{-2})$. $\square$

Remark 5.14. Note that if $0 < s < 1$, then a similar statement is true in $H^{1-2s}(\Omega^s)$. We will need only the following statement:

$$\|P_l\| = O(e^{-l/2}) \text{ if } 0 < s < \frac{3}{4},$$

and we will use it in the case $s = \frac{1}{4}$ only.

Corollary 5.15. *Consider a family of disjoint closed disks $K_i \subset \mathbb{CP}^1$, $i \in \mathbb{N}$, with conformal distance between K_i and K_j being l_{ij} . Put $l_{ii} = 0$. Let $\mathcal{L}$ be a linear bundle on $\mathbb{CP}^1$. Suppose that the closure of $\bigcup K_i$ does not coincide with $\mathbb{CP}^1$. Suppose that $\inf_{i \neq j} l_{ij} > 0$, and that the matrix $(e^{-l_{ij}/2})$ corresponds to a bounded operator $l_2 \rightarrow l_2$. Then the inclusions $\mathring{H}^{-1/2}(K_i, \mathcal{L}) \hookrightarrow H^{-1/2}(\mathbb{CP}^1, \mathcal{L})$ extend to a continuous injection*

$$\bigoplus_{l_2} \mathring{H}^{-1/2}(K_i, \mathcal{L}) \hookrightarrow H^{-1/2}(\mathbb{CP}^1, \mathcal{L}).$$

Dually, the restrictions $H^{1/2}(\mathbb{CP}^1, \mathcal{L}) \rightarrow H^{1/2}(K_i, \mathcal{L})$ extend to a continuous surjection

$$H^{1/2}(\mathbb{CP}^1, \mathcal{L}) \rightarrow \bigoplus_{l_2} H^{1/2}(K_i, \mathcal{L}).$$

Proof. If $\mathcal{L} = \Omega^{3/4}$ in the first part of the theorem, or $\mathcal{L} = \Omega^{1/4}$ in the second one, then everything is proved. Otherwise let U be an open subset of $\mathbb{CP}^1 \setminus \bigcup \overline{K_i}$. An isomorphism of $\mathcal{L}$ and $\Omega^{3/4}$ (or $\Omega^{1/4}$) on $\mathbb{CP}^1 \setminus U$ proves the rest. $\square$

Using the results of Remark 5.14 we obtain the following statement:

Proposition 5.16. *In the conditions of Corollary 5.15*

$$\bigoplus_{l_2} \mathring{H}^{1/2}(K_i, \mathcal{L}) \rightarrow H^{1/2}(\mathbb{CP}^1, \mathcal{L}).$$

(components being inclusions) is an isomorphism to its image. The image of this mapping is $\mathring{H}^{1/2}(\bigcup_i K_i, \mathcal{L})$.

6. GENERALIZED HARDY SPACE

6.1. Hilbert space of holomorphic half-forms. In this section we are going to construct a Hilbert space (or at least a space with Hilbert topology) which models global holomorphic half-forms on a curve which is not necessarily compact.

Consider a family of disjoint closed disks $K_i \subset \mathbb{CP}^1$, $i \in I$. Let $\varepsilon > 0$, $\tilde{K}_i$ be the concentric disk to K_i of radius $e^{2\varepsilon} \cdot \text{radius}(K_i)$. Suppose that the disks $\tilde{K}_i$ do not intersect, and denote the conformal distance between $\tilde{K}_i$ and $\tilde{K}_j$ by l_{ij} . Put $l_{ii} = 0$. Suppose that

Condition A . Say that the collection of disks is *well-separated* if

1. $\inf_{i \neq j} l_{ij} > 0$;
2. $\bigcup \tilde{K}_i \neq \mathbb{CP}^1$; and
3. the matrix $(e^{-l_{ij}/2})$ gives a bounded operator $l_2 \rightarrow l_2$;

Note that the interior of the closure $\overline{\mathbb{CP}^1 \setminus \bigcup K_i}$ is non-empty by the above conditions, and

Proposition 6.1. *If disks K_i are well-separated, then $\sum_i \text{radius}(K_i) < \infty$, thus Cauchy kernel defines a mapping from $L_2(\bigcup \partial K_i, \Omega^{1/2} \otimes \mu)$ to the space of analytic functions in $\mathbb{CP}^1 \setminus \bigcup \overline{K_i}$.*

Proof. Indeed, since rows of $(e^{-l_{ij}/2})$ are in l_2 , $\sum_j e^{-l_{ij}} < \infty$. On the other hand, $e^{-l_{ij}}$ is approximately proportional to radius of K_j when i is fixed. $\square$

Moreover, if we slightly decrease the value of ε , then the first two conditions on the family $\{K_i\}$ will be automatically satisfied, so only the last condition is the new one.

Definition 6.2. The *generalized Hardy space* $\mathcal{H}(\mathbb{CP}^1, \{K_i\})$ is the space

$$\left\{ f(z) \in H^{1/2} \left(\mathbb{CP}^1 \setminus \bigcup_i K_i, \omega^{1/2} \right) \mid \bar{\partial} f = 0 \in H^{-1/2} \left(\mathbb{CP}^1 \setminus \bigcup_i K_i, \omega^{1/2} \otimes \bar{\omega} \right) \right\}.$$

Note that elements of $\mathcal{H}(\mathbb{CP}^1, \{K_i\})$ are holomorphic half-forms in $\mathbb{CP}^1 \setminus \bigcup_i \widetilde{K}_i$. Note also that the Sobolev space in the definition is a generalized Sobolev space, since $\mathbb{CP}^1 \setminus \bigcup_i K_i$ is not open if the set I is infinite.

Remark 6.3. Note that if I consists of one element, then $\mathcal{H}$ is the usual Hardy space from Section 2.7 with modifications outlined in Remark 2.10.

In Section 6.7 we introduce a slightly weaker condition on domains K_i , and will use it instead of Condition A. Note that we are going to mention the space $\mathcal{H}$ only in cases when the Condition A is satisfied.

6.2. Hilbert operator. A section of $\Omega^{1/2} \otimes \mu$ on a circle can be decomposed into the Fourier series

$$f(t) dt^{1/2} = \sum_{n \in \mathbb{Z} + 1/2} a_n e^{2\pi i n t} dt^{1/2}$$

(half-integers appear because of the factor μ), hence it can be written as a sum of a component f_+ which can be holomorphically extended to a section of $\omega^{1/2}$ inside the circle, and a component f_- which may be extended outside of the circle. Moreover, norms of $f_{\pm}$ are bounded by the norm of f , and the decomposition is unique.

We denote this decomposition $L_2^+(\gamma, \Omega^{1/2} \otimes \mu) \oplus L_2^-(\gamma, \Omega^{1/2} \otimes \mu)$.

Proposition 6.4. Consider the operator $\mathbf{K}_+$ with Cauchy kernel acting in the space $\bigoplus_{l_2} L_2(\partial K_i, \Omega^{1/2} \otimes \mu)$. Let $\mathbf{K}$ be the operator $\widetilde{\mathbf{K}}$ with diagonal blocks $L_2(\partial K_i, \Omega^{1/2} \otimes \mu) \rightarrow L_2(\partial K_i, \Omega^{1/2} \otimes \mu)$ removed. If the disks K_i are well-separated, then $\mathbf{K}$ is bounded. If we decompose

$$L_2(\partial K_i, \Omega^{1/2} \otimes \mu) = L_2^+(\partial K_i, \Omega^{1/2} \otimes \mu) \oplus L_2^-(\partial K_i, \Omega^{1/2} \otimes \mu),$$

then the only nonzero blocks of $\mathbf{K}$ act from $L_2^-(\partial K_i, \Omega^{1/2} \otimes \mu)$ to $L_2^+(\partial K_j, \Omega^{1/2} \otimes \mu)$, $i \neq j$.

This statement is an immediate corollary of the first part of

Lemma 6.5. Consider a Hilbert space $H = \bigoplus_{l_2} H_i$ and a linear operator $A: H \rightarrow H$ with bounded blocks $A_{ij}: H_j \rightarrow H_i$. Let $a_{ij} = \|A_{ij}\|$. Suppose that the matrix (a_{ij}) gives a bounded operator $\alpha: l_2 \rightarrow l_2$. Then

1. A is bounded, $\|A\| \leq \|\alpha\|$;

2. A is compact if α is compact and each of operators A_{ij} is compact.

Proof. To bound the operator A it is enough to bound $|(x, Ay)|$ for $|x| = |y| = 1$, $x, y \in H$. On the other hand, decomposition of H gives $x = (x_i)$, $y = (y_i)$, and $|x|^2 = \sum |x_i|^2$, $|y|^2 = \sum |y_i|^2$. This gives

$$|(x, Ay)| = \left| \sum_{ij} (x_i, A_{ij}y_j) \right| \leq \sum_{ij} |(x_i, A_{ij}y_j)| \leq \sum_{ij} \|A_{ij}\| |x_i| |y_j|$$

and the latter sum is just $(\xi, \alpha\eta)$ with $\xi = (|x_i|) \in l_2$, similarly for η .

To prove the second part it is enough to show that we can approximate A (in norm) by compact operators. Since α is compact, we can approximate it (in norm) by a matrix with finite number of non-zero elements. The relation of norm of A and norm of α shows that A can be approximated by an operator A' which has the same blocks as A in finite number of places, all the rest is 0.

Since blocks of A are compact, A' is compact as well. We conclude that A can be approximated with arbitrary precision by a compact operator, thus A is compact. $\square$

Corollary 6.6. *Any block-row and block-column of $\mathbf{K}$ gives a compact operator.*

6.3. Boundary map. Here we consider which of the facts from Section 2.7 have sense for the generalized Hardy space as well.

Fix $i \in I$, let $S_i = \partial K_i$. Consider the mollified restriction mapping

$$H^{1/2}(\tilde{K}_i \setminus K_i, \omega^{1/2}) \xrightarrow{\tilde{r}_i} L_2(S_i, \Omega^{1/2} \otimes \mu).$$

Since any element f of the generalized Hardy space is $H^{1/2}$ in $\tilde{K}_i \setminus K_i$ (and holomorphic inside this annulus), the restriction $\tilde{r}_i(f) = f|_{S_i}$ of this half-form on S_i is an L_2 -section of $\Omega^{1/2}(S_i) \otimes \mu$. In the rest of this section we are going to abuse notations and denote $\tilde{r}_i(f)$ as $f|_{S_i}$.

Theorem 6.7. *Suppose that the disks K_i are well-separated. Then the mappings $\tilde{r}_i$ taken together provide a mapping*

$$\mathcal{H}(\mathbb{CP}^1, \{K_i\}) \xrightarrow{\tilde{r}} \coprod L_2(S_i, \Omega^{1/2} \otimes \mu),$$

which is fact is a continuous mapping

$$\mathcal{H}(\mathbb{CP}^1, \{K_i\}) \xrightarrow{\tilde{r}} \bigoplus_{l_2} L_2(S_i, \Omega^{1/2} \otimes \mu).$$

The image of $\tilde{r}$ is closed. Moreover, the mapping $\tilde{r}$ is invertible onto its image.

Proof. In fact, all the main ingredients for the proof of this theorem are already here. An element $f \in \mathcal{H}(\mathbb{CP}^1, \{K_i\})$ is by definition a restriction of some $H^{1/2}$ -section g of $\omega^{1/2}$ on the whole sphere $\mathbb{CP}^1$. The restrictions of g to $\tilde{K}_i$ give an element of $\bigoplus_{l_2} H^{1/2}(\tilde{K}_i, \omega^{1/2})$ by Corollary 5.15. On the other hand, given f , the restriction of

g on $\tilde{K}_i$ is defined up to a section with support on K_i . Hence the mollified restriction on S_i is correctly defined, and it has a norm majorated by some multiple of the norm of the restriction on $\tilde{K}_i$. Hence the restriction mapping $\tilde{r}$ is bounded indeed.

To show that $\text{Im } \tilde{r}$ is closed, let us construct a left inverse l this operator. Then $\tilde{r}l$ is going to be a projection on $\text{Im } \tilde{r}$, which will prove the closeness.

Consider the mapping λ_i from Section 5.4, associated to the disks $K_i, \tilde{K}_i$. Then $g - \lambda_i(g)$

1. vanishes outside of $\tilde{K}_i$;
2. coincides with g inside of K_i ;
3. has a norm bounded by $C \cdot \|g|_{\tilde{K}_i}\|$;
4. depends on values of g outside of K only.

Combining all this together, we get

$$F = g + \sum_i (\lambda_i(g) - g)$$

which is a half-form of the norm bounded by

$$\|g\| + C' \cdot \left(\sum \|g|_{\tilde{K}_i}\|^2 \right)^{1/2} = O(\|g\|).$$

The half-form F is equal to g (thus to f) outside of $\bigcup \tilde{K}_i$, and is equal to 0 inside all K_i . Moreover, since f is holomorphic, one can calculate $\bar{\partial}F|_{\tilde{K}_i}$ as $\tilde{e}_i(f|_{\partial K_i})$.

Proposition 6.8. *Let f be a half-form from the generalized Hardy space. There exists a half-form $F \in H^{1/2}(\mathbb{CP}^1)$ such that:*

1. $\|F\| < C \cdot \|f\|$;
2. $f = F$ outside of $\bigcup \tilde{K}_i$;
3. $F = 0$ inside of K_i for any i ;
4. $\bar{\partial}F = \sum \tilde{e}_i(f|_{\partial K_i})$.

Proof. The only part which needs proof is the last one. First of all, the sum in the right-hand side obviously converges in $H^{-1/2}$, since the disks $\tilde{K}_i$ are separated far enough, and norms of the restrictions of f onto ∂K_i form a sequence in l_2 . The same arguments show that the right-hand side is contained in the generalized Sobolev subspace $\mathring{H}^{-1/2}(\bigcup \tilde{K}_i)$. Moreover, $\bar{\partial}(F - g)$ is contained in the same space, and $\bar{\partial}f$ is there by the definition of the generalized Hardy space. Since g coincides with f outside of $\bigcup K_i$, $\bar{\partial}F \in \mathring{H}^{-1/2}(\bigcup \tilde{K}_i)$.

Thus we know that the difference of the right-hand side and left-hand side is contained in $\mathring{H}^{-1/2}(\bigcup \tilde{K}_i)$ and is 0 inside any $\tilde{K}_i$. On the other hand, since $\tilde{K}_i$ are well-separated, $\mathring{H}^{-1/2}(\bigcup \tilde{K}_i) = \bigoplus_{l_2} \mathring{H}^{-1/2}(\tilde{K}_i)$, what finishes the proof. $\square$

Remark 6.9. It is obvious that $f \mapsto F$ is a continuous linear mapping $\mathcal{H}(\mathbb{CP}^1, \{K_i\}) \rightarrow H^{1/2}(\mathbb{CP}^1, \omega^{1/2})$. Moreover, it is an injection. To show this, one needs only to prove that $f|_{\bigcup \tilde{K}_i}$ is determined by F (and bounded by the norm of F). Since the disks $\tilde{K}_i$ are well-separated, it is enough to show this for one particular disk $\tilde{K}_i$.

Since $\tilde{e}_i$ is an injection, $\bar{\partial}F$ determines $\tilde{r}_i(f)$, thus $\lambda_i(g)$ (by construction of λ), thus $(F - f)|_{\tilde{K}_i}$. We obtained

Corollary 6.10. *The relation $f \mapsto F$ is an injection $\mathcal{H}(\mathbb{CP}^1, \{K_i\}) \rightarrow H^{1/2}(\mathbb{CP}^1, \omega^{1/2})$. Since $\bar{\partial}: H^{1/2}(\mathbb{CP}^1, \omega^{1/2}) \rightarrow H^{-1/2}(\mathbb{CP}^1, \omega^{1/2} \otimes \bar{\omega})$ is an isomorphism, the relation $f \mapsto \bar{\partial}F$ gives an injection $\mathcal{H}(\mathbb{CP}^1, \{K_i\}) \rightarrow H^{-1/2}(\mathbb{CP}^1, \omega^{1/2} \otimes \bar{\omega})$. Moreover, the last mapping may be pushed through $\mathcal{H}(\mathbb{CP}^1, \{K_i\}) \rightarrow \bigoplus_{l_2} H^{-1/2}(\tilde{K}_i, \omega^{1/2} \otimes \bar{\omega})$.*

Since the operator $\bar{\partial}$ has no null-space on $H^{1/2}(\mathbb{CP}^1, \omega^{1/2})$, we obtain

Corollary 6.11. *$\bar{\partial}^{-1}(\sum \tilde{e}_i(f|_{\partial K_i}))$ equals f outside of $\bigcup \tilde{K}_i$, i.e., modulo $\mathring{H}^{1/2}(\bigcup \tilde{K}_i)$.*

Since $\bar{\partial}F$ is determined by $\tilde{r}(f)$, we found a left inverse to the mapping $\tilde{r}$, thus it is an injection and the image is closed. This finishes the proof of Theorem 6.7. $\square$

The next step is to describe the image of the operator in question. Let $\gamma_i = \partial K_i$, $\gamma = \bigcup \gamma_i$. We claim that an element of the image of the mapping $\tilde{r}$ is uniquely determined by the minus-components, and any collection of minus-components of bounded norm is possible.

Proposition 6.12. *Consider a decomposition of the space $L_2(\gamma_i, \Omega^{1/2}) = L_2^+(\gamma_i) \oplus L_2^-(\gamma_i)$ into subspaces of half-forms which may be holomorphically extended inside the circle and outside the circle. The image of $\tilde{r}$ consists of sequences $(f_i^\pm)$ such that*

$$f_i^+ = \sum_{j \neq i} \mathbf{K} f_j^-.$$

Here $\mathbf{K}$ is the Hilbert operator, i.e., the operator with Cauchy kernel.

Proof. First of all, note that the proof of Theorem 6.7 together with Propositions 3.7 and 6.1 shows that knowing $\tilde{r}(f)$ one can write f by an explicit formula outside of $\bigcup \tilde{K}_i$. On the other hand, it is easy to see that if we throw away the restriction that we want our operators to be continuous in H^s -topology, one can reconstruct f outside of $\bigcup \tilde{K}_i$.

Indeed, since the curve γ has a finite length, restriction of any smooth section α of $\omega^{1/2}$ to γ is in $L_2(\gamma)$, thus $\int \alpha|_S \tilde{r}(f)$ is correctly defined. By duality, this means that extension $e(\tilde{r}(f))$ of $\tilde{r}(f)$ to $\mathbb{CP}^1$ by δ -function is a correctly defined generalized section of $\omega^{1/2} \otimes \bar{\omega}$. (Note that we consider e , not $\tilde{e}$!)

Lemma 6.13. *Half-form f coincides with $\bar{\partial}^{-1}e(\tilde{r}(f))$ outside of $\bigcup \tilde{K}_i$.*

Proof. Indeed, by the construction of $\tilde{e}_i$, $\bar{\partial}^{-1}\tilde{e}_i(\tilde{r}_i(f))$ coincides with $\bar{\partial}^{-1}e(\tilde{r}_i(f))$ outside of $\tilde{K}_i \setminus K_i$, thus $\bar{\partial}^{-1}e(\tilde{r}(f))$ coincides with f outside of $\bigcup \tilde{K}_i$. On the other hand, $\bar{\partial}^{-1}e(\tilde{r}(f))$ is holomorphic outside of $\bar{\gamma}$, thus this equality can be extended up to γ . $\square$

Consider now $f = \bar{\partial}^{-1}e(\tilde{r}(f))$ near γ_i . Breaking $\tilde{r}(f)$ into two components, $(\tilde{r}(f) - \tilde{r}_i(f))$ and $\tilde{r}_i(f)$, we conclude that

$$f = \bar{\partial}^{-1}e(\tilde{r}(f) - \tilde{r}_i(f)) + \bar{\partial}^{-1}e(\tilde{r}_i(f)).$$

The first summand is holomorphic inside $\tilde{K}_i$ and coincides with $\sum_{j \neq i} \mathbf{K} f_j^-$, since $\mathbf{K}$ kills f_j^+ . The second summand is holomorphic outside of ∂K_i , so its $+$ -part vanishes on ∂K_i , which finishes the proof of Proposition 6.12. $\square$

6.4. Gluing conditions. We continue using notations of Section 6.1. Suppose that the set I has an involution $': I \rightarrow I$ which interchanges two subsets I_+ and I'_+ , $I = I_+ \amalg I'_+$. Thus all the disks K_i are divided into pairs $(K_i, K_{i'})$, $i \in I_+$. Fix a fraction-linear identification φ_i of $\partial K_{i'}$ and ∂K_i which reverses the orientation of the circles, $\varphi_{i'} = \varphi_i^{-1}$. Let φ_i identifies the boundary of the disk $\tilde{K}_{i'}$ with the boundary of $\mathring{K}_i$, and $\partial \tilde{K}_i$ with $\partial \mathring{K}_{i'}$, thus $\mathring{K}_i \subset K_i \subset \tilde{K}_i$.

Let R_i be the annulus between $\tilde{K}_i$ and $\mathring{K}_i$. The mapping φ_i identifies R_i with $R_{i'}$. Let S be the part of $\mathbb{CP}^1$ which lies outside of all the disks $\mathring{K}_i$. Glue the annuli $R_i \subset S$ with $R_{i'} \subset S$ using φ_i .

Definition 6.14. A *model space* $\bar{M}$ is the set obtained from S by identifying the annuli R_i and $R_{i'}$ using φ_i .

Note that $\bar{M}$ consists of two parts: a smooth manifold M which is the image of $S \setminus \bigcup \mathring{K}_\bullet$, and the rest, which one should consider as “infinity” M_∞ of the manifold M (compare with Section 0.7). Unfortunately, the topology on $\bar{M}$ in neighborhood of infinity is not suitable for studying the Riemann–Roch theorem, so we will not consider it in this paper.

Note that Section 6.8 suggests a different topology on $\bar{M}$. We will use some features of this topology when we discuss a mapping into universal Grassmannian.

6.5. Strong sections and duality. In the notations of the previous section consider now holomorphic functions ψ_i defined in R_i . Suppose that ψ_i are nowhere 0, and $\psi_{i'} \circ \varphi_i = \psi_i^{-1}$.

Define $\mathcal{L}$ to be a sheaf on $\bar{M}$ associated with gluing conditions ψ_i , i.e., for $U \subset \bar{M}$ the section of $\mathcal{L}$ on U is a function f on $\tilde{U} \subset \mathbb{CP}^1$ such that $f(\varphi_i(x)) = \psi_i(x) f(x)$ whenever both sides have sense (here $\tilde{U}$ is an appropriate covering subset of $\mathbb{CP}^1$).

Let $\mathcal{L}^{-1}$ be the sheaf associated with gluing conditions ψ_i^{-1} . Similarly define the tensor product of two sheaves defined via gluing conditions.

The identifications φ_i fixes an identification φ_i^* of half-forms on R_i and on $R_{i'}$ up to a sign. Choose this sign for all $i \in I_+$.¹⁵ Let $\omega^{1/2} \otimes \mathcal{L}$ be the sheaf on $\bar{M}$ consisting of half-forms on S such that

$$(6.1) \quad \varphi_i^* (\alpha|_{R_{i'}}) = \psi_i \cdot \alpha|_{R_i}, \quad i \in I.$$

Similarly define $\omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega}$.

To write an analogue of (6.1) for sections in Sobolev spaces, consider a subannulus $\mathring{R}_i \subset R_i$ such that $\mathring{R}_{i'} = \varphi_i \left(\mathring{R}_i \right)$. We suppose that one can find numbers $C, D > 0$ and a sequence (s_i) , $i \in I$, $|s_i| + C + D < 1$, such that the annulus $\mathring{R}_i$ is described as $((s_i - C)\varepsilon, (s_i + C)\varepsilon) \times S^1$ in the conformal coordinate system such that R_i is $(-\varepsilon, \varepsilon) \times S^1$. Let K'_i be the disk bounded by the inner boundary of $\mathring{R}_i$.

Consider a subset $\mathring{S} \subset S$, $\mathring{S} = \mathbb{CP}^1 \setminus \bigcup_i K'_i$. One obtains the same manifold $\bar{M}$ by gluing $\mathring{R}_i \subset \mathring{S}$ as by gluing $R \subset S$, but in what follows it will be more convenient to have a choice of annuli $\mathring{R}_i$, and have them separated from boundary of R_i .

Definition 6.15. Let $H^s(\bar{M}, \omega^{1/2} \otimes \mathcal{L})$ be the subspace of $H^s(\mathbb{CP}^1 \setminus \bigcup_i K'_i, \omega^{1/2})$ consisting of sections which satisfy the gluing conditions (6.1), similarly define $H^s(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega})$.

Dually,

Definition 6.16. Let $\mathring{H}^s(\bar{M}, \omega^{1/2} \otimes \mathcal{L})$ be the quotient of $\mathring{H}^s(\mathbb{CP}^1 \setminus \bigcup_i K'_i, \omega^{1/2})$ by the subspace of $\mathring{H}^s(\bigcup_i \mathring{R}_i, \omega^{1/2})$ consisting of sections which satisfy the gluing conditions

$$(6.2) \quad \varphi_i^* (\alpha|_{\mathring{R}_{i'}}) = -\psi_i \cdot \alpha|_{\mathring{R}_i}, \quad i \in I,$$

similarly define $\mathring{H}^s(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega})$.

Indeed, these definitions are dual due to

Lemma 6.17. *If the disks K_i are well-separated, the spaces $H^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L})$ and $\mathring{H}^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}^{-1} \otimes \bar{\omega})$ are mutually dual w.r.t. the pairing $\int \alpha \beta$.*

Proof. The spaces $H^s(\mathbb{CP}^1 \setminus \bigcup_i K'_i, \omega^{1/2} \otimes \mathcal{L})$ and $\mathring{H}^{-s}(\mathbb{CP}^1 \setminus \bigcup_i K'_i, \omega^{1/2} \otimes \mathcal{L}^{-1} \otimes \bar{\omega})$ are mutually dual w.r.t. this pairing by definition. What remains to prove is that the orthogonal complement to the subspace of $H^s(\mathbb{CP}^1 \setminus \bigcup_i K'_i, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega})$ given by (6.1) is given by

$$(6.3) \quad \varphi_i^* (\alpha|_{R_{i'}}) = -\psi_i^{-1} \cdot \alpha|_{R_i}, \quad i \in I,$$

in $\mathring{H}^{-s}(\mathbb{CP}^1 \setminus \bigcup_i K'_i, \omega^{1/2} \otimes \mathcal{L}^{-1} \otimes \bar{\omega})$, provided $s = 1/2$. This statement is obvious for any fixed $i \in I$.

¹⁵This corresponds to picking a representative of φ_i in 2-sheet cover $\mathrm{SL}(2, \mathbb{C}) \rightarrow \mathrm{PGL}(2, \mathbb{C})$.

On the other hand, solutions to (6.3) form a direct sum over i , and by the second part of Corollary 5.15, solutions to (6.1) form a “direct” intersection (i.e., an intersection of subspaces with almost orthogonal complements). $\square$

Lemma 6.18. *Suppose that the disks K_i are well-separated. The natural mapping*

$$(6.4) \quad \mathring{H}^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega}) \rightarrow H^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega})$$

is an isomorphism. Dually,

$$\mathring{H}^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) \rightarrow H^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L})$$

is an isomorphism.

Proof. We may suppose that all the rings $\mathring{R}_i$ have conformal distance between boundaries greater than $2C\varepsilon$, $C > 0$. Consider a function $\sigma(x)$, $x \in (-C\varepsilon, C\varepsilon)$, such that $\sigma(x) = 0$ near the left end, $\sigma(x) + \sigma(1-x) = 1$. This gives a cut-off function in all the rings $\mathring{R}_i$, and we may extend it by 0 into all K'_i , and by 1 into S . We obtain a function on $\mathbb{CP}^1$.

By the results of Section 5.8 and Amplification 2.7 the multiplication by this function is a bounded operator in $H^{1/2}$. Clearly, this operator provides an inverse to (6.4). $\square$

The operator $\bar{\partial}$ gives mappings

$$\begin{aligned} \mathring{H}^s(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) &\rightarrow \mathring{H}^{s-1}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega}), \\ H^s(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) &\rightarrow H^{s-1}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega}), \end{aligned}$$

which by the previous lemma induce mappings

$$\begin{aligned} \mathring{H}^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) &\xrightarrow{\bar{\partial}} H^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega}), \\ H^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) &\xrightarrow{\bar{\partial}} \mathring{H}^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega}) \end{aligned}$$

(obviously, it does not matter which of isomorphisms of Lemma 6.18 we use to obtain these mappings).

Definition 6.19. Define the space $\Gamma_{\text{strong}}(\bar{M}, \omega^{1/2} \otimes \mathcal{L})$ of *strong global holomorphic sections* of $\omega^{1/2} \otimes \mathcal{L}$ as

$$\Gamma_{\text{strong}}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) = \text{Ker} \left(H^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) \xrightarrow{\bar{\partial}} H^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega}) \right).$$

Note that the conditions on strong global sections α are: they are holomorphic sections of $\omega^{1/2} \otimes \mathcal{L}|_M$ (since $\bar{\partial}\alpha$ vanishes on M), they do not grow very quick near M_∞ (since they belong to $H^{1/2}$), and they have no residue on M_∞ (since $\bar{\partial}\alpha$ vanishes on M_∞).

6.6. Weak sections. The space Γ_{strong} from the last section is good for studying the duality conditions, but it is not suitable for description of global sections via boundary conditions.

Definition 6.20. Define the space $\Gamma_{\text{weak}}(\bar{M}, \omega^{1/2} \otimes \mathcal{L})$ of *weak global holomorphic sections* of $\omega^{1/2} \otimes \mathcal{L}$ as forms from the generalized Hardy space $\alpha \in \mathcal{H}(\mathbb{CP}^1, \{K_i\})$ such that the (mollified) restrictions on the circles ∂K_i satisfy the gluing conditions. In other words,

$$\Gamma_{\text{weak}}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) = \{\alpha \in \mathcal{H}(\mathbb{CP}^1, \{K_i\}) \mid \psi_i \tilde{r}_i(\alpha) = \varphi_i^*(\tilde{r}_{i'}(\alpha))\}.$$

There is a natural mapping

$$\Gamma_{\text{strong}} \rightarrow \Gamma_{\text{weak}}.$$

Theorem 6.21. *Suppose that the disks K_i are well-separated, and for some $A > 1$ either $|\psi_i(z)| < A$, or $|\psi_i(z)| > 1/A$ for any i and $z \in R_i$. Then the above mapping is an isomorphism for an appropriate choice of annuli $\mathring{R}_i$.*

Proof. Indeed, any weak section α is a holomorphic form inside $\mathbb{CP}^1 \setminus \overline{\bigcup_i K_i}$. Consider $\psi_i^{-1} \cdot \varphi_i^*(\alpha)$, it is a holomorphic form inside $K_i \setminus \mathring{K}_i$. Since α satisfies gluing conditions, $\alpha|_{\tilde{K}_i \setminus K_i}$ has the same Laurent coefficients as $\psi_i^{-1} \cdot \varphi_i^*(\alpha)$ (we used compatibility of $\tilde{r}$ with r on holomorphic forms), thus these two forms are restrictions of the same holomorphic form defined inside $\tilde{K}_i \setminus \mathring{K}_i$. We see that α can be extended (as a holomorphic form) into $\mathbb{CP}^1 \setminus \bigcup_i \mathring{K}_i$, and this holomorphic form satisfies the gluing conditions (6.1).

What remains to prove is that we can bound the norm of this extension. On the other hand, this is a local statement, since one can represent $\alpha = \alpha_1 + \alpha_2$, and $\alpha = 0$ inside an appropriate circle concentric with $\tilde{K}_i$, $\alpha_2 = 0$ outside of $\bigcup \tilde{K}_i$. Thus the only thing we need to prove is that inside R_i the form α can be extended across ∂K_i without increasing its norm too much.

Take $C = \frac{1}{4}$, let $\varepsilon_i = C$ if $|\psi_i(z)| < A$, $\varepsilon_i = -C$ otherwise. With this choice of $\mathring{R}_i$ we know that the $H^{1/2}$ -norms of $\alpha|_{\tilde{K}_i \setminus K_i}$ and $\alpha|_{K_i \setminus K'_i}$ are bounded by the norms of $\alpha|_{\tilde{K}_i \setminus K_i}$ and $\alpha|_{\tilde{K}_{i'} \setminus K_{i'}}$. Now the theorem becomes a corollary of the following

Lemma 6.22. *Fix two numbers $A > a > 0$. Consider an annulus R with concentric boundaries and conformal distance between boundaries between A and a . Let V_R be the space of holomorphic half-forms in R which belong to $H^{1/2}(R)$. Then the mapping of taking boundary value*

$$b: H^{1/2}(R, \omega^{1/2}) \rightarrow L_2(\partial R, \Omega^{1/2}) : \alpha \mapsto \alpha|_{\partial R}$$

is an invertible mapping to its (closed) image, and the norms of this mapping and its inverse are bounded by numbers depending on A and a only.

This lemma is a variation of what we did in Section 2.7, with a disk substituted with an annulus. It can be proven in the same way as the case of a disk.

This finishes the proof of the theorem. $\square$

Remark 6.23. To complement the notion of weak holomorphic sections, let us define the spaces of “weak” $H^{\pm 1/2}$ -section in such a way that

$$\Gamma_{\text{weak}}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) = \text{Ker} \left(H_{\text{weak}}^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) \xrightarrow{\bar{\partial}} H_{\text{weak}}^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega}) \right) :$$

Definition 6.24. Let $H_{\text{weak}}^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L})$ be the subspace of $H^{1/2}(\mathbb{C}P^1 \setminus \bigcup_i K_i, \omega^{1/2})$ consisting of sections α which satisfy the gluing conditions:

$$\psi_i \tilde{r}_i(\alpha) = \varphi_i^*(\tilde{r}_{i'}(\alpha)), \quad i \in I.$$

Let $H_{\text{weak}}^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega}) = H^{-1/2}(\mathbb{C}P^1 \setminus \bigcup_i K_i, \omega^{1/2} \otimes \bar{\omega})$.

6.7. Finite-degree bundles. Let d_i be the *index* of ψ_i , i.e., the degree of the mapping $\arg \psi_i: K_i \rightarrow S^1$. We say that the collection of gluing data $\{\psi_i\}$ is of *finite degree* if $d_i = 0$ for all but the finite number of indices $i \in I$. The collection $\{\psi_i\}$ is *semibounded* if for some fixed number C and any i either $|\psi_i(z)|$ or $|\psi(z)|_i^{-1}$ is bounded by C if $z \in R_i$. The *degree* of the finite-degree collection $\{\psi_i\}$ is the sum

$$\sum_{i \in I_+} d_i = \frac{1}{2} \sum_{i \in I} d_i.$$

Definition 6.25. The *degree* of a bundle $\mathcal{L}$ defined by gluing conditions ψ_i is the degree of the collection $\{\psi_i\}$.

Amplification 6.26. *In what follows we are going to use the following generalization of these constructions: we allow a substitution of a finite number of simply-connected domains with smooth boundaries instead of disks K_i . (For such an i one should substitute any bigger domain instead of $\tilde{K}_i$.) However, we still require that the identifications φ_i are fraction-linear.*

6.8. Stratification of infinity. Consider a smooth curve γ on $\mathbb{C}P^1$ of finite length. Suppose that a metric on $\mathbb{C}P^1$ is fixed, and $z \notin \gamma$. Let $\rho_k(z, \gamma) = \|\text{dist}(z, \gamma)\|_{L_2(\gamma)}^{-k}$, here $y \in \gamma$.

Lemma 6.27. $\rho_k(x, \gamma)$ is a semicontinuous function of x , thus

$$D_{k,R} = \{z \in \mathbb{C}P^1 \mid \rho_k(x, \gamma) \leq R\}$$

is a compact subset of $\mathbb{C}P^1 \setminus \gamma$.

Let $D_k = \bigcup_{R \in \mathbb{R}} D_{k,R}$. It is a subset of $\mathbb{C}P^1 \setminus \gamma$, moreover, $\mathbb{C}P^1 \setminus \bar{\gamma} \subset D_k$. Let $\mathring{D}_{k,R} = D_{k,R} \cap (\mathbb{C}P^1 \setminus \bar{\gamma})$, $\mathring{D}_k = \bigcup_{R \in \mathbb{R}} \mathring{D}_{k,R}$. Since length of γ is finite, $\mathring{D}_0 = \mathbb{C}P^1$.

Definition 6.28. Define a filtration of $\mathcal{K} = \bar{\gamma} \setminus \gamma$ by $\mathcal{K}^{(k)} = \mathcal{K} \cap \mathring{D}_{k+1}$, $k \geq -1$. Let $\mathcal{K}^{(\infty)} = \bigcap \mathcal{K}^{(k)}$.

Suppose that γ is the boundary of a well-separated family of circles, $\gamma = \bigcup \partial K_i$.

Theorem 6.29. Let $z_0 \in \mathcal{K}^{(k)}$, z be a coordinate system near z_0 , $f \in \mathcal{H}(\mathbb{CP}^1, \{K_i\})$. Then f has an asymptotic decomposition

$$f(z) dz^{-1/2} = f_0 + f_1(z - z_0) + f_2(z - z_0)^2 + \cdots + f_k(z - z_0)^k + o(z^k)$$

when $z \rightarrow z_0$ along $\mathring{D}_{k,R}$ for an appropriate $R \gg 0$.

Proof. The half-form f is holomorphic inside $\mathbb{CP}^1 \setminus \overline{\bigcup K_i}$. The Cauchy formula show that inside $\mathring{D}_{k,R}$ the derivatives $f^{(l)}$, $l \leq k$, are bounded. Moreover, these derivatives are given by some integrals along γ , and these integrals remain well-defined in $D_{k,R}$ as well. Let f_l be the values of these integrals in $z_0 \in D_{k,R_1}$ (here $R_1 \gg 0$).

Since $z_0 \in \mathring{D}_{k+1}$, it is in the closure of $\mathring{D}_{k+1,R}$ for an appropriate R . It is easy to see that $f^{(k)}(z)$ has a limit f_k when $z \rightarrow z_0$ along $\mathring{D}_{k+1,R}$. Same is true for $f^{(l)}(z)$, $l \leq k$.

Consider the integral for

$$\frac{f(z) dz^{-1/2} - \left(f_0 + f_1(z - z_0) + f_2(z - z_0)^2 + \cdots + f_k(z - z_0)^k \right)}{(z - z_0)^k}.$$

It is

$$\int_{\gamma} K_k(z, z_0, \zeta) f(\zeta),$$

here

$$\begin{aligned} K(z, \zeta) &= \frac{d\zeta^{1/2}}{\zeta - z}, \\ K_k(z, z_0, \zeta) &= \frac{K(z, \zeta) - \sum_{l=0}^k \frac{d^l K}{dz^l} \Big|_{z=z_0} \frac{(z - z_0)^l}{l!}}{(z - z_0)^k}, \\ &= \frac{z - z_0}{(\zeta - z_0)^{k+1}} K(z, \zeta) = \frac{1}{(\zeta - z_0)^k} (K(z, \zeta) - K(z_0, \zeta)). \end{aligned}$$

We need to show that this integral goes to 0 when $z \rightarrow z_0$ along $\mathring{D}_{k+1,R}$. Consider a function $\rho(r)$ such that $\lim_{r \rightarrow 0} \rho(r) = 0$, $\lim_{r \rightarrow 0} \rho(r)/r = \infty$. Here γ_r is the intersection of γ with the disk of radius r about z_0 , instead of $\tilde{r}_{\gamma}(f) \in L_2(\gamma)$ we write just f .

Break the integral into three parts: two (which we do not want to separate yet) along $\gamma_{\rho(|z-z_0|)}$, the other along $\gamma \setminus \gamma_{\rho(|z-z_0|)}$. Since

$$K_k = \frac{z - z_0}{\zeta - z_0} \frac{K(z, \zeta)}{(\zeta - z_0)^k} = o\left(\frac{K(z_0, \zeta)}{|\zeta - z_0|^k}\right)$$

along the second part of γ if $|z - z_0| \rightarrow 0$, and since

$$\frac{K(z_0, \zeta)}{|z_0 - \zeta|^k} \in L_2(\gamma)$$

as a function of ζ , it is enough to show that the first two part of the integral go to 0. Since $\lim_{r \rightarrow 0} \int_{\gamma_{\rho(r)}} |f(\zeta)|^2 = 0$, we need to show only that the L_2 -norm of $K_k(z, z_0, \zeta)$ is bounded when $\zeta \in \gamma_{\rho(|z-z_0|)}$.

Subdivide $\gamma_{\rho(|z-z_0|)}$ once more: into $\gamma_{\varepsilon|z-z_0|}$ and $\gamma_{\rho(|z-z_0|)} \setminus \gamma_{\varepsilon|z-z_0|}$. Here $\varepsilon \ll 1$. Since on the second part $K(z_0, \zeta) = O(K(z, \zeta))$, we see that $K_k(z, z_0, \zeta) = O\left(\frac{K(z, \zeta)}{(\zeta - z)^k}\right)$, thus K_k has a bounded L_2 -norm. We conclude that the integral along the second part goes to 0 when $z \rightarrow z_0$. On the first part $K(z, \zeta) = O(K(z_0, \zeta))$, thus

$$K_k(z, z_0, \zeta) = \frac{1}{(\zeta - z_0)^k} (K(z, \zeta) - K(z_0, \zeta)) = O\left(\frac{K(z_0, \zeta)}{(\zeta - z_0)^k}\right) = O\left(\frac{1}{(z - z_0)^{k+1}}\right),$$

so it has a bounded L_2 -norm as well. $\square$

Definition 6.30. Consider a model $\bar{M}$ of a curve. Say that a point $z \in M_\infty$ is of smoothness C^k , if $z \in \mathcal{K}^{(k)}$, here $\mathcal{K} = M_\infty$.

Remark 6.31. Note that the stratification we used is related to the following inclusion of M into $L_2(\gamma, \Omega^{1/2} \otimes \mu)$:

$$z \mapsto \tilde{r}_\gamma(K_z),$$

here $K_z(\zeta) = \frac{d\zeta^{1/2}}{\zeta - z}$ is the Cauchy kernel. In other words,

$$z \mapsto \frac{d\zeta^{1/2}}{\zeta - z}|_\gamma$$

after a choice of coordinate ζ on $\mathbb{CP}^1$. The points of smoothness C^0 correspond to limit points of this inclusion, the points of smoothness C^k correspond to limit points of k -jets continuation of this mapping.

7. RIEMANN–ROCH THEOREMS

7.1. Abstract Riemann–Roch theorem. We say that two vector subspaces V_1, V_2 of a topological vector space H satisfy the Riemann–Roch theorem if $\dim V_1 \cap V_2 < \infty$, and $\operatorname{codim} (\overline{V_1 + V_2}) < \infty$. We call the number

$$\dim V_1 \cap V_2 - \operatorname{codim} (\overline{V_1 + V_2})$$

the *index* of two subspaces. If $V_1 + V_2 = \overline{V_1 + V_2}$, we say that V_1, V_2 satisfy the *strong form* of the theorem.

Remark 7.1. Note that if V_1, V_2 satisfy the strong form of the theorem, then the natural mapping $V_1 \rightarrow V/V_2$ is a Fredholm mapping with the index being the index of V_1, V_2 . If V_1, V_2 satisfy the weak form of the theorem, then this mapping is a continuous mapping p with $\dim \operatorname{Ker} p - \dim \operatorname{Coker} p$ being the index of V_1, V_2 . Here Coker is the quotient by the closure of the image.

Consider a direct sum of two Hilbert spaces $H = H_1 \oplus H_2$. Consider two closed vector subspaces $V_1, V_2 \subset H$ such that the projection of V_i on H_i has no null-space and a dense image. This means that one can consider V_1 as a graph of a mapping $A_1: H_1 \rightarrow H_2$, similarly V_2 is a graph of $A_2: H_2 \rightarrow H_1$. Mappings $A_{1,2}$ are closed, but not necessarily bounded.

Lemma 7.2 (abstract finiteness). *If A_2 is bounded, and $A_1 \circ A_2$ is compact, then $V_1 \cap V_2$ is finite dimensional.*

Proof. The projection of $V_1 \cap V_2$ to H_2 is a subspace of $\operatorname{Ker} (A_1 \circ A_2 - \mathbf{1})$, thus is finite-dimensional. $\square$

Proposition 7.3 (strong form). *If A_1, A_2 are bounded, and $A_1 \circ A_2$ is compact, then V_1 and V_2 satisfy the strong form of Riemann–Roch theorem with index 0.*

Proof. The projection of $V_1 \cap V_2$ to H_2 is $\operatorname{Ker} (A_1 \circ A_2 - \mathbf{1})$, which implies the statement about $\dim V_1 \cap V_2$. The statement about $\operatorname{codim} \overline{V_1 + V_2}$ follows from the fact that the orthogonal complements to V_1 and V_2 satisfy the same conditions as V_2 and V_1 with linear mappings being $-A_1^*, -A_2^*$.

To show that $\dim V_1 \cap V_2 = \operatorname{codim} (\overline{V_1 + V_2})$ note that $\operatorname{Ker} (A_1 \circ A_2 - \mathbf{1})$ is dual to $\operatorname{Coker} (A_2^* \circ A_1^* - \mathbf{1})$. $\square$

Proposition 7.4 (weak form). *If A_2 is bounded, and both $A_1 \circ A_2$ and $A_1^* \circ A_2^*$ are compact, then V_1 and V_2 satisfy the Riemann–Roch theorem with index 0.*

Proof. We already know that $\dim V_1 \cap V_2$ and $\operatorname{codim} \overline{V_1 + V_2}$ are finite. The only thing to prove is that

$$\dim \operatorname{Ker} (A_1 \circ A_2 - \mathbf{1}) = \dim \operatorname{Coker} (A_1^* \circ A_2^* - \mathbf{1}).$$

Obviously, $(A_1^* \circ A_2^*)^*$ is the closure of $A_2 \circ A_1$, thus

$$A_2(\text{Ker}(A_1 \circ A_2 - \mathbf{1})) \subset \text{Ker}((A_1^* \circ A_2^*)^* - \mathbf{1}),$$

hence

$$\dim \text{Ker}(A_1 \circ A_2 - \mathbf{1}) \leq \dim \text{Ker}((A_1^* \circ A_2^*)^* - \mathbf{1}) = \dim \text{Coker}(A_1^* \circ A_2^* - \mathbf{1}).$$

Application of the same argument to the dual operators shows the opposite inequality. $\square$

We say that two vector subspaces V, V' of a vector space H are *comparable*, if $V \cap V'$ is of finite codimension in both V and V' . The *relative dimension* $\text{reldim}(V, V')$ is $\text{codim}(V \cap V' \subset V) - \text{codim}(V \cap V' \subset V')$. The following theorem is a direct corollary of the above statement:

Theorem 7.5 (Riemann–Roch theorem). *Consider two vector subspaces $V_{1,2} \subset H$ of a Hilbert space $H = H_1 \oplus H_2$. Suppose that V_1 is comparable with the graph of a closed mapping $A_1: H_1 \rightarrow H_2$ and the relative dimension of V_1 and this graph is d_1 . Suppose V_2 is comparable with the graph of a closed mapping $A_2: H_2 \rightarrow H_1$ and the relative dimension of V_2 and this graph is d_2 .*

1. **(weak form)** *If A_2 is bounded, and both $A_1 \circ A_2$ and $A_1^* \circ A_2^*$ are compact, then V_1 and V_2 satisfy the Riemann–Roch theorem with the index being $d_1 + d_2$.*
2. **(strong form)** *If both A_1 and A_2 are bounded, and $A_1 \circ A_2$ is compact, then V_1 and V_2 satisfy the strong form of Riemann–Roch theorem with the index being $d_1 + d_2$.*

7.2. Riemann problem. Consider a holomorphic function $\psi(z)$ defined in an annulus $U = \{z \mid 1 - \varepsilon \leq |z| \leq 1 + \varepsilon\}$, let $S^1 = \{z \mid |z| = 1\}$. Suppose that $\psi(z)$ is nowhere 0, and consider the subspace V_ψ in $L_2(S^1, \omega^{1/2}) \oplus L_2(S^1, \omega^{1/2})$ consisting of pairs of the form $(f(z), \psi(z)f(z))$. The Hilbert space $L_2(S^1, \omega^{1/2})$ is a direct sum of subspaces $L_2^\pm(S^1, \omega^{1/2})$ consisting of forms which can be holomorphically continued into two regions S^1 divides $\mathbb{CP}^1$ into.

Let p be the projection of V_ψ to

$$L_2^+(S^1, \omega^{1/2}) \oplus L_2^-(S^1, \omega^{1/2}) \subset L_2(S^1, \omega^{1/2}) \oplus L_2(S^1, \omega^{1/2}).$$

Lemma 7.6. *If $\text{ind } \psi = 0$, then p is invertible.*

Proof. Consider a linear bundle $\mathcal{L}$ over $\mathbb{CP}^1$ with isomorphisms to $\omega^{1/2}$ over $U^+ = \{|z| < 1 + \varepsilon\}$ and $U^- = \{|z| > 1 - \varepsilon\}$, and the gluing data being $l^+ = \psi l^-$. Since $\deg \mathcal{L} = \deg \omega^{1/2} + \text{ind } \psi$, and a linear bundle over $\mathbb{CP}^1$ is determined by its degree up to an isomorphism, we see that $\mathcal{L} \simeq \omega^{1/2}$ if $\text{ind } \psi = 0$.

Since $\text{Ker } p$ consists of global sections of $\mathcal{L}$, $\text{Ker } p = \{0\}$. Similarly, consideration of orthogonal complement to V_ψ shows that $\text{Coker } p = \{0\}$. This finishes the proof, since the operator is obviously Fredholm. $\square$

Consider now another linear bundle $\mathcal{L}'$ with trivializations over U^+ and over U^- with gluing data $l^+ = \psi l^-$. The same arguments as above show that $\mathcal{L}'$ is trivial, thus it has a (unique up to multiplication by a constant) global section. This means that there are functions $l^\pm$ defined on $U^\pm$ such that $l^+ = \psi l^-$. Since this global section has no zeros, $l^\pm$ have no zeros inside the domain of definition, thus $\psi = (l^+)^{-1} l^-$. We see that any function ψ such that $\text{ind } \psi = 0$ can be represented as a product $\psi = \psi_+ \psi_-$ of a parts ψ_+ , ψ_- which can be holomorphically extended inside/outside a circle without zeros.

Let us write the mapping p^{-1} in terms of $\psi_\pm$. For any L_2 -section ω of $\Omega^{1/2} \otimes \mu$ on $\{|z| = 1\}$ let $\omega = \omega_+ + \omega_-$ be the (unique) decomposition of ω into a sum of forms which can holomorphically extended inside/outside of the unit circle. Given ω_+ and ω'_- we want to find ω_- and ω'_+ from the equality

$$\psi(\omega_+ + \omega_-) = \omega'_+ + \omega'_-, \quad \text{or} \quad \psi_-(\omega_+ + \omega_-) = \psi_+^{-1}(\omega'_+ + \omega'_-).$$

Taking the $+$ -part we see that $(\psi_-\omega_+)_+ = \psi_+^{-1}\omega'_+ + (\psi_+^{-1}\omega'_-)_+$, thus

$$\omega'_+ = \psi_+(\psi_-\omega_+)_+ - \psi_+(\psi_+^{-1}\omega'_-)_+ = \psi\omega_+ - \psi_+(\psi_-\omega_+)_- - \psi_+(\psi_+^{-1}\omega'_-)_+,$$

similarly $(\psi_-\omega_+)_- + \psi_-\omega_- = (\psi_+^{-1}\omega'_-)_-$, thus

$$\omega_- = \psi_+^{-1}(\psi_+^{-1}\omega'_-)_- - \psi_+^{-1}(\psi_-\omega_+)_- = \psi_+^{-1}\omega'_- - \psi_+^{-1}(\psi_+^{-1}\omega'_-)_+ - \psi_+^{-1}(\psi_-\omega_+)_-.$$

These two formulae express ω_- and ω'_+ in terms of ω'_- and ω_+ , thus give an inverse mapping to p .

Let $|\psi|_{++} = \frac{\max|\psi_+|}{\min|\psi_+|}$, $|\psi|_{--} = \frac{\max|\psi_-|}{\min|\psi_-|}$, $|\psi|_{+-} = \max|\psi_+| \max|\psi_-|$, $|\psi|_{-+} = \max|\psi_+|^{-1} \max|\psi_-|^{-1}$, $|\psi|_0 = \max(|\psi|_{++}, |\psi|_{--}, |\psi|_{+-}, |\psi|_{-+})$. Then the norm of p^{-1} is bounded by $C \cdot |\psi|_0$.

Definition 7.7. The *Riemann norm* $\|\psi\|_{\mathbf{R}}$ of ψ is the norm of the operator p^{-1} .

The following lemma is a corollary of the fact that one can find factorization $\psi = \psi_+ \psi_-$ using integral operators applied to $\log \psi$:

Lemma 7.8. *Let $\log \psi(z)$ is defined using any branch of logarithm. Then*

$$\|\psi\|_{\mathbf{R}} < C \exp C \max_{z \in U} |\log \psi(z)|$$

for an appropriate C (which depends on ε only).

Let Π_ψ be the composition of p^{-1} with the projection of $L_2(S^1, \omega^{1/2}) \oplus L_2(S^1, \omega^{1/2})$ to $L_2^-(S^1, \omega^{1/2}) \oplus L_2^+(S^1, \omega^{1/2})$,

$$\Pi_\psi: L_2^+(S^1, \omega^{1/2}) \oplus L_2^-(S^1, \omega^{1/2}) \rightarrow L_2^-(S^1, \omega^{1/2}) \oplus L_2^+(S^1, \omega^{1/2}).$$

In other words, $((f_+, g_+), (f_-, g_-))$ lies on the graph of Π_ψ if $g_+ + g_- = \psi(f_+ + f_-)$.

We see that the norm of Π_ψ is bounded by $C \cdot |\psi|_0$. Moreover, Π_ψ can be written as a sum

$$(7.1) \quad \Pi_\psi = \begin{pmatrix} 0 & \psi^{-1} \\ \psi & 0 \end{pmatrix} + k.$$

Obviously, k is compact (as any Hankel operator). Indeed, components of k look like $f_+ \mapsto (mf_+)_-$. The Hilbert operator $f = f_+ + f_- \xrightarrow{H} f_+ - f_-$ is a pseudodifferential operator of degree 0, thus components of K may be written as $[m, H]$, thus are pseudodifferential operators of degree -1 , thus compact.

Lemma 7.9. *Consider an invertible function ψ defined in a neighborhood of $|z| = 1$ and having $\text{ind} = k$. Let $V_\psi \subset H = L_2(S^1) \oplus L_2(S^1)$ be*

$$V_\psi = \{(f_1, f_2) \mid f_2 = \psi f_1\}.$$

Define $H^\pm = L_2(S^1)^\pm \oplus L_2(S^1)^\mp$. Then for an appropriate bounded operator $\pi: H^+ \rightarrow H^-$ the graph (π) is compatible with V_ψ , and $\text{reldim}(H_2, \text{graph}(\pi)) = k$.

Proof. Let $\psi(z) = z^k \psi_0(z)$. The function ψ_0 has $\text{ind} = 0$, thus the corresponding subspace V_{ψ_0} is the graph of Π_{ψ_0} . Let $H_0^\pm = L_2(S^1)^\pm \oplus z^k L_2(S^1)^\mp$. We see that V_ψ is a graph of a bounded mapping $H_0^+ \rightarrow H_0^-$. Since H_0^+ is compatible with H_0 of relative dimension k , we momentarily obtain the required statement about V_ψ . $\square$

Remark 7.10. Note that $\|\pi\|$ may be bounded in the same way as in Lemma 7.8, but neither this result, nor (7.1) are going to be needed in what follows.

7.3. Finiteness theorem. Consider a family of disks and gluing conditions $K_\bullet, \varphi_\bullet, \psi_\bullet$ from Section 6.4. Let $\mathcal{H}^\pm = \bigoplus_{i \in I} L_2^\pm(\partial K_i, \Omega^{1/2} \otimes \mu)$, $i \in I$. Then the operator with Cauchy kernel defines a Hilbert mapping $\mathbf{K}: \mathcal{H}^- \rightarrow \mathcal{H}^+$ (see Section 6.2). This mapping depends on the circles ∂K_i only, not on the gluing conditions $\varphi_\bullet, \psi_\bullet$.

Since the Hilbert structure on $L_2(\partial K_i, \Omega^{1/2} \otimes \mu)$ is invariant w.r.t. fraction-linear mappings (as is decomposition into $\pm$ -parts), the identification $\varphi_i: \partial K_{i'} \rightarrow \partial K_i$ gives an isomorphism of Hilbert spaces $L_2(\partial K_i, \Omega^{1/2} \otimes \mu)$ and $L_2(\partial K_{i'}, \Omega^{1/2} \otimes \mu)$ which interchanges $+$ -part and $-$ -part.

Suppose that $\text{ind } \psi_i = 0$. Then the operator Π_{ψ_i} from Section 7.2 together with an identification given by φ_i gives a mapping Π_{φ_i, ψ_i}

$$L_2^+(\partial K_i, \Omega^{1/2} \otimes \mu) \oplus L_2^+(\partial K_{i'}, \Omega^{1/2} \otimes \mu) \rightarrow L_2^-(\partial K_i, \Omega^{1/2} \otimes \mu) \oplus L_2^-(\partial K_{i'}, \Omega^{1/2} \otimes \mu),$$

the graph of this mapping consists of half-forms on ∂K_i and $\partial K_{i'}$ which differ by multiplication by ψ_i . In other words, $((f_+, g_+), (f_-, g_-))$ lies on this graph if

$$(g_+ + g_-)(\varphi_i^{-1}(z)) = \psi_i(z)(f_+ + f_-)(z), \quad z \in \partial K_i.$$

If $\text{ind } \psi_i \neq 0$, instead of Π_{φ_i, ψ_i} consider an arbitrary operator between the same spaces such that the graph of P is comparable with the set of half-forms which differ

by multiplication by ψ_i (see Lemma 7.9). The only fact important in what follows is that this is a bounded operator (see Corollary 6.6).

Let $\Pi = \Pi_{\{\psi\}} = \bigoplus_{i \in I_+} \Pi_{\varphi_i, \psi_i} : \mathcal{H}^+ \rightarrow \mathcal{H}^-$. Similarly, let $\Pi_{\{\psi^{-1}\}}$ corresponds to gluing data φ_i, ψ_i^{-1} . Since Π consists of bounded diagonal blocks, it is a closed operator.

Let $\bar{M}$ be a curve determined by gluing conditions $\varphi_\bullet$, $\mathcal{L}$ be a bundle on $\bar{M}$ determined by gluing conditions $\psi_\bullet$.

Theorem 7.11. *Suppose that the disks K_i are well separated, $\mathcal{L}$ is a finite-degree bundle, and $\Pi_{\{\psi\}} \circ \mathbf{K}$ is compact. Then both $\Gamma_{\text{strong}}(\bar{M}, \omega^{1/2} \otimes \mathcal{L})$ and $\Gamma_{\text{weak}}(\bar{M}, \omega^{1/2} \otimes \mathcal{L})$ are finite-dimensional.*

Proof. We give only a sketch of a proof, since the details are the same as in the case of Riemann–Roch theorem (see Section 7.5). Since Γ_{strong} is identified with a subspace of Γ_{weak} , it is enough to show that $\dim \Gamma_{\text{weak}} < \infty$. On the other hand, Γ_{weak} is defined as an intersection of the generalized Hardy space $\mathcal{H}(\mathbb{CP}^1, \{K_i\})$ with the subspace of forms which satisfy the gluing conditions. We are going to reduce the statement of the theorem to the abstract finiteness theorem.

To do this, note that Proposition 6.12 identifies $\mathcal{H}(\mathbb{CP}^1, \{K_i\})$ with the graph of $\mathbf{K} : \mathcal{H}^- \rightarrow \mathcal{H}^+$, thus the only thing to note is that fact that the subspace of forms which satisfy the gluing conditions is compatible with the graph of $\Pi_{\{\psi\}}$. $\square$

7.4. Duality. Consider the curve $\bar{M}$ and a sheaf $\mathcal{L}$ from the previous section and the dual sheaf $\mathcal{L}^{-1}$ with inverse gluing conditions $\{\psi_i^{-1}\}$.

Theorem 7.12. *Suppose that the disks K_i are well-separated, and $\mathcal{L}$ is of finite degree. The mappings*

$$H^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) \xrightarrow{\bar{\partial}} H^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega})$$

and

$$H^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}^{-1}) \xrightarrow{-\bar{\partial}} H^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}^{-1} \otimes \bar{\omega})$$

are mutually dual, thus dimension of null-space of one mapping is equal to the dimension of cokernel of another one.

Proof. This is a direct corollary of Lemmas 6.17 and 6.18. $\square$

Using Theorems 7.12 and 6.21 together with selfduality of $\mathbf{K}$ and the fact that $\Pi_{\psi}^t = \Pi_{\psi^{-1}}$, we obtain

Corollary 7.13. *Suppose that the disks K_i are well separated, and $\mathcal{L}$ is finite-degree and semibounded, and both $\Pi_{\{\psi\}} \circ \mathbf{K}$ and $\Pi_{\{\psi^{-1}\}} \circ \mathbf{K}$ are compact. Then the bounded operator*

$$H_{\text{weak}}^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) \xrightarrow{\bar{\partial}} H_{\text{weak}}^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega})$$

has finite-dimensional null-space and cokernel.

7.5. Riemann–Roch for curves. Now we are ready to state

Theorem 7.14 (Riemann–Roch). *Suppose that the disks K_i are well-separated, and ψ_i is a finite-degree family.*

1. *If both $\Pi_{\{\psi\}} \circ \mathbf{K}$ and $\Pi_{\{\psi^{-1}\}} \circ \mathbf{K}$ are compact operators then the mapping*

$$H_{weak}^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) \xrightarrow{\bar{\partial}} H_{weak}^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega})$$

has finite-dimensional null-space and cokernel, and

$$\dim \text{Ker } \bar{\partial} - \dim \text{Coker } \bar{\partial} = \sum_{i \in I_+} \text{ind } \psi_i.$$

2. *If $\Pi_{\{\psi\}}$ is bounded, and $\mathbf{K}$ is compact, then the mapping is Fredholm of index $\sum_{i \in I_+} \text{ind } \psi_i$.*

Proof. The only thing we need to do is to describe the null-space and the image of $\bar{\partial}$. We are going to reduce this description to the abstract Riemann–Roch theorem. First of all, the null-space consists of half-forms in the generalized Hardy space, so $\tilde{r}$ maps it injectively to a subspace of $\bigoplus_{l_2} L_2(\partial K_i, \Omega^{1/2} \otimes \mu)$. On the other hand, the image of $\text{Ker } \bar{\partial}$ in $\bigoplus_{l_2} L_2(\partial K_i, \Omega^{1/2} \otimes \mu)$ is described as intersection of the image of the generalized Hardy space and a subspace in $\bigoplus_{l_2} L_2(\partial K_i, \Omega^{1/2} \otimes \mu)$ consisting of half-forms which satisfy the gluing conditions.

The first condition can be written as

$$f_i^+ = \sum_{j \neq i} \mathbf{K}_{ij} f_j^-,$$

here $f_i^\pm$ are $\pm$ -parts of the component of f in $L_2(\partial K_i, \Omega^{1/2} \otimes \mu)$. The second condition is

$$f_{i'}' = \psi_i f_i.$$

Here $f(t)' = f(-t)$, and we suppose that we use compatible parameterizations of ∂K_i and of $\partial K_{i'}$, i.e., such that φ_i send parameter t on ∂K_i to parameter $-t$ on $\partial K_{i'}$.

Consider the decomposition $\mathcal{H} = \mathcal{H}^+ \oplus \mathcal{H}^-$. Vectors which satisfy the first condition are in the graph of operator $\mathbf{K}: \mathcal{H}^- \rightarrow \mathcal{H}^+$. Consider the vector space of vectors $\mathcal{H}_2$ which satisfy the second conditions. It is a direct sum over $i \in I_+$ of subspaces of

$$L_2(\partial K_i, \Omega^{1/2} \otimes \mu) \oplus L_2(\partial K_{i'}, \Omega^{1/2} \otimes \mu),$$

and all the components but a finite number have ψ with $\text{ind} = 0$, thus are described as graphs of Π_{φ_i, ψ_i} . As a corollary, we conclude that this subspace is compatible with the graph of the mapping Π .

To finish the description of the null-space the only thing which remains to prove is to show that the relative dimension of $\mathcal{H}_2$ and $\text{graph}(\Pi)$ is $\sum_{i \in I_+} \text{ind } \psi_i$. However, because of decomposition of $\mathcal{H}_2$ into a direct sum it follows from the corresponding fact for each component, i.e., from Lemma 7.9.

To describe the image, consider

$$\alpha \in H_{\text{weak}}^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega}) = H^{-1/2}\left(\mathbb{CP}^1 \setminus \bigcup K_i, \omega^{1/2} \otimes \bar{\omega}\right).$$

Since $\bar{\partial}$ is an isomorphism

$$H^{1/2}(\mathbb{CP}^1, \omega^{1/2}) \rightarrow H^{-1/2}(\mathbb{CP}^1, \omega^{1/2} \otimes \bar{\omega}),$$

the element $\bar{\partial}^{-1}\alpha \in H^{1/2}(\mathbb{CP}^1, \omega^{1/2})$ is defined up to addition of an element of $\bar{\partial}^{-1} \mathring{H}^{-1/2}(\bigcup K_i, \omega^{1/2} \otimes \bar{\omega})$. On the other hand, the latter space is $\mathcal{H}(\mathbb{CP}^1, \{K_i\})$, i.e., $\bar{\partial}^{-1}$ gives a correctly defined isomorphism

$$H^{-1/2}\left(\mathbb{CP}^1 \setminus \bigcup K_i, \omega^{1/2} \otimes \bar{\omega}\right) \rightarrow H^{1/2}(\mathbb{CP}^1, \omega^{1/2}) / \mathcal{H}(\mathbb{CP}^1, \{K_i\}),$$

hence $\bar{\partial}^{-1}\alpha$ is an element of the latter space. Thus

$$\alpha \in \text{Im} \left(H_{\text{weak}}^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) \xrightarrow{\bar{\partial}} H_{\text{weak}}^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega}) \right)$$

is equivalent to

$$\bar{\partial}^{-1}\alpha \in \mathcal{H}_2 / \mathcal{H}(\mathbb{CP}^1, \{K_i\}) \stackrel{\text{def}}{=} (\mathcal{H}(\mathbb{CP}^1, \{K_i\}) + \mathcal{H}_2) / \mathcal{H}(\mathbb{CP}^1, \{K_i\}).$$

In particular, if $\mathcal{H}(\mathbb{CP}^1, \{K_i\}) + \mathcal{H}_2$ is closed, then $\text{Im } \bar{\partial}$ is closed, and in any case the codimension of the closure of $\text{Im } \bar{\partial}$ is equal to codimension of the closure of $\mathcal{H}(\mathbb{CP}^1, \{K_i\}) \oplus \mathcal{H}_2$. $\square$

From Lemma 7.8 we momentarily obtain

Corollary 7.15. *Suppose that $\mathbf{K}$ is compact, and $\psi_i = \exp \Phi_i$ for all i but a finite number. If for an appropriate C and any i*

$$|\Phi_i| < C,$$

then the operator

$$H_{\text{weak}}^{1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L}) \xrightarrow{\bar{\partial}} H_{\text{weak}}^{-1/2}(\bar{M}, \omega^{1/2} \otimes \mathcal{L} \otimes \bar{\omega})$$

is Fredholm.

Definition 7.16. Call a pair $(\bar{M}, \mathcal{L})$ *admissible* if $\bar{M}$ is given by a family of well-separated disks, $\mathcal{L}$ is of finite degree, and both $\Pi_{\{\psi\}} \circ \mathbf{K}$ and $\Pi_{\{\psi^{-1}\}} \circ \mathbf{K}$ are compact. Call $\bar{M}$ *admissible* if $(\bar{M}, \mathbf{1})$ is admissible. Here $\mathbf{1}$ is a bundle over $\bar{M}$ with $\psi_i \equiv 1$.

7.6. Criterion of admissibility of a curve. Recall that in Section 6.1 we considered a matrix $(e^{-l_{ij}/2})$ which was supposed to give a bounded operator in l_2 .

Proposition 7.17. *If the matrix $(e^{-l_{ij}/2} - \delta_{ij})$ gives a compact operator $l_2 \rightarrow l_2$, then $\bar{M}$ is admissible.*

Proof. Indeed, for the bundle $\mathcal{L} = \mathbf{1}$ the operator Π is an isometry, thus we need to show that $\mathbf{K}$ is compact, which is a corollary of Lemma 6.5. $\square$

Corollary 7.18. *If $\sum_{i \neq j} e^{-l_{ij}} < \infty$, then $\bar{M}$ is admissible.*

Corollary 7.19. *Fix a locally discrete subset $I \subset \mathbb{CP}^1$ (i.e., for any point i of I there is a punctured neighborhood of i which does not intersect I). Then there exists a family of disks K_i , $i \in I$ with center $(K_i) = i$ such that for any involution $'$ and any gluing data φ_i the corresponding curve $\bar{M}$ is admissible. Moreover, one can chose K_i in such a way that M_∞ consists of points of smoothness C^∞ (see Section 6.8).*

Corollary 7.20. *For an arbitrary nowhere dense subset N of $\mathbb{CP}^1$ there exists an admissible curve $\bar{M}$ such that $M_\infty = N$. Moreover, it is possible to make every point of N to be of smoothness C^∞ .*

Remark 7.21. While the above statements are obvious, note that construction of examples and counterexamples may be simplified a lot by an additional restriction:

$$\text{dist}(i, j) \geq \varepsilon \cdot \text{dist}(i, N) \quad \text{for any } i, j \in I, i \neq j.$$

Here $\varepsilon \ll 1$. To construct such a family I for a given nowhere dense set N , let $N_k = \{z \mid 2^{-k-1} \leq \text{dist}(z, N) \leq 2^{-k}\}$. Fix k , and let $\delta = 2^{-k}$. Consider a $\delta/8$ -net for N_k . By removing some points from this net one can obtain $\delta/4$ -net such that it does not have two points closer than $\delta/8$. This net is necessarily finite.

Now consider the union of these finite sets over $k \in \mathbb{N}$, and again remove net points from N_{k+1} which are closer than $\delta/8$ to net points in N_k . One obviously obtains a set I with required properties.

Now to chose the radius of K_i denote by n_k the number of points in $I \cap N_k$. Let radius $(K_i) = \frac{f(k)}{n_k}$ if $i \in N_k$, here $f(k)$ is rapidly decreasing function. Picking appropriate $f(k)$, one obtains disks which satisfy the given above requirements.

The following property of admissibility is obvious:

Fact . *If we change a finite number of contours ∂K_i and/or a finite numbers of identifications φ_i , this does not change the admissibility of the resulting curve.*

7.7. Filling the gap. This section contains heuristics only, so anyone interested exceptionally in exact results should proceed directly to Section 8.

Consider once more the criterion of admissibility of a curve. It says that if the operator $\mathbf{K}$ is compact, the curve is admissible (together with any bundle which is defined by gluing functions ψ_i with bounded Riemann norm $|\psi|_{\mathbf{R}}$). On the other hand, to obtain this result we study the generalized Hardy space, which is correctly defined if $\mathbf{K}$ is bounded.

Thus the tools we use leave a gap between the objects for which the analysis is applicable (i.e., $\mathbf{K}$ is bounded), and objects for which we get the admissibility (i.e., $\mathbf{K}$ is compact). How can we use the existence of this gap?

We propose to consider this gap as a confirmation that the Riemann–Roch theorem we obtained is *almost unimprovable*. Indeed, we know that the finiteness condition can be written as $\dim \text{Ker}(\Pi \circ \mathbf{K} - 1) < \infty$. On the other hand, the invertible operator Π depends on the family (ψ_i) which has very high degree of freedom (even if we consider the strong form of Riemann–Roch theorem, so Π is required to be bounded), and one should expect that the condition

$$\dim \text{Ker}(\Pi \circ \mathbf{K} - 1) < \infty \text{ for every choice of } \psi_i \text{ with } |\psi|_{\mathbf{R}} < \infty$$

should be very close to the condition

$$(7.2) \quad \mathbf{K} \text{ has discrete spectrum near } |\lambda| = 1.$$

In other words, the pairs $(\Pi, \mathbf{K})$ with an invertible Π , bounded $\mathbf{K}$, and *infinite-dimensional* null-space of $\Pi\mathbf{K} - 1$ are plentiful (at least if we drop “geometric” conditions on Π and $\mathbf{K}$, and consider abstract operators), and they form a “natural boundary” of the set of curves for which Riemann–Roch theorem has a chance to be true. This natural boundary is quite close to the boundary of the set of compact operators, which is another confirmation of our thesis.

Note that (7.2) may lead to a different description of possible $\mathbf{K}$, like $\frac{\mathbf{K}}{1+\mathbf{K}^2}$ being compact. It is unclear, however, whether the above boundary separates the set of compact operators as a connected component of the set of operators $\mathbf{K}$ with compact $\frac{\mathbf{K}}{1+\mathbf{K}^2}$.

7.8. Moduli space. The description of a complex curve by disks $\{K_i\}$, involution σ and gluings φ_i leaves a feeling of being incomplete, since in the case of finite genus it is enough to provide just gluings φ_i which generate a subgroup of $\text{SL}(2, \mathbb{C})$. To describe the quotient by this subgroup one can take any fundamental domain for the subgroup, and different choices of the fundamental domain result in the same geometric data.

To get a similar description in the case of infinite genus, note that in Section 6.1 instead of the restriction that $\tilde{K}_i$ is a concentric with K_i disk of radius $e^{2\varepsilon} \cdot \text{radius}(K_i)$ one can require that $\tilde{K}_i$ *contains* such a circle. This leads to the construction of strong

sections of $\omega^{1/2} \otimes \mathcal{L}$ in the same way as in Section 6.5, and Theorem 6.21 can be refined as

Amplification 7.22. *Fix a metric on $\mathbb{CP}^1$ and $\varepsilon > 0$. Consider a family of elements φ_i of $\mathrm{SL}(2, \mathbb{C})$. Suppose that there exists a family of disjoint domains $\tilde{K}_i \subset \mathbb{CP}^1$, $i \in I$, which satisfy the following properties:*

1. *All $\tilde{K}_i$ but a finite number are disks;*
2. *Let K'_i be a concentric with $\tilde{K}_i$ disk of radius $(1 - \varepsilon)$ radius $(\tilde{K}_i)$ (or any domain in $\tilde{K}_i$ if $\tilde{K}_i$ is not a disk). Suppose that for an involution $\prime: I \rightarrow I$ one has $\varphi_{i'} = \varphi_i^{-1}$, $\varphi_i(K'_i) \cup K'_{i'} = \mathbb{CP}^1$;*
3. *Let ψ_i be holomorphic functions defined inside $\tilde{K}_i \cap \varphi_i^{-1}(\tilde{K}_{i'})$ which satisfy $\psi_{i'} \circ \varphi_i = \psi_i^{-1}$, and $\mathrm{ind} \psi_i = 0$ for all but a finite number of $i \in I$.*
4. *Suppose that the pairwise conformal distances l_{ij} between $\tilde{K}_i, \tilde{K}_j$ satisfy the condition that the matrix $(e^{-l_{ij}/2} - \delta_{ij})$ gives a compact mapping $l_2 \rightarrow l_2$, and all the functions ψ_i are bounded taken together.*

Let $\bar{M}$ be a curve given by gluing $S = \bigcap \varphi_i^{-1}(K'_i)$ together via φ_i , $\mathcal{L}$ be a bundle on $\bar{M}$ given by cocycle ψ_i . Define strong sections of $\omega^{1/2} \otimes \mathcal{L}$ associated with family $\{\tilde{K}_i\}$ as forms α in $H^{1/2}(\bigcap \varphi_i^{-1}(K'_i), \omega^{1/2})$ which satisfy $\varphi_i^(\alpha) = \psi_i \cdot \alpha$ whenever both sides have sense.*

Let $I = I_+ \amalg I'_+$. For any choice of circles γ_i in $K'_i \cap \varphi_i^{-1}(K'_{i'})$, $i \in I_+$, let K_i be the disk bounded by γ_i inside K'_i (with appropriate modifications if K_i is not a disk). Let $K_{i'} = \mathbb{CP}^1 \setminus \varphi_i(K_i)$, $i \in I_+$. Then the space of strong sections of $\omega^{1/2} \otimes \mathcal{L}$ coincides with the set of weak sections associated to the family $\{K_i\}$.

In particular, the Riemann–Roch theorem (in the strong form) is valid for strong sections associated to the family $\{\tilde{K}_i\}$.

The proof of this statement is a corollary of the proof of Theorem 6.21.

We see that at least for bundles described by bounded cocycles one does not need to specify *precisely* the circles which cut the curve, one can vary them in wide ranges (which depend on the Kleinian group) without any change to the geometric data. One possible objection to usability of the above theorem is that for different choices $\{\gamma_i^{(1)}\}, \{\gamma_i^{(2)}\}$ of the circles the cocycles ψ_i which describe the bundle need to be defined everywhere between the circles for the theorem to be applicable. However, in Section 9.1 we will see that any bounded bundle of degree 0 may be described by constant cocycles, thus this objection becomes void. (If the degree is not 0, one can take all the cocycles to be constant except one pair described by rational functions.)

Call a Kleinian group *admissible* if it has generators φ_i , $i \in I_+$, which satisfy the conditions of the amplification. The following conjecture would show that the curve (with a fixed family of A -cycles) is completely described by the corresponding

Kleinian group, and the restriction on distances between $\tilde{K}_i$ is in fact the restriction on the Kleinian group.

Consider an admissible Kleinian group. A *fundamental family* is a family of domains $\tilde{K}_\bullet$ which satisfies the conditions of amplification. If $\tilde{K}_i^{(1)} \subset \tilde{K}_i^{(2)}$, $i \in I$, then call these families *equivalent*, and continue this relation by transitivity. The amplification shows that equivalent families lead to the same spaces of sections of bundles.

Conjecture 7.23. Any two fundamental families for an admissible Kleinian group are equivalent.

Let the *fine moduli space* be the set of admissible Kleinian groups up to conjugation in $\mathrm{PGL}(2, \mathbb{C})$.

Conjecture 7.24. Consider a complex curve M . Let $\gamma_i^{(1)}$, $i \in I^{(1)}$, be a disjoint family of smooth embedded cycles in M such that $M \setminus \bigcup \gamma_i$ is conformally equivalent to a fundamental domain of an admissible Kleinian group. Let $\gamma_i^{(2)}$ be a different family which satisfies same conditions. Then all the cycles $\gamma_i^{(1)}$ except a finite number are homotopic to cycles in $\gamma_i^{(2)}$.

Conjecture 7.25. Two families of cuts from the previous conjecture lead to the same set of C^n -points at infinity (see Section 6.8) for any $n \geq 0$.

Definition 7.26. Suppose that families $\{K_i^{(1)}, \varphi_i^{(1)}\}$ and $\{K_i^{(2)}, \varphi_i^{(2)}\}$ have all the accumulation points of class C^0 . Say that these families *describe the same curve* if the set of finite points of the corresponding curves are equivalent as complex manifolds, and the equivalence continues by continuity to the points at infinity. Let the *moduli space* be the set of equivalence classes of such families up to relationship that they describe the same curve.

Note that it is not reasonable to drop the consideration of points at infinity.

Example 7.27. Indeed, consider a curve $\bar{M}$ with the Serpinsky carpet as the set M_∞ of accumulation points of disks. The Serpinsky carpet breaks $\mathbb{CP}^1$ into a union of triangles. Suppose that each triangle contains exactly one disk K_i . Then the smooth points on M form a disjoint union of tubes, so the only invariant is the conformal lengths of these tubes. This is one parameter per handle, much smaller than three parameters per handle as one would expect to have from finite-genus theory. The remaining parameters must be contained in the data for gluing the boundary of each smooth component to the set of points at infinity.

8. SET OF ADMISSIBLE BUNDLES

Here we investigate the structure of the set $\mathfrak{L}$ of admissible bundles $\mathcal{L}$ over the given curve (described by the model space $\bar{M}$, as in Section 6.4). The mapping $\mathcal{L} \mapsto$

$(\Pi_{\{\psi\}} \circ \mathbf{K}, \Pi_{\{\psi^{-1}\}} \circ \mathbf{K})$ into a pair of compact operators allows one to consider the topology on $\mathfrak{L}$ induced by the operator topology on the space of compact operators.

Thus $\mathfrak{L}$ is a topological space. We will describe some remarkable subsets of $\mathfrak{L}$ and algebraic structures on these subsets.

8.1. Exceptional indices. Since operators Π_{φ_i, ψ_i} are bounded, and row-blocks and column-blocks of the operator $\mathbf{K}$ are compact, we can conclude that instead of compactness of $\Pi\mathbf{K}$ one can require compactness of restriction of $\Pi\mathbf{K}$ to the direct sum of all but a finite number of $L_2(\partial K_i, \Omega^1 \otimes \mu)$. We call the indices of excluded contours *exceptional indices*.

In particular, we can include all the non-circular contours and all contours ∂K_i such that $\text{ind } \psi_i \neq 0$ in the set of exceptional indices. Thus whenever we discuss admissibility conditions we can suppose that $\text{ind } \psi_i = 0$ and all the K_i are disks.

The following facts are obvious:

Proposition 8.1. *If we change a finite number of functions ψ_i , this does not change the admissibility of the resulting bundle. If we multiply functions ψ_i by constants c_i with $|c_i|, |c_i^{-1}|$ being bounded, this does not change the admissibility of the bundle.*

8.2. Hilbert–Schmidt bundles. In practice the condition of admissibility is very hard to use directly, since there is no practically useful criterion of compactness which is necessary and sufficient. The closest simple-to-check approximation is the Hilbert–Schmidt condition.

Definition 8.2. We say that a curve with a bundle $(\bar{M}, \mathcal{L})$ is *Hilbert–Schmidt* if both operators $\Pi_{\{\psi\}}\mathbf{K}$ and $\Pi_{\{\psi^{-1}\}}\mathbf{K}$ are Hilbert–Schmidt operators.

Remark 8.3. Note that since the Hilbert structure on the sections of the bundle of half-forms on a curve is canonically defined, so is the notion of Hilbert–Schmidt operator.

8.3. Involution. Consider an admissible bundle $\mathcal{L}$ and the dual bundle $\mathcal{L}^{-1}$ (defined by inverse gluing conditions). The following statement is a direct corollary of definitions:

Proposition 8.4. *If $\mathcal{L}$ is admissible, so is $\mathcal{L}^{-1}$. If $\mathcal{L}$ is Hilbert–Schmidt, so is $\mathcal{L}^{-1}$.*

8.4. Hilbert–Schmidt criterion. From the description of a solution of the Riemann problem one can easily get the following criterion:

Theorem 8.5. *Consider a family of disks K_i , $i \in I$, with an involution $'$ on I and gluing data φ_i, ψ_i .*

1. *If*

$$\sum_{i \neq j} \left(|\psi_i|_{\mathbf{R}}^2 + |\psi_i^{-1}|_{\mathbf{R}}^2 \right) e^{-l_{ij}} < \infty$$

then $\mathcal{L}$ is Hilbert–Schmidt (here $\|\cdot\|_{\mathbf{R}}$ is the Riemann norm, see Definition 7.7).

2. Suppose that all but a finite number of functions ψ_i are constant. Then the corresponding curve M with a bundle $\mathcal{L}$ is Hilbert–Schmidt iff

$$\sum_{i \neq j} (|\psi_i|^2 + |\psi_i|^{-2}) e^{-l_{ij}} < \infty$$

(here the indices i for which ψ_i is not constant are excluded from summation).

Here l_{ij} is the conformal distance between ∂K_i and ∂K_j .

This theorem follows immediately from the following

Lemma 8.6. Let k_{ij} , $i, j \in I$ be the Hilbert–Schmidt norm of the block of $\mathbf{K}$ which maps $L_2^- (\partial K_i, \omega^{1/2} \otimes \mu) \rightarrow L_2^+ (\partial K_j, \omega^{1/2} \otimes \mu)$. Then

$$k_{ij} = \sum_{\substack{s > 0 \\ s \in \mathbb{Z} + 1/2}} e^{-sl_{ij}} = O(e^{-l_{ij}/2}).$$

Proof. It is enough to prove that the characteristic numbers of the block of $\mathbf{K}$ which is the mapping

$$L_2^- (\partial K_i, \omega^{1/2} \otimes \mu) \rightarrow L_2^+ (\partial K_j, \omega^{1/2} \otimes \mu)$$

are $e^{-sl_{ij}}$ for $s > 0$, $s \in \mathbb{Z} + \frac{1}{2}$. Since $\mathbf{K}$ is invariant with respect to fraction-linear transformations, we can assume that ∂K_i and ∂K_j bound a tube $S^1 \times (0, l_{ij})$ of circumference 2π and length l_{ij} . Then characteristic vectors of $\mathbf{K}$ correspond to holomorphic $1/2$ -forms

$$e^{isx} e^{-sy} dz^{1/2}, \quad z = x + iy, \quad (x, y) \in S^1 \times (0, l_{ij}),$$

the restrictions on $y = 0$ being the characteristic vectors themselves, the restrictions on $y = l_{ij}$ being their images, which are $e^{-sl_{ij}}$ times smaller. Now the condition that the restriction on $y = 0$ is a section of $\Omega \otimes \mu$ gives the condition that $s \in \mathbb{Z} + \frac{1}{2}$, the condition that this restriction is in $--$ -part of L_2 gives $s > 0$. $\square$

Corollary 8.7. 1. A curve $\bar{M}$ is Hilbert–Schmidt iff

$$\sum_{i \neq j} e^{-l_{ij}} < \infty;$$

2. If a curve $\bar{M}$ allows a Hilbert–Schmidt bundle, it is Hilbert–Schmidt itself.

Proof. Indeed, only the second part requires proof, and it follows from

$$\sum_{ij} k_{ij}^2 \leq \frac{1}{2} \sum_{ij} (|\psi_i|^2 + |\psi_i|^{-2}) k_{ij}^2.$$

$\square$

8.5. **log-convexity.** We were not able to prove the following

Conjecture 8.8. Let $\mathcal{L}$ and $\mathcal{M}$ are two bundles on $\bar{M}$ given by gluing conditions. Suppose that both $\mathcal{M} \otimes \mathcal{L}$ and $\mathcal{M} \otimes \mathcal{L}^{-1}$ are admissible. Then $\mathcal{M}$ is admissible as well.

However, the following statement is true:

Proposition 8.9. *Let $\mathcal{L}$ and $\mathcal{M}$ are two bundles on $\bar{M}$ given by gluing conditions with functions ψ_i and γ_i , and all but a finite number of these functions are constant. Suppose that both $\mathcal{M} \otimes \mathcal{L}$ and $\mathcal{M} \otimes \mathcal{L}^{-1}$ are Hilbert–Schmidt. Then $\mathcal{M}$ is Hilbert–Schmidt as well.*

Proof. We may suppose that all ψ_i and γ_i are constant. Then in notations of Lemma 8.6 we know that

$$\sum_{ij} (|\psi_i \gamma_i|^2 + |\psi_i \gamma_i|^{-2}) k_{ij}^2 < \infty,$$

$$\sum_{ij} (|\psi_i \gamma_i^{-1}|^2 + |\psi_i \gamma_i^{-1}|^{-2}) k_{ij}^2 < \infty,$$

and want to prove that

$$\sum_{ij} (|\psi_i|^2 + |\psi_i|^{-2}) k_{ij}^2 < \infty.$$

However, this is an obvious corollary of relation between geometric mean and arithmetic mean. $\square$

8.6. **Types of admissible bundles.** In what follows we are going to study Hilbert–Schmidt bundles, thus we may assume that the curve is Hilbert–Schmidt itself.

Definition 8.10. 1. Call the bundle $\omega \otimes \mathcal{L}$ on a curve $\bar{M}$ *real* if all the gluing functions ψ_i for this bundle are constants of magnitude 1.
 2. Call the bundle $\omega \otimes \mathcal{L}$ on a curve $\bar{M}$ *bounded* if all the gluing functions ψ_i for this bundle taken together are bounded in $\|\cdot\|_{\mathbf{R}}$ -norm.
 3. Call the bundle $\omega \otimes \mathcal{L}$ on a curve $\bar{M}$ *strongly Hilbert–Schmidt* if the bundle $\omega \otimes \mathcal{L}^n$ is Hilbert–Schmidt for any $n \in \mathbb{Z}$.

Lemma 8.11. *If the bundles $\omega \otimes \mathcal{L}$ and $\omega \otimes \mathcal{M}$ on a curve $\bar{M}$ are strongly Hilbert–Schmidt, then $\omega \otimes \mathcal{L} \otimes \mathcal{M}$ is also strongly Hilbert–Schmidt.*

Proof. Since $\omega \otimes \mathcal{L}^{2n}$ and $\omega \otimes \mathcal{M}^{2n}$ are Hilbert–Schmidt, such is $\omega \otimes \mathcal{L}^n \otimes \mathcal{M}^n$ by the log-convexity. $\square$

Lemma 8.12. *Any real bundle is bounded. Any bounded bundle is strongly Hilbert–Schmidt.*

Proposition 8.13. *Consider an admissible curve $\bar{M}$ and a bundle $\mathcal{L}$ defined by gluing conditions ψ_i . Suppose that for any $N > 0$*

$$\sum_{i \neq j} \left(\max_{R_i} |\psi_i|^N + \max_{R_i} |\psi_i|^{-N} \right) e^{-l_{ij}} < \infty.$$

Then $\mathcal{L}$ is strongly Hilbert-Schmidt.

Proof. Indeed, this is a direct corollary of Lemma 7.8. $\square$

Topology on $\mathfrak{L}$ defines a topology on the set of bounded bundles. Note that the latter topology is very easy to describe. Fix a number $M > 1$, and consider the subset of bundles with $|\psi_i|_{\mathbf{R}} < M$, $i \in I$. Then the topology on this subset is the topology of direct product. This topology is important for the description of Jacobian in Section 9.7.

8.7. Multipliers, equivalence and Jacobians. Consider what can play a rôle of a mapping $\mathfrak{m}: \mathcal{L}_1 \rightarrow \mathcal{L}_2$ between two bundles on $\bar{M}$ defined by gluing conditions. Inside the “smooth” part of the curve such mapping should be a multiplication by a section a of a holomorphic bundle, thus $\mathfrak{m}$ is determined by a function a which is holomorphic inside $\mathbb{CP}^1 \setminus \bigcup \overline{K_i}$.

We start with discussing heuristics for the properties of the function a . On one hand, $\mathfrak{m}$ should send holomorphic sections to holomorphic one. If $\bar{\partial}a$ is defined in a neighborhood of the infinity M_∞ , and is not 0, then by Leibniz rule $\mathfrak{m}$ will not send holomorphic sections of $\mathcal{L}_1$ to holomorphic section of $\mathcal{L}_2$.

On the other hand, $\mathfrak{m}$ should send $H^{1/2}$ -sections to $H^{1/2}$ -sections. Since $1 \in H^{1/2}$, $a \in H^{1/2}$. Moreover, if $\mathfrak{m}$ is bounded, multiplication by a should send L_2 -sections of $\Omega^{1/2}(\bigcup_i \partial K_i) \otimes \mu$ to L_2 -sections. Since the only multipliers in L_2 are essentially bounded functions, we conclude that the restriction that a is bounded on the domain of definition and is in $H^{1/2}$ looks like a particularly good candidate.

Proposition 8.14. *Consider a function a which is holomorphic inside $\mathbb{CP}^1 \setminus \bigcup \overline{K_i}$. Identify $\mathcal{H}(\mathbb{CP}^1, \{K_i\})$ with a subspace of the space of analytic functions on $\mathbb{CP}^1 \setminus \bigcup \overline{K_i}$. Suppose that the disks K_i are well separated. Multiplication by a preserves $\mathcal{H}(\mathbb{CP}^1, \{K_i\})$ if and only if a is a restriction of an element $\tilde{a} \in H^{1/2}(\mathbb{CP}^1 \setminus \bigcup K_i)$ such that $\bar{\partial}\tilde{a} = 0 \in H^{-1/2}(\mathbb{CP}^1 \setminus \bigcup K_i)$ and the restriction $\tilde{r}(\tilde{a}) \in L_2(\bigcup \partial K_i)$ of $\tilde{a}$ to the boundary is essentially bounded.*

Proof. Indeed, we may suppose that $\infty \in K_i$ for some $i \in I$. Let us show the “only if” part first. Considering $a \cdot dz^{1/2}$ we see that if a is a multiplier in $\mathcal{H}(\mathbb{CP}^1, \{K_i\})$, then $\tilde{a}$ satisfying the first two conditions of the proposition exists. Since the operator of multiplication by a is automatically bounded in $\mathcal{H}(\mathbb{CP}^1, \{K_i\})$, we see that the boundary value of $\tilde{a}$ is essentially bounded by the norm M of this operator.

Indeed, if $\varepsilon > 0$ and the essential supremum of $\tilde{r}(\tilde{a})$ is bigger than $M/(1 - \varepsilon)$, then there is an arc in $\bigcup \partial K_i$ such that $|\tilde{r}(\tilde{a})| > M/(1 - \varepsilon)$ on a subset of this arc

of relative measure greater than $1 - \varepsilon$. Taking $\|a \cdot \frac{dz^{1/2}}{z - z_0}\|_{L_2(\bigcup \partial K_i)}$ with z_0 close to this arc and inside $\bigcup K_i$, we obtain a contradiction.

The proof of the “if” part consists of three parts. First, let us show that if $\alpha \in \mathcal{H}(\mathbb{CP}^1, \{K_i\})$, then $a\alpha$ is a generalized half-form correctly defined up to addition of a half-form with support in $\bigcup \partial K_i$ (here we say that the support of a generalized function β is in a set U —not necessarily closed—if it is a weak limit of generalized functions β_n such that $\text{Supp } \beta_n \subset U$). Indeed, for any manifold M the formula $\langle \alpha\beta, \varphi \rangle \stackrel{\text{def}}{=} \langle \alpha, \varphi\beta \rangle$, $\varphi \in \mathcal{D}(M)$, $\alpha \in H^s(M)$, $\beta \in H^{-s}(M, \Omega^{\text{top}})$, shows that there is a natural pairing $(\alpha, \beta) \mapsto \alpha\beta$ of $H^s(M)$ with $H^{-s}(M, \Omega^{\text{top}})$ with values in $\mathcal{D}'(M)$.

This pairing is weakly bicontinuous, moreover, for any smooth vector field $v \in \text{Vect}(M)$ the Leibniz identity

$$v(\alpha\beta) = (v\alpha)\beta + \alpha(v\beta)$$

holds if $\alpha \in H^s(M)$, $\beta \in H^{1-s}(M, \Omega^{\text{top}})$. Since both a and α are of smoothness $H^{1/2}$, we see that $a\alpha$ is indeed a generalized function defined with the described above ambiguity (take $s = 1/2$, and make appropriate changes to adjust the above discussion to half-forms). Moreover, $\bar{\partial}(a\alpha) = (\bar{\partial}a)\alpha + a(\bar{\partial}\alpha)$, thus $\bar{\partial}(a\alpha) = 0$ as a generalized function defined up to addition of a function with support in $\bigcup \partial K_i$. What remains to prove is that $\bar{\partial}(a\alpha) \in H^{-1/2}(\mathbb{CP}^1, \omega^{1/2} \otimes \bar{\omega})$ after an appropriate choice of continuation of $a\alpha$ to $\mathbb{CP}^1$.

As a second step, fix $i \in I$ and show that $a\alpha$ has an appropriate extension into K_i . Consider restriction of $a\alpha$ to a small collar outside ∂K_i .

Lemma 8.15. *Consider two concentric circles $K \subset \tilde{K} \subset \mathbb{C}$, and a holomorphic half-form β in $\tilde{K} \setminus K$. Consider a L_2 -half-form B on ∂K , and suppose that Laurent coefficients of β coincide with Fourier coefficients of B . Then β has an $H^{1/2}$ -continuation into K with the $H^{1/2}$ -norm being $O(\|B\|_{L_2})$.*

Proof. Indeed, we know that positive Fourier coefficients of B are $O(e^{-\varepsilon k})$, and negative are in l_2 . Consider positive and negative parts of β separately. The positive part automatically continues into K , and the bound is a corollary of results of Section 2.7.

Consider now the negative part. We can suppose that $K = \{z \mid |z| < 1\}$, and β is $\sum B_k z^{-k} dz^{1/2}$, $(B_k) \in l_2$. Extend β into K as $\sum B_k \bar{z}^k dz^{1/2}$. Let us show that this extension satisfies the conditions of the lemma.

It is sufficient to show that the $H^{-1/2}$ -norm of $\bar{\partial}\beta = \sum k B_k \bar{z}^{k-1} \vartheta_{K_1} dz^{1/2} d\bar{z}$ is bounded (here ϑ_{K_1} is the characteristic function of K_1). Since different components of this form are perpendicular in $H^{-1/2}$, it is sufficient to bound the norm of one component. To do this it is sufficient to apply the methods of Section 2.7. $\square$

Application of this lemma to $a\alpha$ shows that $a\alpha$ may be extended into every disk K_i , moreover, that after these extensions the norms $\|\bar{\partial}(a\alpha)|_{K_i}\|_{H^{-1/2}}$ form a sequence in l_2 . Indeed, $\tilde{r}_i(a)$ is a bounded function on ∂K_i , thus $\tilde{r}_i(a)\tilde{r}_i(\alpha)$ has its L_2 -norm

bounded by L_2 -norm of $\tilde{r}_i(\alpha)$, and the latter norms (for different i) form a sequence in l_2 .

Third, we need to show that the above extensions can be glued together to an extension to $\mathbb{CP}^1$. Consider an arbitrary generalized-function-extension of $a\alpha$ to $\mathbb{CP}^1$ and a generalized function $\bar{\partial}(a\alpha)$. It is a generalized function with support in the disjoint union $\bigcup K_i$, thus the components $\bar{\partial}(a\alpha)|_{K_i}$ are well defined generalized forms b_i , $\text{Supp } b_i \subset K_i$. On the other hand, above we constructed an $H^{1/2}$ -extension of $a\alpha$ to K_i , let $\mathring{b}_i$ be $\bar{\partial}(a\alpha)$ obtained from this extension.

Lemma 8.16. $b_i - \mathring{b}_i = \bar{\partial}\beta_i$, β_i being a generalized function with support in K_i .

This lemma is an obvious corollary of the fact that restriction of $\bar{\partial}^{-1}(b_i - \mathring{b}_i)$ to a collar around ∂K_i can be holomorphically extended inside of K_i . Since both series $\sum b_i$ and $\sum \mathring{b}_i$ converge (one in $\mathcal{D}'$, another in $H^{-1/2}$), we conclude that $\sum (b_i - \mathring{b}_i)$ converges in $\mathcal{D}'$, thus $\sum \beta_i$ converges in $\mathcal{D}'$.

We conclude that $\beta = a\alpha - \sum \beta_i$ is a generalized function such that

1. $\bar{\partial}\beta$ has support in $\bigcup K_i$;
2. β coincides with $a\alpha$ inside $\mathbb{CP}^1 \setminus \overline{\bigcup K_i}$;
3. $\bar{\partial}\beta$ is in $H^{-1/2}$.

Last condition implies $\beta \in H^{1/2}$, which shows that $a\alpha$ can be extended to an element of $H^{1/2}(\mathbb{CP}^1, \omega^{1/2})$, thus $a\alpha \in \mathcal{H}(\mathbb{CP}^1, \{K_i\})$. This finishes the proof of Proposition 8.14. $\square$

Definition 8.17. Call a bounded operator $\mathcal{M}: \mathcal{H}(\mathbb{CP}^1, \{K_i\}) \rightarrow \mathcal{H}(\mathbb{CP}^1, \{K_i\})$ a *multiplicator*, if for some point $z_0 \in \mathbb{CP}^1 \setminus \overline{\bigcup K_i}$ there is a formal power series ν at z_0 such that for any $f \in \mathcal{H}(\mathbb{CP}^1, \{K_i\})$ formal power series $\mathcal{M}f$ and νf coincide.

(This is just a formal way to say that $\mathcal{M}$ is a bounded operator of multiplication by a holomorphic function.)

Definition 8.18. Let $\mathcal{H}^\infty(\mathbb{CP}^1, \{K_i\}) \subset H^{1/2}(\mathbb{CP}^1 \setminus \bigcup K_i)$ consists of functions f such that the restriction $\tilde{r}(f)$ to $\bigcup \partial K_i$ (which is automatically in $L_2(\bigcup \partial K_i)$) is essentially bounded. Define a norm on this space by taking the essential maximum of $|\tilde{r}(f)|$.

Amplification 8.19. *Multiplicators form an algebra. The set of multiplicators coincides with $\mathcal{H}^\infty(\mathbb{CP}^1, \{K_i\})$. One can choose a norm in $\mathcal{H}(\mathbb{CP}^1, \{K_i\})$ in such a way that the operator norm of any multiplicator coincides with the $\mathcal{H}^\infty$ -norm.*

Proof. The only statement which needs a proof is the last one, and the norm in question is the norm induced by the inclusion into $\bigoplus_{l_2} L_2(\partial K_i, \Omega^{1/2} \otimes \mu)$. $\square$

Corollary 8.20. *Let $z_0 \in \mathbb{CP}^1 \setminus \bigcup \overline{K_i}$, $f \in \mathcal{H}^\infty(\mathbb{CP}^1, \{K_i\})$. Then $|f(z_0)| \leq \|f\|_{\mathcal{H}^\infty}$.*

Proof. One can suppose that $\infty \in K_i$ for some $i \in I$, so that $dz^{1/2} \in \mathcal{H}$. Since L_2 -norm on $\mathbb{CP}^1 \setminus \bigcup \overline{K_i}$ is majorated by $H^{1/2}$ -norm, we see that $\|f^n dz^{1/2}\|_{L_2} \leq C \cdot \|f\|_{\mathcal{H}^\infty}^n$, $n \in \mathbb{N}$. Hence $|f(z_0)|^n \leq C \cdot \|f\|_{\mathcal{H}^\infty}^n$, thus $|f(z_0)| \leq \|f\|_{\mathcal{H}^\infty}$. $\square$

Last conditions on $\mathbf{m}$ is that it should preserve the gluing conditions. In particular, if $\mathcal{L}_1$ is defined by gluing conditions ψ_i , and $\mathcal{L}_2$ by ξ_i , then

$$(8.1) \quad \xi_i = \frac{a \circ \varphi_i}{a} \psi_i.$$

Definition 8.21. Say that linear bundles $\mathcal{L}_1, \mathcal{L}_2$ defined by gluing conditions $\{\psi_i\}$ and $\{\xi_i\}$ are *bounded-equivalent* if there exists a function $a \in \mathcal{H}^\infty(\mathbb{CP}^1, \{K_i\})$ such that $\xi_i = \frac{a \circ \varphi_i}{a} \psi_i$ and $a^{-1} \in \mathcal{H}^\infty(\mathbb{CP}^1, \{K_i\})$.

Remark 8.22. It is obvious that a bundle which is bounded-equivalent to a bounded bundle is bounded itself. Moreover, if a bundle is bounded-equivalent to a (strongly) Hilbert–Schmidt bundle, it is (strongly) Hilbert–Schmidt itself.

Definition 8.23. The *Jacobian* is the set of equivalence classes of admissible bundles of degree 0. The *bounded Jacobian* is the subset of Jacobian which consists of classes of bounded bundles, similarly for *(strongly) Hilbert–Schmidt Jacobian*, and *real Jacobian*. *Constant Jacobian* is formed from classes of bundles defined by constant gluing functions, similarly one can define different flavors of constant Jacobians.

Multiplication by an appropriate rational function with zeros and poles inside $\bigcup \overset{\circ}{K}_i$ shows that

Proposition 8.24. *Any admissible bundle of degree 0 is bounded-equivalent to a bundle with all the gluing functions ψ_i of index 0.*

8.8. Divisors. Consider a model $(\mathbb{CP}^1, \{K_\bullet\}, \{\varphi_\bullet\})$ of a curve $\bar{M}$, and a rational function a on $\mathbb{CP}^1$ with a divisor D . Suppose that the part of D inside $\bigcup R_i$ is invariant w.r.t. $\varphi_\bullet$, and D does not intersect with the infinity $M_\infty \subset \mathbb{CP}^1$ of the curve M . If the bundle $\mathcal{L}$ with gluing data $\{\psi_i\}$ is admissible, so is the bundle $\mathcal{L}(D)$ given by the gluing data

$$\xi_i = \frac{a \circ \varphi_i}{a} \psi_i.$$

Obviously, $\deg \mathcal{L}(D) = \deg \mathcal{L} + \deg' D$, if $\deg' D$ is the degree of the part of D inside $M \setminus \bigcup R_i$ plus half the degree of the part of D inside $\bigcup R_i$. Moreover, if we change the part of D inside $\bigcup \overset{\circ}{K}_i$, the bundle $\mathcal{L}(D)$ will change to a bounded-equivalent one.

In particular, to any finite subset D_0 of M (with integer multiplicities fixed for any point of D_0) we associate a transformation $\mathcal{L} \mapsto \mathcal{L}(D_0)$ where the right-hand

side is defined up to equivalence. Note that if we fix a point $Z \in \bigcup \overset{\circ}{K}_i$, then we can complete any divisor D_0 on M to a divisor on $\mathbb{C}P^1$ of degree 0 by adding some multiple of Z , thus one can define $\mathcal{L}(D_0)$ uniquely. Moreover, if D_0 is of degree 0, then $\mathcal{L}(D_0)$ does not depend on the choice of Z .

Consider now two sequences of points $(x_k), (y_k) \subset \mathbb{C}P^1$. Let $a_k(z) = \frac{z-x_k}{z-y_k}$, and $a(z) = \prod_k a_k(z)$. If x_k is sufficiently close to y_k , and both these points are in appropriate neighborhood of $\overset{\circ}{K}_k$, then the infinite product converges and defines a bounded function on $\mathbb{C}P^1 \setminus \bigcup \overset{\circ}{K}_i$. We conclude that it is possible to consider also some “infinite” divisors (of finite degree) on M .

Consider now a finite divisor $D \subset \mathbb{C}P^1$ such that $D \cap M_\infty$ consists of one point $z_0 \in M_\infty$ (with some multiplicity). Changing D to $D - (\deg D) \cdot Z$, we can consider D as a divisor of a rational function a , thus $\mathcal{L}(D)$ is correctly defined by gluing conditions (8.1). For $\mathcal{L}(D)$ to be admissible for any such D and any admissible $\mathcal{L}$ it is sufficient that for some constant C

$$\text{dist}(z_0, \text{center}(K_{i'})) \leq C \cdot \text{dist}(z_0, \text{center}(K_i)), \quad i \in I$$

(since the disks K_i are well-separated). On the other hand, suppose that all the bundles with gluing functions

$$\Psi_i = \left(\frac{\text{dist}(z_0, \text{center}(K_{i'}))}{\text{dist}(z_0, \text{center}(K_i))} \right)^N, \quad n \in \mathbb{Z},$$

are Hilbert–Schmidt. Then by log-convexity the bundle $L(D)$ is strongly Hilbert–Schmidt if $\mathcal{L}$ is strongly Hilbert–Schmidt.

Definition 8.25. Say that the point $z_0 \in M_\infty$ is *bounded* if for any $N \in \mathbb{Z}$ the sheaf with constant gluing functions

$$\Psi_i = \left(\frac{\text{dist}(z_0, \text{center}(K_{i'}))}{\text{dist}(z_0, \text{center}(K_i))} \right)^N, \quad n \in \mathbb{Z}, i \in I,$$

is Hilbert–Schmidt.

We see that if the point $z_0 \in M_\infty$ is bounded, then the function $a_N(z) = \left(\frac{z-z_0}{z-Z} \right)^N$ defines a strongly Hilbert–Schmidt sheaf of degree 1. In particular, one can consider finite divisors on $\bar{M}$ which consist of points of M and bounded points on M_∞ . Similarly, one can also consider some infinite divisors on $\bar{M}$.

8.9. Universal Grassmannian and bundles.

Definition 8.26. Let V is the vector space of sequences (a_k) , $k \in \mathbb{Z}$, such that $a_k = 0$ for $k \ll 0$. The vector space V carries a natural topology of inductive limit of projective limits. Let

$$V_+ = \{(a_k) \mid a_k = 0 \text{ if } k < 0\}, \quad V_- = \{(a_k) \in V \mid a_k = 0 \text{ if } k \geq 0\}.$$

Say that the vector subspace $W \subset V$ is *admissible* if $W \cap V_-$ is finite-dimensional, and $W + V_-$ is closed and has a finite codimension in V . Let *universal Grassmannian* $\mathcal{G}$ be the set of admissible vector subspaces with natural topology.

Consider an admissible pair $(\bar{M}, \mathcal{L})$ and a smooth point $z_0 \in M$. Pick up a coordinate system z in neighborhood of z_0 , and a half-form f defined in the same neighborhood. Now any global section of $\mathcal{L}$ may be written as $g(z)f(z)$ with $g(z)$ being a holomorphic function which is correctly defined in the neighborhood of z_0 .

Call f a *local section* of $\mathcal{L}$ near z_0 . If $\mathcal{L}'$ is an equivalent to $\mathcal{L}$ bundle, and the equivalence is given by multiplication by a , we obtain a local section af of $\mathcal{L}'$. Say that local sections f and af are *equivalent*.

Consider now a bundle $\mathcal{L}(k \cdot z_0)$, defined using a point $Z \in \mathring{K}_i$ (as in the previous section). A section $h(\zeta)$, $\zeta \in \mathbb{C}P^1$, of this bundle can be identified (via multiplication by $\left(\frac{\zeta - Z}{\zeta - z_0}\right)^k$) with a “meromorphic section” of $\mathcal{L}$, i.e., one can write it as $g(z)f(z)$, $g(z)$ being a meromorphic function correctly defined in the neighborhood of z_0 . Moreover, $g(z)$ has poles only at z_0 . One can momentarily see that $g(z)$ does not depend on the choice of the point Z , more precise, for a different choice of Z there is a different choice of h which gives the same $g(z)$.

Associate to any such function $g(z)$ the sequence of its Laurent coefficients. Consider this sequence as an element of V . Let W be the vector subspace of V spanned by all possible functions $g(z)$ for bundles $\mathcal{L}(k \cdot z_0)$, $k \in \mathbb{Z}$. The Riemann–Roch theorem momentarily implies that W is admissible. Indeed, the condition that $W \cap V_-$ is finite-dimensional means that $\mathcal{L}$ has a finite-dimensional space of global sections. The condition on $W + V_-$ is implied by the following fact:

Lemma 8.27. *Consider a admissible semibounded bundle $\mathcal{L}$ on a curve $\bar{M}$, and a point $z_0 \in M$. Then*

1. *for big enough k the bundle $\mathcal{L}(-k \cdot z_0)$ has no sections.*
2. *for big enough k the bundle $\mathcal{L}((k+1) \cdot z_0)$ has one more section than $\mathcal{L}(k \cdot z_0)$.*

Proof. The second statement is a corollary of the first one, of Riemann–Roch theorem and duality. The first one is obvious, since the global sections of $\mathcal{L}(-k \cdot z_0)$ are naturally identified with global sections of $\mathcal{L}$ which have a zero of k -th order at z_0 . $\square$

We obtained

Proposition 8.28. *To each admissible pair $(\bar{M}, \mathcal{L})$ with a fixed smooth point $z_0 \in M$, a coordinate system near z_0 , and a local section f of $\mathcal{L}$ one can associate a point $W \in \mathcal{G}$. If we change $\mathcal{L}$ and f to an equivalent bundle with a local section, W does not change.*

One can generalize this proposition to some points at infinity. If $z_0 \in M_\infty$, define a *local section* of $\mathcal{L}$ near z_0 in the same way as above, i.e., as a non-zero section of $\omega^{1/2}$ near z_0 . In what follows we use only the ∞ -jet of this section.

Amplification 8.29. *Consider an admissible pair $(\bar{M}, \mathcal{L})$ with a fixed point at infinity $z_0 \in M_\infty$, a coordinate system near z_0 , and a local section f of $\mathcal{L}$. Suppose that $\mathcal{L}$ is Hilbert–Schmidt, and z_0 is bounded and of class C^∞ . To this data we can associate a point $W \in \mathcal{G}$. If we change $\mathcal{L}$ and f to an equivalent bundle with a local section, W does not change.*

Proof. Since z_0 is bounded, and $\mathcal{L}$ is Hilbert–Schmidt, $\mathcal{L}(k \cdot z_0)$ is Hilbert–Schmidt too, thus the Riemann–Roch theorem is applicable. Since z_0 is of class C^∞ , any section of $\mathcal{L}(k \cdot z_0)$ has a (formal) Taylor series near z_0 , thus the corresponding section of $\mathcal{L}$ has a (formal) Laurent series.

What remains to be proved is the fact that equivalence between bundles can be pushed to infinity points of class C^∞ . Note that one can associate an element of the generalized Hardy space to any bounded function a . Indeed, suppose that $\zeta = \infty \in K_i$, then $a \cdot d\zeta^{1/2}$ is an element of $\mathcal{H}$. In particular, a has an asymptotic expansion near z_0 , thus multiplication by a maps (formal) Laurent series at z_0 to themselves. $\square$

9. STRUCTURE OF JACOBIAN

9.1. Constant Jacobian. Consider the involution $': I \rightarrow I$. It defines a transposition matrix $t = (t_{ij})$, $i, j \in I$, $t_{ij} = \delta_{ij'}$. We use the results of toy theory (see Section 4) to obtain the following

Theorem 9.1. *Let l_{ij} be pairwise conformal distances between disks K_i . Suppose that the matrix $\mathcal{R} = (e^{-l_{ij}} - \delta_{ij})$ defines a compact mapping $l_2 \rightarrow l_2$ and a bounded mapping $l_\infty \rightarrow l_\infty$, and that for some $N > 0$ the matrix $(t\mathcal{R})^N$ defines a mapping $l_\infty \rightarrow l_2$. Then for any admissible curve obtained by gluing circles ∂K_i the constant bounded Jacobian coincides with the bounded Jacobian.*

Proof. Indeed, to show this we need to show that for any cycle $\{\psi_i\}_{i \in I}$ with $\deg \psi_i = 0$ and $|\psi_i| < C$ one can find a bounded collection of constants c_i such that $c_i \psi_i = \frac{a \circ \varphi_i}{a}$ for some a such that $a, a^{-1} \in \mathcal{H}^\infty(\mathbb{CP}^1, \{K_i\})$.

Taking logarithms, we see that it is sufficient to show that the mapping

$$\mathcal{J}_b: f + C \mapsto \left(f|_{\partial K_j} - \varphi^* \left(f|_{\partial K_{j'}} \right) + C_j \right)_{j \in I_+}$$

from bounded analytic functions (modulo constants) to functions on boundary (modulo constants) with bounded $\pm$ -parts is surjective. In Section 4.5 we have seen that (given the first condition of the theorem) a similar mapping

$$\mathcal{J}: \mathcal{H}^{(1)} / \text{const} \rightarrow \bigoplus_{\substack{l_2 \\ j \in I_+}} H^{1/2}(\partial K_j) / \text{const}$$

is a bounded mapping of index 0. We are going to prove the surjectivity by using a combination of following lemmas:

Lemma 9.2. *If $(e^{-l_{ij}} - \delta_{ij})$ gives a compact operator $l_2 \rightarrow l_2$, then the mapping $\mathcal{J}$ is a bijection.*

Proof. Since we know the index of $\mathcal{J}$, it is sufficient to show that $\text{Ker } \mathcal{J} = 0$. Let $f \in \mathcal{H}^{(1)}/\text{const}$ and $\mathcal{J}f = 0$. We are going to show that $\|f\|_{H^1} = 0$. Since ∂ is elliptic, and $\bar{\partial}f = 0$, it is sufficient to show that $\|\partial f\|_{L_2} = 0$, i.e., that $\int \partial f \wedge \bar{\partial} \bar{f} = 0$, the integral is taken over $\mathbb{C}P^1 \setminus \bigcup K_i$. Denote the domain of integration by S . Note that the above integral is well defined, since $\partial f \in H^0(S) = L_2(S)$.

First of all, since $\wedge$ -product of any two sections of ω is 0, one can add $\partial \bar{f}$ to $\bar{\partial} \bar{f}$ without changing the integral, thus it is sufficient to show that $\int_S \partial f \wedge d\bar{f} = 0$. Additionally, since $\bar{\partial}f = 0$ on S , one can change ∂f to df without changing the integral, thus it is sufficient to show that $\int_S df \wedge d\bar{f} = 0$. Second, use the duality identity $\int_S d\alpha = \int_{\partial S} \alpha$. Applying it (at first formally), we get

$$\begin{aligned} \int_S df \wedge d\bar{f} &= \int_S d(f d\bar{f}) = \int_{\partial S} f d\bar{f} = - \sum_{i \in I} \int_{\partial K_i} f d\bar{f} \\ &= - \sum_{i \in I_+} \left(\int_{\partial K_i} f d\bar{f} + \int_{\partial K_{i'}} f d\bar{f} \right). \end{aligned}$$

We want to show that $\int_{\partial K_i} f d\bar{f} + \int_{\partial K_{i'}} f d\bar{f} = 0$. Indeed, $\int_{\partial K_{i'}} f d\bar{f} = - \int_{\partial K_i} \varphi_{i'}^* (f d\bar{f})$, the sign appears since φ_i changes the orientations of $\partial K_\bullet$. Now the above equality becomes obvious, since $f \circ \varphi_i = f + C_i$ when both sides are defined, thus the sum of the integrals is $-C_i \int_{\partial K_i} d\bar{f} = 0$.

What remains to be proved is that one can indeed apply the formula $\int_S d\alpha = \int_{\partial S} \alpha$ in our situation, when α is not smooth, and S has non-smooth boundary. Consider an H^1 -extension g of f to $\mathbb{C}P^1$. Since $dg \wedge d\bar{g} \in L_1(\Omega^{\text{top}})$, the integral $\int_S df \wedge d\bar{f}$ can be represented as

$$\int_{\mathbb{C}P^1} dg \wedge d\bar{g} - \int_{\bigcup K_i} dg \wedge d\bar{g} = \int_{\mathbb{C}P^1} dg \wedge d\bar{g} - \sum \int_{K_i} dg \wedge d\bar{g}.$$

It remains to prove that $\int_{\mathbb{C}P^1} dg \wedge d\bar{g} = 0$, and $\int_{K_i} dg \wedge d\bar{g} = \int_{\partial K_i} g d\bar{g}$. Note that in both identities the boundary is already smooth (it is empty in the first one!).

Let us prove that if M is a two-dimension manifold, S is a compact subset of M with a smooth boundary, and $g_1, g_2 \in H^1(M)$, then

$$\int_S dg_1 \wedge dg_2 = \int_{\partial S} g_1 dg_2.$$

Here we understand the right-hand side as a natural pairing between $g_1|_{\partial S} \in H^{1/2}(\partial S)$ and $d(g_2|_{\partial S}) \in H^{-1/2}(\partial S, \Omega_{\partial S}^{\text{top}})$. However, both sides define bounded bilinear functionals on $H^1(M)$, thus it is sufficient to check them on a dense subset $C^\infty(M)$, where they are true due to de Rham theory. $\square$

To continue the proof of the theorem, note that the operator $\mathcal{J}$ is related to the operator $\mathbf{K}$ in the following way: identify $\mathcal{H}^{(1)}/\text{const}$ with $\bigoplus_{i \in I} H_-^{1/2}(\partial K_i)/\text{const}$.

Let Π_i be the identification of $H_+^{1/2}(\partial K_i)$ with $H_-^{1/2}(\partial K_{i'})$ via φ^* , let $\Pi^{(1)}$ be the direct sum of such identifications. As we have seen it in Section 4.5, the image of $\mathcal{J}$ coincides with the image of $\Pi^{(1)} \circ \mathbf{K} - 1$ (which is a mapping

$$\bigoplus_{l_2} H_-^{1/2}(\partial K_i)/\text{const} \rightarrow \bigoplus_{l_2} H_-^{1/2}(\partial K_i)/\text{const},$$

but we identify $\bigoplus_{l_2} H_-^{1/2}(\partial K_i)/\text{const}$ with $\bigoplus_{j \in I_+} H_-^{1/2}(\partial K_j)/\text{const}$ via $\varphi_\bullet^*$).

- Lemma 9.3.** 1. Consider two disjoint disks K_1, K_2 on $\mathbb{CP}^1$ of conformal distance l . The operator $\mathbf{K}$ with smooth kernel $\frac{dy}{y-x}$ defines bounded operators $H^{1/2}(\partial K_1) \rightarrow L_\infty(\partial K_2)$, $L_\infty(\partial K_1) \rightarrow L_\infty(\partial K_2)$, $H^{1/2}(\partial K_1) \rightarrow H^{1/2}(\partial K_2)$, $L_\infty(\partial K_1) \rightarrow H^{1/2}(\partial K_2)$ with the norms being $O(e^{-l})$.
2. Consider a family of disjoint disks $\{K_i\}$ on $\mathbb{CP}^1$ with conformal distances l_{ij} , and suppose that the matrix $\mathcal{R} = (e^{-l_{ij}} - \delta_{ij})$ defines bounded mappings $l_2 \rightarrow l_2$ and $l_\infty \rightarrow l_\infty$, and for some $N > 0$ the matrix $(t\mathcal{R})^N$ defines a mapping $l_\infty \rightarrow l_2$. Then the operator $\mathbf{K}$ gives bounded mappings $\bigoplus_{l_2} H^{1/2}(\partial K_i) \rightarrow \bigoplus_{l_2} H^{1/2}(\partial K_i)$ and $L_\infty(\bigcup \partial K_i) \rightarrow L_\infty(\bigcup \partial K_i)$, moreover, the operator $(\Pi^{(1)}\mathbf{K})^N$ gives a bounded mapping $L_\infty(\bigcup \partial K_i) \rightarrow \bigoplus_{l_2} H^{1/2}(\partial K_i)$.
3. In the conditions of the previous part of the lemma let $f_- \in L_\infty(\bigcup \partial K_i)$. Suppose that $+$ -parts of f_- on all the ∂K_i vanish. Then $\mathbf{K}f_-$ is a bounded analytic function.

Now we are able to invert the mapping $\mathcal{J}_b$. Consider a fixed function f on $\bigcup \partial K_i$ with bounded $+$ -part and bounded $-$ -part. If $f - f \circ \varphi_i$ is a constant for any $i \in I$, then $+$ -part of $f|_{\partial K_i}$ can be reconstructed basing on $-$ -part of $f|_{\partial K_{i'}}$, thus one can identify f (modulo constants) with the collection of $-$ -parts of f on $\bigcup_{i \in I} \partial K_i$ (modulo constants).

The restrictions on l_{ij} show that the radii of disks K_i for a sequence from l_1 . Hence the Cauchy kernel allows one to construct an analytic function $\tilde{f}$ on S such that the $-$ -parts of $\tilde{f}|_{\partial K_i}$ and $f|_{\partial K_i}$ coincide (modulo constants). Moreover, $\mathcal{J}_b \tilde{f} = (\Pi^{(1)} \circ \mathbf{K} - 1)(f)$. Since f is bounded, it is in L_2 , thus $\tilde{f} \in H^{1/2}$. Since $\tilde{f}|_{\bigcup \partial K_i}$ is bounded, $\tilde{f}$ is a multiplier.

We see that $\Pi^{(1)} \circ \mathbf{K}(f)$ is a well-defined bounded function on $\bigcup_{j \in I_+} \partial K_j$, moreover, $(\Pi^{(1)} \circ \mathbf{K})^N(f) \in \bigoplus_{l_2} H^{1/2}(\partial K_j)$. In particular, $(\Pi^{(1)} \circ \mathbf{K})^N(f) = \mathcal{J}F$ for some function $F \in \mathcal{H}^{(1)}$. To show that f is in the image of $\Pi^{(1)} \circ \mathbf{K} - 1$, it remains to prove is that $f - (\Pi^{(1)} \circ \mathbf{K})^N(f)$ is in the image of $\Pi^{(1)} \circ \mathbf{K} - 1$, what is obvious.

This finishes the proof of Theorem 9.1. $\square$

To demonstrate a particular case in which the assumptions of Theorem 9.1 hold, consider

Lemma 9.4. *Suppose that the matrix (a_{ij}) gives a Hilbert–Schmidt operator $l_2 \rightarrow l_2$, i.e., $\sum |a_{ij}|^2 < \infty$. Then the matrix (a_{ij}^2) gives a bounded operator $l_\infty \rightarrow l_2$.*

If $a_{ij} > 0$ for any $i, j \in I$, and the matrix (a_{ij}) gives a compact mapping $l_2 \rightarrow l_2$, then (a_{ij}^2) also gives a compact mapping $l_2 \rightarrow l_2$.

Corollary 9.5. *If $\bar{M}$ is a Hilbert–Schmidt curve, then the bounded Jacobian of $\bar{M}$ coincides with the constant bounded Jacobian.*

In Section 9.7 we show that under suitable restrictions on $\bar{M}$ the above Jacobians are in 1-to-1 correspondence with quotients of topological vector spaces by $\mathbb{Z}$ -lattices. On the other hand, this statement is obvious when applied to real Jacobian, since it is a direct product of circles $|\psi_i| = 1$. To construct an isomorphism of this product with a quotient by a lattice, one can take the Hilbert space l_2 with the standard basis e_i , and the lattice spanned by $\alpha_i e_i$, with $\alpha_i \rightarrow 0$.

The description of the real part of Jacobian was first obtained in [7] in the case of a real hyperelliptic curve of a special form.

9.2. The partial period mapping. Consider a theory similar to one discussed in Section 4.4, but related to holomorphic 1-forms instead of holomorphic functions. Consider a family $\{K_i\}$ of disjoint disks in $\mathbb{CP}^1$.

Definition 9.6. We say that a generalized-function section α of the linear bundle ω on $\mathbb{CP}^1$ is H^0 -holomorphic in $\mathbb{CP}^1 \setminus \bigcup K_i$ if $\alpha \in H^0(\mathbb{CP}^1 \setminus \bigcup K_i, \omega) = L_2(\mathbb{CP}^1 \setminus \bigcup K_i, \omega)$, and $\bar{\partial}\alpha = 0 \in H^{-1}(\mathbb{CP}^1 \setminus \bigcup K_i, \omega \otimes \bar{\omega})$. Denote the space of H^0 -holomorphic forms in $\mathbb{CP}^1 \setminus \bigcup K_i$ by $\mathcal{H}^{(0)}$.

Since one can extend a H^0 -holomorphic form α into any disk K_i by 0 without increasing H^0 -norm, α has a canonical extension to $\mathbb{CP}^1$, and $\bar{\partial}\alpha$ is concentrated on $\bigcup \partial K_i$. Moreover, if K_i has a neighborhood which does not intersect with other disks, one can define a *boundary value* $\alpha|_{\partial K_i}$ of α on ∂K_i , which is an element of $H^{-1/2}(\partial K_i, \Omega_{\partial K_i}^1)$ (note that if γ is a curve in $\mathbb{CP}^1$, then $\omega|_\gamma \simeq \Omega_\gamma^1$; one can get the $H^{-1/2}$ -restriction on the smoothness of the boundary value in the same way as in Section 2.7). Now $\bar{\partial}\alpha$ can be described as the extension of $-\alpha|_{\partial K_i}$ from ∂K_i to $\mathbb{CP}^1$ by δ -function (see (3.1)). (Such an extension is a correctly defined continuous mapping $H^{-1/2}(\partial K_i, \Omega_{\partial K_i}^1) \rightarrow H^{-1}(\mathbb{CP}^1, \omega \otimes \bar{\omega})$.)

Since the norm in $H^1(\mathbb{CP}^1)$ can be described by a local (non-invariant) formula $\|\alpha\|_{l_2}^2 + \|\alpha_{x,y}\|_{l_2}^2 + \|\alpha_{y,x}\|_{l_2}^2$ (here $x+iy$ is the local coordinate on $\mathbb{CP}^1$), functions with non-intersecting support are orthogonal in H^1 . Dually, if U_i are pairwise non-intersecting, then the natural mapping

$$H^{-1}(\mathbb{CP}^1, \omega \otimes \bar{\omega}) \rightarrow \bigoplus_{l_2} H^{-1}(U_i, \omega \otimes \bar{\omega})$$

is a continuous epimorphism.

Definition 9.7. Fix $\varepsilon > 0$. Say that disks $\{K_i\}$ have a *thickening* $\{U_i\}$ if K_i have non-intersecting neighborhoods U_i such that a pair $K_i \subset U_i$ is conformally equivalent to $\{|z| < 1\} \subset \{|z| < 1 + \varepsilon\}$. Say that $\{K_i\}$ have a *uniform thickening* if the neighborhood U_i can be picked up to be concentric to K_i circles (assume that a metric on $\mathbb{CP}^1$ is fixed).

(Note that the existence of uniform thickening does not depend on the metric on $\mathbb{CP}^1$.)

In particular, if $\{K_i\}$ have a thickening $\{U_i\}$, then the above arguments show that $(\|\alpha|_{\partial K_i}\|_{H^{-1/2}})_{i \in I} \in l_2$. Note also that $H^{-1/2}(\partial K_i, \Omega_{\partial K_i}^1)$ is dual to $H^{1/2}(\partial K_i)$, thus the subspace $H_{f=0}^{-1/2}(\partial K_i, \Omega_{\partial K_i}^1)$ of forms with vanishing integral has an $\text{PGL}(2, \mathbb{C})$ -invariant Hilbert structure (in the same sense as in Section 4.3).

Definition 9.8. Call the mapping $\alpha \mapsto \left(\int_{\partial K_i} \alpha\right)_{i \in I}$, $H^0(\mathbb{CP}^1 \setminus \bigcup K_i, \omega) \rightarrow l_2$ the *integration mapping*. Fix an involution $': I \rightarrow I$ which interchanges two parts of I , $I = I_+ \amalg I'_+$, and fraction-linear identifications $\varphi_i: \partial K_i \rightarrow \partial K_i$ (with the same conditions as in Section 4.5), then a *global holomorphic form* α is an element of $\mathcal{H}^{(0)}$ such that $\varphi_i^*(\alpha|_{\partial K_{i'}}) = \alpha|_{\partial K_i}$. Define the *partial period mapping* P by restriction of the integration mapping to global holomorphic forms, and taking integrals only for $i \in I_+$. Let $\bar{M}$ be the curve obtained by gluing ∂K_i together via φ_i . Denote by $\Gamma(\bar{M}, \omega)$ the space of global holomorphic forms on $\bar{M}$, i.e., the space of global holomorphic forms compatible with $\varphi_\bullet$.

The significant difference of this case and the case of Section 4.4 is that the mapping $\bar{\partial}: H^0(\mathbb{CP}^1, \omega) \rightarrow H^{-1}(\mathbb{CP}^1, \omega \otimes \bar{\omega})$ has no null-space, but has 1-dimensional cokernel, thus it is not easy to reconstruct α basing on $\bar{\partial}\alpha$ by local formulae. However, it is easy to prove

Lemma 9.9. *Suppose that $\{K_i\}$ has a thickening. Then the partial period mapping P satisfies the identity*

$$\dim \text{Ker } P = \dim \text{Ker } \mathcal{J}: \mathcal{H}^{(1)} / \text{const} \rightarrow \bigoplus_{l_2} H^{1/2}(\partial K_j) / \text{const}.$$

Proof. We claim that ∂ gives a mapping from one null-space to another one. Let $\alpha \in \text{Ker } P$. Then $\alpha|_{\partial K_i} \in H^{-1/2}(\partial K_i, \Omega_{\partial K_i}^1)$ has an antiderivative $f_i \in H^{1/2}(\partial K_i)$, and the harmonic extension of f_i into K_i has a bounded H^1 -norm. Since α is closed in $U_i \setminus K_i$, it has an antiderivative. Since ∂ is elliptic, the antiderivative has smoothness H^1 . It is possible to pick up the constant in such a way that two antiderivatives coincide on ∂K_i . Taking de Rham differential of resulting H^1 -function, we see that one can extend α into K_i preserving the closeness, and the related increase of the norm of α is bounded by $\|\alpha|_{\partial K_i}\|_{H^{-1/2}}$.

Repeating this operation for all the K_i , we obtain a closed extension of α to $\mathbb{CP}^1$ with the H^0 -norm bounded by $\|\alpha\|_{H^0}$ (here we use locality of L_2 -norm). Thus α has a H^1 -antiderivative f on $\mathbb{CP}^1 \setminus \bigcup K_i$, thus $\alpha = df$, hence $\alpha = \bar{\partial}f$. Obviously, $f \in \text{Ker } \mathcal{J}$.

On the other hand, if $f \in \text{Ker } \mathcal{J}$, then $\bar{\partial}f \in \text{Ker } P$. $\square$

Remark 9.10. As we have seen in Section 9.1, in assumptions of Theorem 4.11 the dimension in the lemma is 0, thus P defines an inclusion of global holomorphic forms into l_2 . Note that we do not claim that this inclusion is a monomorphism—estimates in Section 9.3 show that it is not if $\text{card}(I) = \infty$.

9.3. A-Periods. To describe the image of the integration mapping, consider the mapping $\bar{\partial}^{-1}: H_{f=0}^{-1}(\mathbb{CP}^1, \omega \otimes \bar{\omega}) \rightarrow H^0(\mathbb{CP}^1, \omega)$. It is a continuous elliptic operator. To make the formulae localizable, fix a top form β_0 with integral 1, and extend $\bar{\partial}^{-1}$ to any form on $\mathbb{CP}^1$ by $\bar{\partial}^{-1}\beta \stackrel{\text{def}}{=} \bar{\partial}^{-1}(\beta - \beta_0 \int_{\mathbb{CP}^1} \beta)$. This operator is still elliptic, but is not an isomorphism any more.

Let $\alpha \in \mathcal{H}^{(0)}$, consider $\alpha|_{\partial K_i}$. In contrast with the cases of Section 4.3 and Proposition 6.12 the $--$ -components of $(\alpha|_{\partial K_i})_{i \in I}$ are not arbitrary. To describe possible values of $(\alpha|_{\partial K_i})_{i \in I}$, break $\alpha|_{\partial K_i}$ into two parts, one $\alpha_i^{(0)}$ with integral 0, another one $\alpha_i^{(1)}$ proportional to some fixed 1-form μ_i on ∂K_i . Use the same letters for δ -function-extensions of these forms to $\mathbb{CP}^1$. Suppose that 1-forms μ_i (one per disk boundary) are normalized to have integral 1 and have uniformly bounded $H^{-1/2}$ -norms when i varies, and that disks K_i have a thickening. Then $\sum \alpha_i^{(1)}$ converges in $\bigoplus_{l_2} H^{-1/2}(\partial K_i)$, same for $\sum \alpha_i^{(0)}$. Note that if conditions of Theorem 4.11 hold, then $\alpha^{(0)} = \bar{\partial}^{-1}(\sum \alpha_i^{(0)}) \in H^0(\mathbb{CP}^1, \omega)$. Since $\alpha = \bar{\partial}^{-1} \sum_i \alpha|_{\partial K_i}$, $\alpha^{(1)} = \bar{\partial}^{-1}(\sum \alpha_i^{(1)}) = \alpha - \bar{\partial}^{-1}(\sum \alpha_i^{(0)})$ is also in $H^0(\mathbb{CP}^1, \omega)$.

We see that any form $\alpha \in \mathcal{H}^{(0)}$ can be represented as a sum $\alpha^{(0)} + \alpha^{(1)}$, $\alpha^{(0)}, \alpha^{(1)} \in \mathcal{H}^{(0)}$ such that integrals of $\alpha^{(0)}$ around each disk K_i vanishes, and $\alpha^{(1)}$ is a linear combination of 1-forms $\bar{\partial}^{-1}\mu_i$, $i \in I$. Consider the subspace $\mathcal{H}_0^{(0)} \subset \mathcal{H}^{(0)}$ consisting of 1-forms which satisfy $\int_{\partial K_i} \alpha = 0$ for any $i \in I$, and the subspace $\mathcal{H}_1^{(0)}$ of forms satisfying $\int_{\partial K_i} \alpha = \int_{\partial K'_i} \alpha$ for any $i \in I$. Let $\mathfrak{c} = \mathcal{H}^{(0)}/\mathcal{H}_0^{(0)}$, $\mathfrak{c}'' = \mathcal{H}_1^{(0)}/\mathcal{H}_0^{(0)}$. The decomposition $\alpha = \alpha^{(0)} + \alpha^{(1)}$ gives a splitting of $\mathcal{H}^{(0)}$ into a direct sum of $\mathcal{H}_0^{(0)}$ and the span of $\bar{\partial}^{-1}\mu_i$, $i \in I$. As a corollary, the images of forms $\bar{\partial}^{-1}\mu_i$ form a basis in $\mathfrak{c}$, and the Gram matrix of the pairing in $\mathfrak{c}$ can be calculated as $(\bar{\partial}^{-1}\mu_i, \bar{\partial}^{-1}\mu_j)_{L_2(\mathbb{CP}^1)}$. Similarly, projections of $\bar{\partial}^{-1}\mu_i - \bar{\partial}^{-1}\mu_{i'}$, $i \in I_+$, form a basis in $\mathfrak{c}''$, and one can easily calculate the Gram matrix of this basis. Let us estimate elements of these Gram matrices.

We may assume that $\infty \in K_{i_0}$ for some $i_0 \in I$, and that $\text{Supp } \beta_0 \subset K_{i_0}$. Then the operator $\bar{\partial}^{-1}$ restricted to $\mathbb{CP}^1 \setminus K_{i_0}$ has $\frac{dx}{y-x}$ as a kernel, and we may suppose that

μ_i is $d\vartheta_i/2\pi$, ϑ_i being the natural angle coordinate on $\partial K_i \subset \mathbb{C}$. Let us calculate the Gram matrix (G_{ij}) for $\bar{\partial}^{-1}d\vartheta_i \in \mathcal{H}^{(0)}$. If the radius of K_i is r_i , and the distance between centers of K_i and K_j is d_{ij} , then G_{ij} can be estimated as

$$\int_{\substack{r_i < |z| < C \\ |z - d_{ij}| > r_j}} \frac{1}{z(z - d_{ij})} dx dy$$

here $C \gg 0$ (and depends on β_0), $z = x + iy$. Note that the integral over $|z| > C/2$ is greater than a constant which does not depend on d_{ij} and r_i, r_j , thus this integral is bounded from below. If $i = j$, then it behaves as $\log \frac{1}{r}$. If $i \neq j$, then the part outside the disk of radius $2d_{ij}$ is $\sim \log \frac{1}{d_{ij}}$, and the part inside this disk is scaling-invariant. Thus to estimate the second part one may assume $d_{ij} = 1$. However, the integral converges absolutely inside the whole disk $|z| < 2$, thus this part is bounded.

Thus we have estimates for the elements of the Gram matrix¹⁶. Knowledge of this Gram matrix gives a complete description of the subspace $\mathcal{B}$ of $H^{-1}(\mathbb{CP}^1, \omega \otimes \bar{\omega})$ spanned by extensions-by- δ -function of forms on ∂K_i , $i \in I$.

Similarly one can estimate the elements of the Gram matrix for $\mathfrak{c}''$. Since elements of $\mathfrak{c}''$ correspond to 2-forms with integral 0, this Gram matrix does not change when we apply a conformal transformation to disks K_i . The diagonal elements are $\frac{l_{ii'}}{\pi}$, and the off-diagonal ones are $C \frac{1}{\pi} \log |\lambda(c_i, c_{i'}, c_j, c_{j'})| + O(\sum \varepsilon_k r_k)$, here λ is the double ratio, $c_\bullet$ and $r_\bullet$ are the center and radius of $K_\bullet$. The constants $\varepsilon_\bullet$ can be calculated as $c_i = \frac{c_j - c_{j'}}{(c_i - c_j)(c_i - c_{j'})}$.

Remark 9.11. Note that we have estimates for the elements of this Gram matrix. It is easy to check that if the matrix $(r_i/d_{ij})_{i,j \in I}$ gives a compact operator $l_2 \rightarrow l_2$, these estimates allow one to reconstruct the Hilbert norm up to equivalence, thus to describe the space $\mathfrak{c}''$ completely.

Note that the diagonal entries of the Gram matrix for $\mathfrak{c}$ are not bounded. Since any element in the image of the integration mapping has a finite norm w.r.t. this matrix, this shows, in particular, that the image of the integration mapping does not coincide with l_2 . Similarly, the Gram matrix for $\mathfrak{c}''$ contains arbitrarily big elements, thus $\mathfrak{c}''$ also differs from l_2 , thus image of partial period mapping differs from l_2 . As we will see it in Section 9.5, this image coincides with the space $\mathfrak{c}'$ defined below.

So far the spaces we consider depended on the relative position of disks K_i only. Now suppose that a fraction-linear orientation-changing identification φ_i of ∂K_i and $\partial K_{i'}$ is fixed, and consider the subspace $\mathcal{B}'$ of $\mathcal{B}$ which is spanned by extensions-by- δ -function of forms on ∂K_i , $i \in I$, which are preserved by the identifications φ_i . Let $\mathcal{B}_0$ be the subspace of $\mathcal{B}$ spanned by extensions-by- δ -function of forms on ∂K_i , $i \in I$, with integral 0 along each ∂K_i , let $\mathcal{B}'_0 = \mathcal{B}' \cap \mathcal{B}_0$.

¹⁶Note that these estimates may be not sufficient to describe the Hilbert structure on $\mathfrak{c}$ up to equivalence.

It is clear that $\mathcal{B}/\mathcal{B}_0 \simeq \mathfrak{c}$. Let us describe $\mathcal{B}'/\mathcal{B}'_0$. Obviously, the image of $\mathcal{B}'/\mathcal{B}'_0$ in $\mathfrak{c}$ is a subspace of $\mathfrak{c}''$, however, if the identifications φ_i “squeeze” the circles ∂K_i too much, this image may be a proper subspace of $\mathfrak{c}''$.

Indeed, the dual statement to Remark 4.5 shows that one can define a $\mathrm{PGL}(2, \mathbb{R})$ -invariant norm on $H_{f=0}^{-1/2}(S^1, \Omega^{\mathrm{top}})$ which is equivalent to Hilbert norm. However, there is no $\mathrm{PGL}(2, \mathbb{R})$ -invariant norm on the whole space $H^{-1/2}(S^1, \Omega^{\mathrm{top}})$. For $g \in \mathrm{PGL}(2, \mathbb{R})$ consider how g changes the Sobolev norm on $H^{-1/2}(S^1, \Omega^{\mathrm{top}})$. It is enough to estimate $N(g) = \|g^*(\beta)\|$, β being an arbitrary form with non-zero integral. It is clear that different choices of Sobolev norm on $H^{-1/2}(S^1, \Omega^{\mathrm{top}})$ ¹⁷ and different choices of β would change $N(g)$ to something of the form $CN(g) + O(1)$ only. Since $\mathrm{PGL}(2, \mathbb{R})$ has a compact subgroup $U(1)$ of dimension 1, it is enough to estimate $N(g)$ on double classes $U(1) \backslash \mathrm{PGL}(2, \mathbb{R}) / U(1)$ only, thus one can assume $g = \mathrm{diag}(\lambda, 1)$, $\lambda \geq 1$.

In turn, it is sufficient to estimate $H^{-1/2}$ -norm of $\frac{\lambda dx}{1+\lambda^2 x^2}$ on $\mathbb{R}$, and one can do it explicitly since the Fourier transform can be easily calculated, it is proportional to $e^{-\lambda|\xi|}$. Thus $N(g) = C \log \lambda + O(1)$. Using this, one can easily obtain

Proposition 9.12. *Consider the image $\mathfrak{c}'$ of $\mathcal{B}'/\mathcal{B}'_0$ in $\mathfrak{c} = \mathcal{B}/\mathcal{B}_0$. Consider φ_i as elements of $\mathrm{SL}(2, \mathbb{R}) \subset \mathbb{R}^4$. The space $\mathfrak{c}'$ consists of sequences $(p_i) \in \mathfrak{c}''$ which satisfy $(p_i \log \|\varphi_i\|) \in l_2$.*

We see that $\mathfrak{c}'$ can be described by Gram matrix $G''_{ij} = \delta_{ij} \log \|\varphi_i\| + G''_{ij}$, G'' being the Gram matrix for $\mathfrak{c}''$.

9.4. B-periods. What we defined in Section 9.2 was integration of global holomorphic forms along A -cycles. Finite-genus theory shows that it is important to study additional integrals along B -cycles. They should depend on the choice of cuts on the Riemann surface, and we are not in the conditions when one can easily proceed with such cuts.

Since we have only H^0 -smoothness of global holomorphic forms, one cannot invent a priori bounds on integrals of these forms along arbitrary curves. Indeed, the “infinity” $M_\infty \subset \mathbb{CP}^1$ can break the complex sphere into infinitely many connected components (see the example in Section 0.10), thus one cannot assume that the cuts do not intersect M_∞ . Moreover, even if there is only one connected component, it is not clear how to make infinitely many cuts in $\mathbb{CP}^1$ which would not intersect each other. In fact it *is* possible to make such cuts, but in general the lengths of these cuts form a quickly increasing sequence.

However, if one has a strip $(-\varepsilon, \varepsilon) \times (a, b)$ embedded into the complex curve, then the value of $\int_a^b$ averaged along $(-\varepsilon, \varepsilon)$ is well defined and may be bounded as

¹⁷Recall that it is defined up to equivalence only.

$O(\varepsilon^{-1/2})$. Indeed, this average is L_2 -pairing

$$\frac{1}{2\varepsilon} \int \alpha \wedge dx$$

with dx/ε , here (x, y) are coordinates on the strip $-\varepsilon < x < \varepsilon$.

Thus we are not going to define the B -periods as integrals over curves, but as some averaged integrals. On the level of would-be homology of the surface we will pair the form with cycles on the surface which are not “geometric” cycles, but linear combinations of them.

Consider the involution $'$ and gluings φ_i . Suppose that $0 \in K_{i_0}$, $\infty \in K_{i'_0}$, and that the disks $\{K_i\}$ have a uniform thickening. Parameterize the set of rays going from 0 to ∞ by the angle ϑ , and deform each ray slightly in such a way that it avoids all the circles K_i . We require that deformations of two rays which differ by the angle $\Delta\vartheta$ do not become closer than $\varepsilon \cdot \Delta\vartheta$ (outside of K_{i_0}). Say, let $\tilde{K}_i$ be the concentric with K_i disk of radius $(1 + 2\varepsilon)\text{radius}(K_i)$, and deform the ray $\mathcal{R}$ inside $\tilde{K}_i$ so that it moves along an appropriate arc going between K_i and $\tilde{K}_i$. If the $\text{dist}(\mathcal{R}, \text{center}(K_i)) / \text{radius}(\tilde{K}_i) = \rho \leq 1$, one can take the radius of the arc to be $(1 + \varepsilon + \varepsilon\rho)\text{radius}(K_i)$.

In fact we need to choose whether the arcs are going to leave the disk K_i on the right or on the left. We use the following algorithm: let O_i be the fixed point of φ_i inside K_i . If the ray $\mathcal{R}$ leaves O_i on the right, let the deformation leave K_i on the right, otherwise leave it on the left. (This choice is going to be important in the proof of Proposition 9.21. Note that the choice of the direction of the turnout does not depend on the metric on $\mathbb{CP}^1$.)

Now we assume that 0 and ∞ are fixed points of φ_{i_0} . As a last correction, in a neighborhood of $K_{i'_0}$ change a ray $\mathcal{R}$ to a logarithmic spiral so that the intersection of $\mathcal{R}$ with ∂K_{i_0} and intersection of $\mathcal{R}$ with $\partial K_{i'_0}$ are glued together by φ_{i_0} . Say, if $\partial K_{i'_0} = \{z \mid |z| = R\}$, take the part of a spiral $d\vartheta = C dr/\varepsilon r$ inside $\{z \mid e^{-\varepsilon}R < |z| < R\}$, here $C = \text{Arg } \varphi_{i_0}$ is defined up to addition of a multiple of 2π . Now each deformed ray represents a closed curve after the gluing by φ_{i_0} is performed.

Under these conditions the averaged over $\vartheta \in [0, 2\pi]$ integral over the deformed rays (between ∂K_{i_0} and $\partial K_{i'_0}$) is correctly defined. It represents a combination of cycles, thus is a cycle itself. Denote this linear functional on $\mathcal{H}^{(0)}(\mathbb{CP}^1, \{K_i\})$ by $\mathcal{Q}_{i_0}$.

Restricting this linear functional to global holomorphic forms, we call the integral along this cycle B -period of the global holomorphic form from ∂K_{i_0} to $\partial K_{i'_0}$. The above description shows that

Lemma 9.13. *B -period of α from ∂K_i to $\partial K_{i'}$ is bounded by $C \cdot l_{ii'} \|\alpha\|_{L_2}$, here $l_{ii'}$ is the conformal distance between ∂K_{i_0} and $\partial K_{i'_0}$. The constant C depends on ε only.*

Definition 9.14. Associate to $\alpha \in \Gamma(\bar{M}, \omega)$ the sequence $(q_j)_{j \in I_+}$, q_j being the B -period of α from ∂K_j to $\partial K_{j'}$. Denote this mapping by Q .

Remark 9.15. Note that in the case of finite genus one chooses the B -cycles to be non-intersecting. This assures that the matrix of periods is symmetric. The above construction takes average of different cycles connecting given points, thus one can get an impression that the resulting matrix of periods will have much worse properties than in the standard settings.

However, the properties turn out to be exactly the same, due to our choice of direction of turnouts, and the following surprising result:

Proposition 9.16. *Consider 4 different points $x_0, x_1, y_0, y_1 \in \mathbb{CP}^1$. Parameterize the set of (circular) arcs connecting x_0 with x_1 by the angle at x_0 , and do the same with arcs connecting y_0 with y_1 . This provides a measure on the set of arcs connecting x_0 with x_1 , same for y_0 and y_1 . Then the average index of intersection of an arc x_0x_1 with an arc y_0y_1 is 0.*

9.5. Space of periods. In Section 9.3 we have shown that under mild assumption the space $\mathcal{H}^{(0)}$ (which, loosely speaking, consists of holomorphic forms with jumps along the cuts on the curve) is a sum of two components: forms with integrals 0 along A -cycles (which can be described by locale data on each cut), and some explicitly defined Hilbert subspace $\mathfrak{c}$ of “small” dimension (two basis vectors per each cut on the curve). Only the elements of $\mathcal{H}^{(0)}$ which have equal integrals along two sides of the cut have a chance to correspond to a global holomorphic form on $\bar{M}$, thus only subspace $\mathfrak{c}' \subset \mathfrak{c}$ is interesting for us.

Theorem 9.17. *Suppose that the matrix $(e^{-l_{ij}} - \delta_{ij})$ defines a compact operator $l_2 \rightarrow l_2$, and the disks K_i have a thickening. Then the partial period mapping P sends global holomorphic forms to elements of $\mathfrak{c}'$. Moreover, it is an isomorphism onto $\mathfrak{c}'$.*

Proof. The first statement is an immediate corollary of the description of the elements of $\mathcal{H}^{(0)}$ via the space $\mathcal{B}'$ in Section 9.3. We start the proof of the second one by showing that P is a component of a Fredholm operator of index 0.

In the notations of Section 9.3, consider the mapping $\mathcal{H}_1^{(0)} \xrightarrow{\pi} \mathfrak{c}''$. Let $\mathcal{H}_2^{(0)} = \pi^{-1}(\mathfrak{c}')$. Then the mapping

$$\mathcal{J}_\omega: \mathcal{H}_2^{(0)} \rightarrow \bigoplus_{\substack{l_2 \\ i \in I_+}} H_{f=0}^{-1/2}(\partial K_i, \Omega_{\partial K_i}^1) : \alpha \mapsto (\alpha|_{\partial K_i} - \varphi^*(\alpha|_{\partial K_{i'}})) .$$

is continuous. The same arguments as in the proof of Theorem 4.11 show that $\mathcal{J}_\omega|_{\mathcal{H}_2^{(0)}}$ is a Fredholm operator of index 0. Combining this mapping with the projection $\mathcal{H}_1^{(0)} \xrightarrow{\pi} \mathcal{H}_2^{(0)}/\mathcal{H}_0^{(0)} = \mathfrak{c}'$, we see that $\alpha \mapsto (\mathcal{J}_\omega(\alpha), \pi(\alpha))$ is a Fredholm mapping of index 0. If α is in the null-space of this mapping, then α is a non-trivial global holomorphic form with vanishing A -periods, thus $\alpha \in \text{Ker } P$, and $\alpha = 0$. Hence this mapping is an isomorphism. Since $P = \pi|_{\text{Ker } \mathcal{J}_\omega}$, it is an isomorphism as well. $\square$

Let $\alpha_i = \bar{\partial}^{-1}\beta_i$. The theorem says that for any element $(c_i) \in \mathfrak{c}'$ one can find an element $\alpha \in \mathcal{H}^{(0)}$ such that integrals of α along ∂K_i are 0 for any $i \in I$, and $\sum c_i \alpha_i + \alpha$ is a global holomorphic form (which automatically has A -periods c_i). Moreover, $\|\alpha\|_{\mathcal{H}^{(0)}} \leq C \cdot \|(c_i)\|_{\mathfrak{c}'}$.

Proposition 9.18. *Let $\beta \in \mathcal{H}^{(0)}$, and integrals of β along ∂K_i are 0 for any $i \in I$, let $(c_i) \in \mathfrak{c}'$. Then $\left| \sum_{j \in I_+} c_j \mathcal{Q}_j(\beta) \right| \leq C \|\beta\|_{\mathcal{H}^{(0)}} \cdot \|(c_i)\|_{\mathfrak{c}'}$ for an appropriate C which does not depend on β and (c_i) .*

Proof. Suppose that β can be holomorphically continued into all disks K_i except K_{i_0} . Then $\beta = \partial f$, and f is holomorphic outside K_{i_0} . Let $j, j' \neq i_0$. The construction of $\mathcal{Q}_j$ shows that one can calculate $\mathcal{Q}_j(\beta)$ as $f(y_1) - f(y_0)$, y_0, y_1 being two fixed points of φ_j . Let Ψ_j be the fraction-linear function with a zero and a pole at y_0 and y_1 . One can momentarily see that $\mathcal{Q}_j(\beta) = \int_{\partial K_{i_0}} \beta \log \Psi_j$. Similarly, $\mathcal{Q}_{i_0}(\beta) = 0$.

Similar statements are true for forms β such that they can be holomorphically continued into all the disks K_i except a finite number. Since any form β which satisfies conditions of the proposition can be approximated by such forms, we see that it is sufficient to show that $\sum_{j \neq i, i'} c_j \log \Psi_j$ converges in $\bigoplus_{l_2} H^{1/2}(\partial K_i) / \text{const}$, or that $\sum_{j \neq i, i'} c_j d\Psi_j / \Psi_j$ converges in $\bigoplus H^{-1/2}(\partial K_i, \Omega^1)$. In turn, it is sufficient to show convergence of $\sum_{j \neq i, i'} c_j d\Psi_j / \Psi_j$ in $\bigoplus_{l_2} H^0(K_i, \omega) = L_2(\bigcup K_i, \omega)$. On the other hand, if σ_j is 1 in K_j and $K_{j'}$, and 0 otherwise, then $\bar{\partial} \sum_{j \neq i, i'} c_j \sigma_j d\Psi_j / \Psi_j = \sum_{j \neq i, i'} c_j \beta_j \in H^{-1}$, hence $\sum c_j \sigma_j d\Psi_j / \Psi_j$ converges in L_2 . $\square$

Proposition 9.19. *Let $(d_i), (c_i) \in \mathfrak{c}'$, and $\alpha \in \mathcal{H}^{(0)}$ such that integrals of α along ∂K_i are 0 for any $i \in I$, and $\alpha_c = \sum c_i \alpha_i + \alpha$ is a global holomorphic form on $\bar{M}$ (automatically with A -periods c_i). Then $\sum d_j \mathcal{Q}_j(\alpha_c) \leq C \|(c_i)\|_{\mathfrak{c}'} \cdot \|(d_i)\|_{\mathfrak{c}'}$.*

Proof. It is sufficient to show that $\sum d_j \mathcal{Q}_j(\sum c_i \alpha_i) \leq C_1 \|(c_i)\|_{\mathfrak{c}'} \cdot \|(d_i)\|_{\mathfrak{c}'}$. In the notations of the previous proposition $\alpha_i - \alpha_{i'} = d\Psi_i / \Psi_i$ outside of K_i and $K_{i'}$. One can assume that only a finite number of c_i and d_i is non-zero. Then $\sum_{i \in I} c_i \alpha_i$ is $\sum_{i \in I_+} c_i d\Psi_i / \Psi_i$.

Now it should be obvious that $\mathcal{Q}_j(\alpha_i) = \int \sigma_i \sigma_j \alpha_i \wedge \alpha_j$, which finishes the proof. $\square$

9.6. Period matrix. Since P is an isomorphism $\Gamma(\bar{M}, \omega) \rightarrow \mathfrak{c}'$, one can consider the mapping $\Omega = Q \circ P^{-1}$. Write this mapping using coordinate “basis” in $\mathfrak{c}'$:

Definition 9.20. Let $\tilde{\alpha}_j, j \in I_+$, be the global holomorphic form on $\bar{M}$ such that $P(\tilde{\alpha}_i)$ has 1 on j -th position, -1 at j' -position, 0 at the other positions. Let Ω_{ij} be the B -period of $\tilde{\alpha}_j$ from ∂K_i to $\partial K_{i'}$, $i \in I_+$.

Proposition 9.21. *Let $\alpha \in \Gamma(\bar{M}, \omega)$ and the sequence $(p_j) = P(\alpha)$ has only a finite number of non-zero elements. Let $Q(\alpha) = (q_j)$. Then $\|\alpha\|_{L_2}^2 = i \sum_{I_+} (\bar{p}_j q_j - p_j \bar{q}_j)$.*

Proof. Consider a representative of α in $\mathcal{H}^{(0)}(\mathbb{CP}^1, \{K_i\})$. Since all the A -periods of α but a finite number are 0, α can be extended (without changing the norm too much) into all the disks but a finite number preserving the closeness. Consider the remaining disks. Since the integral of α around K_i is opposite to the integral around $K_{i'}$, we see that if we connect K_i and $K_{i'}$ by a cut, then the integral of α around the resulting hole is 0. Make smooth cuts which connect the remaining disks pairwise (according to ') and do not intersect. One can suppose that two ends of the cut—one on ∂K_i , another on $\partial K_{i'}$ —are identified by φ_i .

After the cuts are performed, on the resulting domain 1-form α is closed, and the integral along any component of (piecewise-smooth) boundary is 0. Thus one can write $\alpha = df$, f being a function of smoothness H^1 . The restriction of f to any smooth curve is well-defined, and is of smoothness $H^{1/2}$. When one goes from one side of the cut to another one along ∂K_i , f grows by p_i . Let $\mathring{q}_i$ be the change of f when one goes from ∂K_i to $\partial K_{i'}$ along the cut (choosing the side of the cut so that the direction is counterclockwise).

Note that f is holomorphic near ∂K_i and $\partial K_{i'}$ thus the *value* of f at points is well-defined, thus the change of f along the cut is well-defined. (In generic point z of the cut $f(z)$ is not correctly defined, since f is only of smoothness $H^{1/2}$.)

Let γ_i be the part of the boundary of the domain consisting of the circles ∂K_i , $\partial K_{i'}$ and both sides of the cut which connects them.

Now take into account that $\|\alpha\|_{L_2}^2 = i \int \alpha \wedge \bar{\alpha} = i \int \partial f \wedge \bar{\partial} \bar{f}$. Proceeding as in Section 9.1, we see that the only change to the arguments is that instead of taking integrals along $\partial K_i \cup \partial K_{i'}$, one needs to take some integrals along γ_i . As there, the integral along $\partial K_i \cup \partial K_{i'}$ vanishes, so what remains is

$$\|\alpha\|_{L_2}^2 = -i \sum_i \int_{\gamma_i} f d\bar{f}$$

summation being over i such that $p_i \neq 0$. The cycle γ_i consists of 4 parts: two going around ∂K_i and $\partial K_{i'}$, another two going along sides of the cut. On the first two parts $d\bar{f}$ are identified via φ_i (with opposite signs), and f differs by $\mathring{q}_i$, thus the total integral is $-\bar{p}_i \mathring{q}_i$. On the second two $d\bar{f}$ coincide (but the orientation is opposite), and f differs by p_i , thus the total integral is $p_i \overline{\mathring{q}_i}$.

What remains to prove is that we may substitute q_i instead of $\mathring{q}_i$. To do this one needs to investigate the relationship between q_i and $\mathring{q}_i$. First, one can describe $\mathring{q}_i$ as an integral of df along one side of the cut. Indeed, though $d(f|_{\gamma_i})$ is of smoothness $H^{-1/2}$ on each part of γ_i , it is actually analytic near circles ∂K_i , $\partial K_{i'}$, thus the pairing with the fundamental cycle of the interval (which has jumps at the ends of the interval!) is well-defined.

Second, one can suppose that the cuts are in fact piecewise-smooth (as far as non-smooth points are in $\mathbb{CP}^1 \setminus \overline{\bigcup K_i}$), and consist of arcs of circles. Let us recall that q_i is the average integral of α along “rays” which connect 0 and ∞ (after an

appropriate choice of coordinate system). Here “rays” are curves with differ from rays inside circles $\tilde{K}_j$ only, and consist of arcs of circles. What is more, one does not need to deform a ray into a “ray” inside the circle K_j as far as $p_j = 0$, since a closed continuation of α inside K_j is already fixed, thus the deformation does not change the value of the integral along the “ray”. Note that each ray intersects cuts along a finite number of points (except a finite number of rays which may contain whole pieces of cuts), thus the same is true for “rays”.

We see that the integral of α along each “ray” is now well defined (recall that in general setting only the average was well-defined), moreover, it is easy to calculate this integral using representation $\alpha = df$. The integral along a “ray” $\mathcal{R}$ is equal to the change of f on the ends minus the jumps of f at the finite number of points where “ray” intersects cuts, thus it is

$$f(\text{end}(\mathcal{R})) - f(\text{start}(\mathcal{R})) - \sum_j n_{ij} p_j, \quad n_{ij} \in \mathbb{Z}, i, j \in I.$$

The change of f is equal to $\overset{\circ}{q}_i$, thus the average value of the integral is $\overset{\circ}{q}_i$ minus sum of some real multiples of p_j :

$$q_i = \overset{\circ}{q}_i - \sum_j \nu_{ij} p_j, \quad \nu_{ij} \in \mathbb{R}, i, j \in I,$$

and ν_{ij} are averaged values of n_{ij} .

Finally, use our choice of direction of turnout around the disk K_i . It insures that the following fact is true:

Lemma 9.22. $\nu_{ij} = \nu_{ji}$ if $i \neq j$.

Proof. Extend the cut between ∂K_j and $\partial K_{j'}$ to fixed points of φ_j (one inside each of K_j and $K_{j'}$) along straight intervals. Call the resulting curve γ'_j . Then $2\pi\nu_{ij}$ is the change of $\text{Arg } z$ along the curve γ'_j (we again suppose that 0 and ∞ are fixed points of φ_i). Similarly, $\pi\nu_{ji}$ is the change of $\text{Arg } \Psi_j(z)$ along γ'_i , here $\Psi_j(z)$ is a fraction-linear function with a pole and a zero at fixed points of φ_j .

We need to show that $\text{Im} \int_{\gamma'_j} \frac{dz}{z} + \int_{\gamma'_i} \frac{d\Psi_j(z)}{\Psi_j(z)} = 0$. However,

$$\int_{\gamma'_j} \frac{dz}{z} = \frac{1}{2\pi i} \int_{\gamma'_j} \frac{dz}{z} \text{Jump}(\log \Psi_j(z)) = \frac{1}{2\pi i} \int_{\gamma''_j} \log \Psi_j(z) d \log z$$

here we take an arbitrary branch of $\log \Psi(z)$ defined outside of γ'_j , and γ''_j is a loop around γ'_j . Now the identity is obvious, since γ''_j is homotopic to $-\gamma''_i$. $\square$

The lemma implies that plugging in $\overset{\circ}{q}_i$ instead of q_i into the formula of the proposition gives the same value, which finishes the proof of Proposition 9.21. $\square$

Proposition 9.23. *Let $\alpha, \beta \in \Gamma(\bar{M}, \omega)$ and the sequences $(p_j) = P(\alpha)$, $(p'_j) = P(\beta)$ have only a finite number of non-zero elements. Let $Q(\alpha) = (q_j)$, $Q(\beta) = (q'_j)$. Then $\sum_{I_+} (p_j q'_j - p'_j q_j) = 0$.*

Proof. The proof of Proposition 9.21 with minor changes is applicable, the integral to consider is $0 = \int \alpha \beta$. $\square$

The following statement is an immediate corollary of Proposition 9.19:

Proposition 9.24. *In the conditions of the previous proposition $\sum_{I_+} p_j q'_j \leq C \|\alpha\| \cdot \|\beta\|$.*

Corollary 9.25. *The period matrix Ω_{ij} is symmetric, $\text{Im } \Omega_{ij}$ is a positive real symmetric matrix, and the matrix Ω_{ij} defines a bounded symmetric form on the Hilbert space with the pairing given by the Gram matrix $\text{Im } \Omega_{ij}$.*

9.7. Bounded Jacobian as a torus. In Section 9.1 we have seen that (under mild assumptions) the Jacobian coincides with the constant Jacobian. Given a topological space S and a set I , let $S_{l_\infty}^I \stackrel{\text{def}}{=} \bigcup_K K^I$, here K runs over compact subsets of S . Obviously, there is a surjection from $(\mathbb{C}^*)_{l_\infty}^{I+}$ to the constant bounded Jacobian, which sends a sequence $(\psi_i)_{i \in I}$ such that $\psi_i = \psi_i^{-1}$ into a corresponding line bundle. Here we are going to show that the kernel of this surjection is a lattice in $(\mathbb{C}^*)_{l_\infty}^{I+}$, as in finite-genus case. To avoid defining a lattice in $(\mathbb{C}^*)_{l_\infty}^{I+}$, consider $(\mathbb{C}^*)_{l_\infty}^{I+}$ as the set of coordinate-wise exponents of $\mathbb{C}_{l_\infty}^{I+}$. We obtain a mapping from $\mathbb{C}_{l_\infty}^{I+}$ to the bounded Jacobian which sends $(2\pi i \mathbb{Z})_{l_\infty}^{I+}$ to the origin in the Jacobian.

We are going to prove that there is a lattice $L \subset \mathbb{C}^{I+}$ which goes to the origin. First, construct generators of this lattice:

Proposition 9.26. *Suppose that the matrix $(e^{-l_{ij}} - \delta_{ij})$ gives a compact operator $l_2 \rightarrow l_2$ and a compact operator $l_1 \rightarrow l_1$. Let $j \in I_+$. There exists a global holomorphic form $\alpha^{(j)}$ on $\bar{M}$ such that the A -periods of $\alpha^{(j)}$ vanish except for the j -th one, which is equal to $2\pi i$. Then $a_j = \exp \int \alpha^{(j)}$ is an element of $\mathcal{H}^\infty$. The corresponding cocycle $\psi_{kj} = a_j \circ \varphi_k / a_j$ is (locally) constant, and coincides with $\exp 2\pi i \Omega_{kj}$.*

Proof. Existence of $\alpha^{(j)}$ is a corollary of results of Section 9.5. The only statement we need to prove is that $a_j \in \mathcal{H}^\infty$. In turn, it is sufficient to prove that $\text{Im } \Omega_{kj}$ is bounded (for a fixed j). On the other hand, $\text{Im } \Omega_{kj}$ consists of two parts which correspond to decomposition $\alpha^{(j)} = 2\pi i \alpha_j + o^{(j)}$, here α_j is defined as in Section 9.5, and $o^{(j)} \in \mathcal{H}^{(0)}$ and has A -periods 0. The description of the cycle for B -period shows that first part of $\text{Im } \Omega_{kj}$ is bounded by $\log |l_{jj'}|$, thus it is sufficient to estimate the second part $\text{Im } \mathcal{Q}_k(o^{(j)})$.

This estimate follows from the following lemma:

Lemma 9.27. *Suppose that the matrix $(e^{-l_{ij}} - \delta_{ij})$ gives a compact operator $l_2 \rightarrow l_2$ and a compact operator $l_1 \rightarrow l_1$.*

1. *Let $\alpha \in H^{-1/2}(\partial K_i, \Omega^{top})$. Denote by the same letter extension-by- δ -function of α to $\mathbb{CP}^1$. Then $\bar{\partial}^{-1}\alpha \in \mathcal{H}^{(0)}$, and $\mathcal{Q}_k(\bar{\partial}^{-1}\alpha) = O(\|\alpha\|_{H^{-1/2}})$ uniformly in $k \neq i$.*
2. *The sequence $(\|o^{(j)}|_{\partial K_k}\|_{H^{-1/2}})_{k \in I} \in l_1$.*

Proof. The first part follows from the explicit construction of cycles for B -periods. Prove the second part.

Since sum of radii of disks K_i is finite, one can estimate that the sequence $(\|\alpha_j|_{\partial K_k}\|_{H^{-1/2}})_{k \in I} \in l_1$. Denote the space of such 1-forms on $\bigcup_i \partial K_i$ by $\mathcal{H}_1$. On the other hand, $o^{(j)}$ is uniquely determined by the conditions $\mathcal{J}_\omega o^{(j)} = -2\pi i \mathcal{J}_\omega \alpha_j$ and $o^{(j)} \in \mathcal{H}_0^{(0)}$, here $\mathcal{J}_\omega$ is the operator from Section 9.5. What remains to prove is that $\mathcal{J}_\omega$ sends the subspace $\mathcal{H}_0^{(0)} \cap \mathcal{H}_1$ onto itself. We know that $\mathcal{J}_\omega$ sends $\mathcal{H}_0^{(0)}$ onto itself. However, from the restrictions on l_{ij} one immediately obtains that $\mathcal{J}_\omega$ gives a Fredholm operator $\mathcal{H}_1 \rightarrow \mathcal{H}_1$ of index 0. Since $\mathcal{J}_\omega$ has no kernel, this operator is an isomorphism. $\square$

This finishes the proof of Proposition 9.26. $\square$

Definition 9.28. Call a multiplier $a \in \mathcal{H}^\infty$ *lattice-like*, if $a^{-1} \in \mathcal{H}^\infty$ and $\psi_k = a^{-1} \cdot (a \circ \varphi_k)$ is a constant function on ∂K_k for every $k \in I$.

Now to each $j \in I_+$ we associated a lattice-like multiplier a_j , which induces a cocycle $(\psi_k)_{k \in I_+}$, $\psi_k = \exp 2\pi i \Omega_{kj}$. The bundle given by this cocycle is isomorphic to a trivial one (via a_j). Since $\text{Im } \Omega_{kj}$ gives a positive Hermitian form (defined at least on real sequences of finite length), columns of $\text{Im } \Omega_{kj}$ generate the subspace which is dense in the set of real sequences (with topology of direct product), which shows in conditions of Proposition 9.26

Proposition 9.29. *Consider the space $\mathbb{C}_{l_\infty}^{I_+}$ of bounded sequences (Ψ_k) with topology induced from the direct product $\mathbb{C}^{I_+}$. Consider the subgroup L generated by rows (Ω_{kj}) , $j \in I_+$, and $\mathbb{Z}_{l_\infty}^{I_+}$. This subgroup is a lattice, i.e., its $\mathbb{R}$ -span is dense in $\mathbb{C}_{l_\infty}^{I_+}$.*

To finish the description of bounded Jacobian, it remains to prove that any lattice-like multiplier a is a product of powers of a_j , $j \in I_+$. Recall that the index of function a is the change of $\frac{\text{Arg } a}{2\pi i}$ along a contour. Note that the *index* of a_j around ∂K_k is ± 1 if $k = j, j'$, and is 0 otherwise. Thus one reconstruct the needed powers of a_j by taking indices of a around ∂K_j , $j \in I_+$. The only things we need to prove is that a lattice-like multiplier a with all indices around ∂K_j being 0 is constant, and only a finite number of indices of a lattice-like multiplier is non-zero. The first statement is a direct corollary of Lemma 9.2 and the following

Lemma 9.30. *Suppose that the conformal distances l_{ij} satisfy the condition that $(e^{-l_{ij}} - \delta_{ij})$ gives a bounded operator $l_\infty \rightarrow l_2$. Then any lattice-like multiplier is of class $\mathcal{H}^{(1)}$.*

Recall that the space $\mathcal{H}^{(1)}$ was defined in Section 4.4.

Proof. Since $a dz^{1/2} \in \mathcal{H}(\mathbb{CP}^1, \{K_j\})$, Theorem 6.7 shows that a can be reconstructed from its restriction to $\bigcup_{j \in I} \partial K_j$ using Cauchy formula. In particular, $+$ -part of ∂a on ∂K_j is given by an integral along $\bigcup_{k \neq j} \partial K_k$. The restrictions on l_{ij} guarantee that the $+$ -parts of ∂a on the circles ∂K_j is in $\oplus_{l_2} H^{-1/2}(\partial K_j)$ modulo constants. Since $a^{-1} \cdot (a \circ \varphi_k)$ is constant and bounded, $-$ -parts of ∂a are also in this space. Since the integral of ∂a along each circle vanishes, arguments similar to ones in Section 4.4 show that $\partial a \in L_2(\mathbb{CP}^1 \setminus \bigcup K_i, \omega)$, thus $a \in \mathcal{H}^{(1)}$. $\square$

Iteration of the procedure used in the proof of the lemma shows that

Amplification 9.31. *Conclusions of Lemma 9.30 remain true if $(e^{-l_{ij}} - \delta_{ij})$ gives a bounded operator $l_\infty \rightarrow l_\infty$, and some power of this operator sends l_∞ to l_2 .*

To prove that only a finite number of indices of a lattice-like multiplier a is non-zero, it is enough to show that $a^{-1} \partial a \in L_2(U, \omega)$, U being homotopic to $\mathbb{CP}^1 \setminus \bigcup K_i$. Indeed, indices of a are proportional to periods of $a^{-1} \partial a$, and periods of L_2 -form form a sequence in l_2 .

In turn, since a^{-1} is a multiplier, thus is bounded, this follows from the fact that $\partial a \in L_2(U, \omega)$. We obtain

Theorem 9.32. *Consider a family of non-intersecting disks $K_i \subset \mathbb{CP}^1$ with pairwise conformal distances. Let $\mathcal{R}$ be the matrix $(e^{-l_{ij}} - \delta_{ij})$. Suppose that $\mathcal{R}$ gives a compact mapping $l_2 \rightarrow l_2$ and a bounded mapping $l_\infty \rightarrow l_\infty$, and that some power of $\mathcal{R}$ gives a bounded mapping $l_\infty \rightarrow l_2$. If disks K_i have a uniform thickening, then the bounded Jacobian coincides with the quotient of $\mathbb{C}_{l_\infty}^{I+}$ by the lattice generated by $\mathbb{Z}_{l_\infty}^{I+}$ and rows of the period matrix (Ω_{ij}) .*

Remark 9.33. Formally speaking, we defined the Jacobians in the case when $(e^{-l_{ij}/2} - \delta_{ij})$ gives a compact operator $l_2 \rightarrow l_2$, thus the statement of theorem is abuse of notations. However, it is easy to define all the ingredients needed for the definition of the bounded Jacobian as far as the conditions of Amplification 9.31 are satisfied.

Let us recall that the conditions of the theorem are automatically satisfied for any Hilbert–Schmidt curve, thus we get a complete description of the bounded Jacobian in this case. It is similar to the finite-genus case, where Jacobian is a quotient of a finite-dimensional complex vector space by a lattice.

Moreover, note that the topology on the constant bounded Jacobian is inherited from the topology of direct product on $\mathbb{C}^{I+}$, thus the above description allows one to reconstruct the topology on the bounded Jacobian as well.

9.8. Rigged Hodge structure. Let us wrap the results of the previous section into the familiar form of Hodge structures. In fact the resulted structure will be a hybrid of a Hodge structure and a structure of a rigged topological vector space.

Let I_+ be an arbitrary set. Let $H_{\mathbb{Z}}^A, H_{\mathbb{R}}^A, H_{\mathbb{C}}^A$ be the spaces of sequences $(p_i)_{i \in I_+}$ with only a finite number of non-zero terms (with integer/real/complex terms), $H_{\mathbb{Z}}^B, H_{\mathbb{R}}^B, H_{\mathbb{C}}^B$ be the spaces of sequences $(q_i)_{i \in I_+}$ without any restriction on growth, and $H_{\mathbb{Z}} = H_{\mathbb{Z}}^A \oplus H_{\mathbb{Z}}^B, H_{\mathbb{R}} = H_{\mathbb{R}}^A \oplus H_{\mathbb{R}}^B, H_{\mathbb{C}} = H_{\mathbb{C}}^A \oplus H_{\mathbb{C}}^B$. One can define an operation of complex conjugation on the spaces $H_{\mathbb{C}}^\bullet$, this operation leaves $H_{\mathbb{R}}^\bullet$ fixed. The spaces $H_{\mathbb{Z}}, H_{\mathbb{R}}, H_{\mathbb{C}}$ have a natural symplectic structure

$$[((p_i), (q_i)), ((p'_i), (q'_i))] = \sum_i (p_i q'_i - p'_i q_i)$$

such that the components $H_{\bullet}^{A,B}$ are Lagrangian and mutually dual. Fix an arbitrary mapping $\Omega: H_{\mathbb{C}}^A \rightarrow H_{\mathbb{C}}^B$. Let $\Omega_1: H_{\mathbb{C}}^A \rightarrow H_{\mathbb{C}}^A \oplus H_{\mathbb{C}}^B = H_{\mathbb{C}}$ be $\text{id} \oplus \Omega$. Denote $\text{Im } \Omega_1 \subset H_{\mathbb{C}}$ by $H^{1,0}$. Suppose that (compare Proposition 9.23) $H^{1,0}$ is Lagrangian. Moreover, suppose (compare Proposition 9.21) that if $\alpha \in H^{1,0}$, then $\text{Im} [\alpha, \bar{\alpha}] \gg 0$ in the following sense:

1. $\text{Im} [\alpha, \bar{\alpha}] \geq 0$ (here $\alpha \mapsto \bar{\alpha}$ is the complex conjugation on $H_{\mathbb{C}}$);
2. the equality is achieved only if $\alpha = 0$;
3. the completion $\mathfrak{h}^{1,0}$ of $H^{1,0}$ w.r.t. the norm $\|\alpha\|^2 = \text{Im} [\alpha, \bar{\alpha}]$ has no vectors of length 0.

Let $H^{0,1} = \{\alpha \in H_{\mathbb{C}} \mid \bar{\alpha} \in H^{1,0}\}$. The mapping Ω_1 gives an inclusion i of $H_{\mathbb{C}}^A$ into $\mathfrak{h}^{1,0}$.

Definition 9.34. A *rigged Hodge structure* is a subspace $H^{1,0} \subset H_{\mathbb{C}}$ which satisfies the above conditions, and such that the projection $p: H^{1,0} \rightarrow H_{\mathbb{C}}^B$ can be continuously extended to a mapping $p: \mathfrak{h}^{1,0} \rightarrow H_{\mathbb{C}}^B$.

Note the mappings $H_{\mathbb{C}}^A \xrightarrow{i} \mathfrak{h}^{1,0} \xrightarrow{p} H_{\mathbb{C}}^B$ equip $\mathfrak{h}^{1,0}$ with a structure of a rigged topological vector space [5]. It is clear that $H^{1,0}$ is a rigged Hodge structure iff Ω is symmetric, $\text{Im } \Omega > 0$, $-C \cdot \text{Im } \Omega < \text{Re } \Omega < C \cdot \text{Im } \Omega$.

From now on suppose that $H^{1,0}$ is a rigged Hodge structure. Let $\mathfrak{h}^{0,1}$ be the complexly conjugated space to $\mathfrak{h}^{1,0}$, $\mathfrak{h}_{\mathbb{C}} = \mathfrak{h}^{1,0} \oplus \mathfrak{h}^{0,1}$. The vector space $\mathfrak{h}_{\mathbb{C}}$ has a natural operation of complex conjugation $(\alpha, \alpha') \mapsto (\bar{\alpha}', \bar{\alpha})$, let $\mathfrak{h}_{\mathbb{R}}$ be the subspace $(\alpha, \bar{\alpha})$ of fixed points of this complex conjugation. There is a natural extension of the projection $\mathfrak{h}^{1,0} \rightarrow H_{\mathbb{C}}^B$ to $\mathfrak{h}_{\mathbb{C}} \rightarrow H_{\mathbb{C}}^B$. This mapping is compatible with the complex conjugation, thus induces a mapping $\mathfrak{h}_{\mathbb{R}} \rightarrow H_{\mathbb{R}}^B$. Let $\mathfrak{h}_{\mathbb{C}}^A$ be the kernel of the mapping $\mathfrak{h}_{\mathbb{C}} \rightarrow H_{\mathbb{C}}^B$, similarly for $\mathfrak{h}_{\mathbb{R}}^A$.

Let $\bar{\Omega}_1: H_{\mathbb{C}}^A \rightarrow \mathfrak{h}^{0,1}$ be the complex conjugate to the mapping $\Omega_1: H_{\mathbb{C}}^A \rightarrow \mathfrak{h}^{1,0}$, $\bar{\Omega}_1 v = \overline{\Omega_1 v}$. Define a pre-Hilbert structure on $H_{\mathbb{C}}^A$ via $\|v\|_A^2 = \|\Omega_1 v\|_{\mathfrak{h}^{1,0}}^2$. Then Ω extends to a mapping from Hilbert completion $(H_{\mathbb{C}}^A)^{\text{compl}}$ of $H_{\mathbb{C}}^A$ to $\mathfrak{h}^{1,0}$, same for $\bar{\Omega}_1$ and $\mathfrak{h}^{0,1}$. Let $\mathfrak{h}_{\mathbb{C}}^B$ be the image of this completion w.r.t. $\Omega_1 - \bar{\Omega}_1$. Identify $\mathfrak{h}_{\mathbb{C}}^B$ with a subspace of $H_{\mathbb{C}}^B$ via the projection $\mathfrak{h}_{\mathbb{C}} \rightarrow H_{\mathbb{C}}^B$.

The restriction on Ω immediately imply

Proposition 9.35. $\mathfrak{h}_{\mathbb{C}}$ carries a natural symplectic structure, the corresponding mapping $\mathfrak{h}_{\mathbb{C}} \rightarrow \mathfrak{h}_{\mathbb{C}}^*$ is invertible, and $\mathfrak{h}_{\mathbb{C}}$ is a direct sum of Lagrangian subspaces $\mathfrak{h}_{\mathbb{C}}^A$ and $\mathfrak{h}_{\mathbb{C}}^B$.

Since the subspace $\mathfrak{h}_{\mathbb{C}}^B$ is stable w.r.t. complex conjugation, one can define $\mathfrak{h}_{\mathbb{R}}^B$. Similarly, define $\mathfrak{h}_{\mathbb{Z}}^A$ as the image of $H_{\mathbb{Z}}^A$ in $\mathfrak{h}_{\mathbb{C}}^A$. Note that $\mathfrak{h}_{\mathbb{C}}^A$ and $\mathfrak{h}_{\mathbb{C}}^B$ are mutually dual, so $\mathfrak{h}_{\mathbb{Z}}^A$ is identified with a lattice in $\mathfrak{h}_{\mathbb{C}}^B$. Say that $v \in \mathfrak{h}_{\mathbb{Z}}^B \subset \mathfrak{h}_{\mathbb{C}}^B$ if $(v, w) = 0$ for any $w \in \mathfrak{h}_{\mathbb{Z}}^A$.

Proposition 9.36. The subset $\mathfrak{h}_{\mathbb{Z}}^B$ generates $\mathfrak{h}_{\mathbb{C}}^B$.

Proof. Let $\mathbf{e}_j$, $j \in I_+$, be the natural basis of $H_{\mathbb{Z}}^A$. Let V_j be a subspace of $\mathfrak{h}_{\mathbb{C}}^A$ generated by $\mathbf{e}_k$, $k \neq j$. If $V_j = \mathfrak{h}_{\mathbb{C}}^A$, then $\mathbf{e}_j$ is a linear combination of $\mathbf{e}_k$, $k \neq j$, which contradicts $\text{Im } \Omega \gg 0$. Appropriate multiple of a normal vector to V_i is in $\mathfrak{h}_{\mathbb{Z}}^B$. It is obvious that these elements generate $\mathfrak{h}_{\mathbb{C}}^B$. $\square$

Definition 9.37. Say that the rigged Hopf structure is *integer* if $\mathfrak{h}_{\mathbb{Z}}^A$ is closed in $\mathfrak{h}_{\mathbb{C}}^A$.

The condition of being integer is a restriction from below on $\text{Im } \Omega$, say, it prohibits $\text{Im } \Omega = \text{diag } (\lambda_i)$, $\lambda_i \in l_2$.

We obtained 4 subspaces $\mathfrak{h}_{\mathbb{C}}^A$, $\mathfrak{h}_{\mathbb{C}}^B$, $\mathfrak{h}^{1,0}$ and $\mathfrak{h}^{0,1}$ of $\mathfrak{h}_{\mathbb{C}}$. Note that $\mathfrak{h}_{\mathbb{R}}^A$, $\mathfrak{h}_{\mathbb{R}}^B$ are generated by $\mathbb{Z}$ -lattices $\mathfrak{h}_{\mathbb{Z}}^A$, $\mathfrak{h}_{\mathbb{Z}}^B$ in them. If we consider $\mathfrak{h}_{\mathbb{C}}$ and $\mathfrak{h}^{0,1}$ as real vector spaces, then $\mathfrak{h}_{\mathbb{C}} = \mathfrak{h}_{\mathbb{R}} \oplus \mathfrak{h}^{0,1}$. Note that $\mathfrak{h}^{1,0}$ is naturally identified with $\mathfrak{h}_{\mathbb{C}}/\mathfrak{h}^{0,1}$, thus $\mathfrak{h}^{1,0} \simeq \mathfrak{h}_{\mathbb{R}}$. Let $\mathfrak{L}$ be the image of $\mathfrak{h}_{\mathbb{Z}}^A \oplus \mathfrak{h}_{\mathbb{Z}}^B \subset \mathfrak{h}_{\mathbb{R}} \subset \mathfrak{h}_{\mathbb{C}}$ in $\mathfrak{h}_{\mathbb{C}}/\mathfrak{h}^{0,1}$, identify $\mathfrak{L}$ with a subgroup of $\mathfrak{h}^{1,0}$. It is clear that $\mathfrak{L}$ is a lattice in $\mathfrak{h}^{1,0} \simeq \mathfrak{h}_{\mathbb{R}}$.

Start from these 4 subspaces of $\mathfrak{h}_{\mathbb{C}}$, the complex conjugation and symplectic structure on $\mathfrak{h}_{\mathbb{C}}$, and lattices in the real parts of the first two spaces. Try to reconstruct the initial rigged Hodge structure. Suppose that $\mathfrak{h}_{\mathbb{C}}^A$, $\mathfrak{h}_{\mathbb{C}}^B$ are stable w.r.t. the complex conjugation, and that $\mathfrak{h}^{1,0}$, $\mathfrak{h}^{0,1}$ are interchanged by complex conjugation. Suppose that the last two subspaces project isomorphically on any subspace of the first two (along the other one). Thus $\mathfrak{h}^{1,0}$ and $\mathfrak{h}^{0,1}$ are graphs of invertible mappings $\Omega_1, \bar{\Omega}_1: \mathfrak{h}_{\mathbb{C}}^A \rightarrow \mathfrak{h}_{\mathbb{C}}^B$, these mappings are mutually complex conjugate.

Now the symplectic structure on $\mathfrak{h}_{\mathbb{C}}$ identifies $\mathfrak{h}_{\mathbb{Z}}^B$ with a subset of the dual lattice to $\mathfrak{h}_{\mathbb{Z}}^A$. This reconstructs the groups $H_{\mathbb{Z}}^A$, $H_{\mathbb{Z}}^B$, together with Ω_1 they allow to reconstruct the mapping Ω from $H_{\mathbb{C}}^A = H_{\mathbb{Z}}^A \otimes \mathbb{C}$ to $H_{\mathbb{C}}^B = H_{\mathbb{Z}}^B \otimes \mathbb{C}$. The positivity of $\text{Im } \Omega$ is translated into positivity of $(\alpha, \bar{\alpha})$, $\alpha \in \mathfrak{h}^{1,0}$, and the condition $\mathfrak{h}^{1,0} \oplus \mathfrak{h}^{0,1} = \mathfrak{h}$.

Now the results of this chapter imply

Proposition 9.38. Consider a family of disks $K_i \subset \mathbb{CP}^1$, $i \in I$, with pairwise conformal distances l_{ij} . Let $\mathcal{R} = (e^{-l_{ij}} - \delta_{ij})$. Suppose that $\mathcal{R}$ gives a compact mapping $l_2 \rightarrow l_2$ and a bounded mapping $l_{\infty} \rightarrow l_{\infty}$, and that some power of $\mathcal{R}$ gives a bounded mapping $l_{\infty} \rightarrow l_2$. Consider an involution $' : I \rightarrow I$, $I = I_+ \amalg I_+'$, glue boundaries of disks pairwise using fraction-linear mappings $\varphi_i = \varphi_{i'}^{-1}$, let (Ω_{jk}) , $j, k \in I_+$, be the matrix of periods of the resulting curve $\bar{M}$. Then Ω_{jk} gives rise to a rigged Hodge structure.

Note that it is natural to call the space $\mathfrak{h}_{\mathbb{C}}$ of this rigged Hodge structure *the first cohomology space* of the curve $\bar{M}$. The space $\mathfrak{h}^{1,0}$ can be identified with $\Gamma(\bar{M}, \omega)$. As in finite-genus case, the quotient $\mathfrak{h}^{1,0}/\mathcal{L}$ is identified with the (bounded) Jacobian of the curve.

REFERENCES

1. D. Bättig, A. M. Bloch, J.-C. Guillot, and T. Kappeler, *On the symplectic structure of the phase space for periodic KdV, Toda, and defocusing NLS*, Duke Math. J. **79** (1995), no. 3, 549–604.
2. A. A. Beĭlinson and Yu. I. Manin, *The Mumford form and the Polyakov measure in string theory*, Comm. Math. Phys. **107** (1986), no. 3, 359–376.
3. J. Feldman, H. Knörrer, and Trubowitz E., *Infinite genus riemann surfaces*, Canadian Mathematical Society 1945-1995 (James B. Carrell and Ram Murty, eds.), no. 3, Canadian Mathematical Society, Ottawa, 1996, pp. 91–112.
4. I. M. Gelfand and G. E. Shilov, *Generalized functions and operations*, Generalized functions, vol. 1, Nauka, Moscow, 1958.
5. I. M. Gelfand and N. Ya. Vilenkin, *Generalized functions. Vol. 4*, Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1964 [1977], Applications of harmonic analysis, Translated from the Russian by Amiel Feinstein.
6. Lars Hörmander, *The analysis of linear partial differential operators. I*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 256, Springer-Verlag, Berlin, 1983, Distribution theory and Fourier analysis.
7. H. P. McKean and C. Trubowitz, *Hill's operator and hyperelliptic function theory in the presence of infinitely many branch points*, Comm. Pure Appl. Math. **29** (1976), no. 1, 143–226.
8. Takahiro Shiota, *Characterization of Jacobian varieties in terms of soliton equations*, Inventiones Mathematicae **83** (1986), 333–382.
9. Ilya Zakharevich, *The direct and inverse scattering transforms for a Sturm - Liouville equation in finite interval as smooth maps. The geometry of a set of finite gap potentials*, Inverse Methods in Action (P. C. Sabatier, ed.), Springer, 1990, p. 246.
10. ———, *The spectral problems for a Sturm - Liouville equation with a parameter. The geometry of a set of finite gap potentials*, Soviet Mathematics—Doklady **43** (1991), no. 2, 439–444.

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, 231 W. 18 AVE, COLUMBUS, OH, 43210

E-mail address: ilya@math.ohio-state.edu